



Partitioning Infinite Hypergraphs into Few Monochromatic Berge-Paths

Sebastián Bustamante¹ · Jan Corsten² · Nóra Frankl²

Received: 13 May 2019 / Revised: 13 May 2019

© The Author(s) 2020

Abstract

Extending a result of Rado to hypergraphs, we prove that for all $s, k, t \in \mathbb{N}$ with $k \geq t \geq 2$, the vertices of every $r = s(k - t + 1)$ -edge-coloured countably infinite complete k -graph can be partitioned into the cores of at most s monochromatic t -tight Berge-paths of different colours. We further describe a construction showing that this result is best possible.

Keywords Graph partitioning · Monochromatic cycle partitioning · Infinite graphs · Berge-paths

1 Introduction

Lehel's conjecture (first seen in [2]) states that the vertices of every 2-edge-coloured complete graph can be partitioned into two monochromatic cycles, one of each colour. Here, single vertices and edges are considered to be cycles and this convention is used throughout this paper. The conjecture was proved for very large graphs by Łuczak et al. [14] in 1998, for large graphs by Allen [1] in 2008, and for all graphs by Bessy and Thomassé [3] in 2010.

Erdős et al. [6] conjectured in 1991 that this can be extended to r colours (allowing r monochromatic cycles). This was however disproved by Pokrovskiy [17], who showed that for every $r \geq 3$, there are infinitely many r -edge-coloured complete graphs whose vertices cannot be covered by r monochromatic vertex-disjoint cycles. Finding the minimum number of monochromatic vertex-disjoint cycles needed to cover the vertices of any r -edge-coloured complete graph remains a big open problem. We note

✉ Nóra Frankl
n.frankl@lse.ac.uk

Jan Corsten
j.corsten@lse.ac.uk

¹ Universidad de Chile, Santiago, Chile

² Department of Mathematics, London School of Economics, Houghton St, London WC2A 2AE, UK

that a priori it is not obvious that this minimum is independent of the size of the complete graph we wish to cover. The fact that this is the case was proved by Erdős et al. [6], who also presented a simple construction in which r cycles are needed. The currently best-known upper bound is $100r \log r$, due to Gyárfás et al. [9].

An infinite analogue of the conjecture of Erdős et al. is true as proved by Rado [18] already in 1978.

Theorem 1.1 ([18]) *The vertices of every countably infinite r -edge-coloured complete graph can be partitioned into r monochromatic paths (infinite or finite), one of each colour.*

Rado’s theorem is best possible, as the construction for finite graphs in [6] can be extended to infinite graphs.

In this note we consider extensions of this result to hypergraphs. A k -uniform hypergraph (or shortly, a k -graph) is a tuple $\mathcal{H} = (V, \mathcal{E})$ where V and \mathcal{E} are sets with $\mathcal{E} \subseteq \binom{V}{k}$. The complete k -graph with vertex-set V , denoted by $\mathcal{K}_V^{(k)}$, is the k -graph with edge-set $\mathcal{E} = \binom{V}{k}$.

There are different notions of paths in hypergraphs, *loose paths*, *tight paths* and *Berge-paths*; all of these coincide with the notion of paths when $k = 2$. In this note, we will mainly consider Berge-paths. Given integers $2 \leq t \leq k$ and $\ell \geq 1$, a finite (k -uniform) t -tight Berge-path of length ℓ is a pair (X, F) defined as follows. $X = \{v_1, \dots, v_{\ell+t-1}\} \subseteq V$ is a set of $\ell+t-1$ distinct vertices, $F = \{e_1, \dots, e_\ell\} \subseteq \mathcal{E}$ is a set of ℓ distinct edges and $v_i, v_{i+1}, \dots, v_{i+t-1} \in e_i$ for all $i \in [\ell]$. For technical reasons we define a t -tight Berge-path of length 0 as a pair $(\{v\}, \emptyset)$. A (one-way) infinite t -tight Berge-path is a pair (X, F) where $X = \{v_i : i \in \mathbb{N}\} \subseteq V$, $F = \{e_i : i \in \mathbb{N}\} \subseteq \mathcal{E}$, and $v_i, v_{i+1}, \dots, v_{i+t-1} \in e_i$ for all $i \in \mathbb{N}$. For a t -tight Berge-path $P = (X, F)$, X is called the *core* of P . A family $P_1 = (X_1, F_1), \dots, P_r = (X_r, F_r)$ of t -tight Berge-paths *core-partitions* V if $X_1 \cup \dots \cup X_r$ is a partition of V . Given an edge-colouring φ of a k -graph \mathcal{H} , a t -tight Berge-path $P = (X, F)$ in \mathcal{H} is said to be *monochromatic* in colour c if $\varphi(f) = c$ for all $f \in F$.

A 2-tight Berge-path is simply called a Berge-path and a k -tight Berge-path is called a tight path. The k -uniform *loose path* of length ℓ consists of $n = k(\ell - 1) + 1$ vertices $\{v_1, \dots, v_n\}$ and the ℓ edges $\{v_{i(k-1)+1}, \dots, v_{i(k-1)+k}\}$ for $i = 0, \dots, \ell - 1$. The *infinite loose path* consists of the vertices $\{v_1, v_2, \dots\}$ and edges $\{v_{i(k-1)+1}, \dots, v_{i(k-1)+k}\}$ for all $i = 0, 1, \dots$

Many extensions of path partition problems to hypergraphs have been studied, considering loose paths [10,11,20], tight paths [4,5] and Berge-paths [10]. Most relevant for the topic of this note are the following two extensions of Theorem 1.1.

Theorem 1.2 (Gyárfás–Sárközy [10]) *The vertices of every countably infinite r -edge-coloured complete k -graph can be partitioned into r monochromatic loose paths (infinite or finite), one of each colour.*

Theorem 1.3 (Elekes–Soukup–Soukup–Szentmiklóssy [5]) *The vertices of every countably infinite r -edge-coloured complete k -graph can be partitioned into r monochromatic tight paths (infinite or finite), one of each colour.*

The latter result answers a question of Gyárfás and Sárközy from [10]. Note that both Theorems 1.2 and 1.3 reduce to Theorem 1.1 when $k = 2$. Our main result extends Theorem 1.1 in a similar way to Berge-paths. It turns out that $\lceil r/(k - 1) \rceil$ paths suffice.

Theorem 1.4 *For all $s, k \in \mathbb{N}$ with $k \geq 2$ and every $r = s(k - 1)$ -edge-colouring of $\mathcal{K}_{\mathbb{N}}^{(k)}$, the vertices can be core-partitioned into s monochromatic Berge-paths of different colours.*

Note that Theorem 1.4 reduces to Theorem 1.1 when $k = 2$ as well. We shall actually prove the following more general result about t -tight Berge-paths.

Theorem 1.5 *For all $s, k, t \in \mathbb{N}$ with $k \geq t \geq 2$ and every $r = s(k - t + 1)$ -edge-colouring of $\mathcal{K}_{\mathbb{N}}^{(k)}$, the vertices can be core-partitioned into s monochromatic t -tight Berge-paths of different colours.*

Note that the case $k = 2$ reduces to Theorem 1.4, and the case $k = t$ reduces to Theorem 1.3. The following theorem shows that Theorem 1.5 is best possible.

Theorem 1.6 *For all $s, k, t \in \mathbb{N}$ with $k \geq t \geq 2$, there is an edge-colouring of $\mathcal{K}_{\mathbb{N}}^{(k)}$ with $r = s(k - t + 1) + 1$ colours in which the vertices cannot be covered by the cores of s monochromatic t -tight Berge-paths.*

We will prove Theorem 1.6 in Sect. 2 and Theorem 1.5 in Sect. 3.

2 Proof of Theorem 1.6

The construction described in the proof generalises the construction from [6].

Proof of Theorem 1.6 We denote the lexicographical ordering of $\binom{[r]}{s}$ by \prec . First, we partition \mathbb{N} into sets $\{B_I : I \in \binom{[r]}{s}\}$ so that all B_I 's but $B_{\{r-s+1, \dots, r\}}$ are finite and $|B_I| \geq st \cdot \sum_{J \prec I} (|B_J| + 1)$ for every $I \in \binom{[r]}{s}$.¹ For $x \in \mathbb{N}$, let $I(x)$ be the s -subset of $[r]$ for which $x \in B_{I(x)}$. We define an r -edge-colouring φ of $\mathcal{K}_{\mathbb{N}}^{(k)}$ as follows.

For every $e \in E(\mathcal{K}_{\mathbb{N}}^{(k)})$ we consider an order x_e^1, \dots, x_e^k of e satisfying $I(x_e^i) \preceq I(x_e^j)$ for all $1 \leq i < j \leq k$, and define $\varphi(e)$ as an arbitrary member of $[r] \setminus \bigcup_{i \leq k-t+1} I(x_e^i)$.

Assume for contradiction that there are monochromatic t -tight Berge-paths P_1, \dots, P_s with cores X_1, \dots, X_s so that $\bigcup_i X_i = \mathbb{N}$ and let $C \subseteq [r]$ be a set of size s which contains all colours used by these t -tight Berge-paths.

Observe that for every edge e with $e \cap B_C \neq \emptyset$ and $\varphi(e) \in C$ we have

$$\left| e \cap \bigcup_{J \prec C} B_J \right| \geq k - t + 1. \tag{2.1}$$

Indeed, if $|e \cap \bigcup_{J \prec C} B_J| < k - t + 1$, then $C \preceq I(x_e^{k-t+1})$ and thus $\varphi(e) \notin C$.

¹ This growth rate of the $|B_I|$ can be improved by a more careful analysis.

For $i \in C$, let F_i be the set of all $f \in \binom{X_i}{t}$ which consist of t consecutive vertices of X_i with at least one element in B_C . Let $f \in F_i$ and let $e \in E(P_i)$ be some edge with $f \subseteq e$. By (2.1), we have $|e \setminus \bigcup_{J < C} B_J| \leq t - 1$ and therefore some vertex in f must be in $\bigcup_{J < C} B_J$. Since every $v \in \mathbb{N}$ is contained in at most t sets $f \in F_i$, it follows that

$$|F_i| \leq t \left| \bigcup_{J < C} B_J \right|. \tag{2.2}$$

Observe now that for all but at most $t - 1$ vertices $v \in X_i \cap B_C$, there is a unique $f \in F_i$ starting at v and thus

$$|X_i \cap B_C| \leq |F_i| + t - 1. \tag{2.3}$$

Combining (2.2) and (2.3), we get

$$|X_i \cap B_C| \leq t \cdot \sum_{J < C} |B_J| + t - 1 < |B_C|/r$$

for every $i \in [r]$ and hence $|B_C| = |B_C \cap (\bigcup_i X_i)| < |B_C|$, a contradiction. \square

3 Proof of Theorem 1.5

Our proof is based on ideas from [5]. First, we need to introduce some notation. An r -multi-colouring of a k -graph G is a function $\chi : E(G) \rightarrow 2^{[r]}$. Given a set $F \subseteq E(G)$, we denote by $\chi(F) = \bigcap_{e \in F} \chi(e)$ the set of colours they have in common and say that F is (χ) -monochromatic if $\chi(F)$ is non-empty. For a given r -colouring φ of $\mathcal{K} := \mathcal{K}_{\mathbb{N}}^{(k)}$ and $i, j \in \mathbb{N}$ with $j < k$, we define an r -multi-colouring $\varphi_{i,j} : \binom{\mathbb{N} \setminus \{i\}}{j} \rightarrow 2^{[r]}$ by

$$\varphi_{i,j}(f) = \{\varphi(e) : e \in E(\mathcal{K}) \text{ and } \{i\} \cup f \subseteq e\}.$$

Furthermore, we call $\{\mathcal{K}_i : i \in \mathbb{N}\}$ a j -clique-chain w.r.t. an r -colouring φ of \mathcal{K} if \mathcal{K}_1 is a $\varphi_{1,j}$ -monochromatic copy of $\mathcal{K}_{\mathbb{N}}^{(j)}$ with $V(\mathcal{K}_1) \subseteq \mathbb{N}$ and \mathcal{K}_i is a $\varphi_{i,j}$ -monochromatic copy of $\mathcal{K}_{\mathbb{N}}^{(j)}$ with $V(\mathcal{K}_i) \subseteq V(\mathcal{K}_{i-1})$ for every $i \in \mathbb{N}$.

Observe that, by Ramsey’s theorem [19] for infinite hypergraphs, there exists a j -clique-chain for every r -colouring of \mathcal{K} and every $j \in [k - 1]$.

For a j -clique-chain $\{\mathcal{K}_i : i \in \mathbb{N}\}$ we define a vertex-multi-colouring $\chi : \mathbb{N} \rightarrow 2^{[r]}$ by $\chi(i) = \bigcap_{e \in E(\mathcal{K}_i)} \varphi_{i,j}(e)$ for every $i \in \mathbb{N}$. We call χ the *clique-colouring induced by* $\{\mathcal{K}_i : i \in \mathbb{N}\}$.

Lemma 3.1 *For all $s, k, t \in \mathbb{N}$ with $k \geq t \geq 2$ and every $r = s(k - t + 1)$ -colouring of $\mathcal{K}_{\mathbb{N}}^{(k)}$, there is a $(t - 1)$ -clique-chain that induces a clique-colouring using at most s colours.*

Proof Let φ be the given r -colouring of $\mathcal{K} := \mathcal{K}_{\mathbb{N}}^{(k)}$. Furthermore, let $C_1 \cup \dots \cup C_r$ be a partition of the set of $r = s(k - t + 1)$ colours into r blocks of size s . We will

show that there is a $(t - 1)$ -clique-chain and some $i \in [r]$ such that for the induced clique-colouring we have $\chi(v) \cap C_i \neq \emptyset$ for all $v \in \mathbb{N}$.

We call an infinite $(t - 1)$ -uniform clique \mathcal{K}' *maximally-monochromatic* w.r.t. a multi-colouring ψ of \mathcal{K}' and a set $C \subseteq [r]$ if there is no infinite clique $\mathcal{K}'' \subseteq \mathcal{K}'$ with $|\{i : \psi(\mathcal{K}'') \cap C_i \neq \emptyset\} \cap C| > |\{i : \psi(\mathcal{K}') \cap C_i \neq \emptyset\} \cap C|$. Note that a maximally-monochromatic clique is not necessarily monochromatic (since all its infinite monochromatic subcliques might have colours not in C). Further note that every infinite clique contains a maximally-monochromatic infinite clique (since r is finite).

We build a $(t - 1)$ -clique-chain as follows. Let \mathcal{K}_1 be any $\varphi_{1,t-1}$ -monochromatic, maximally-monochromatic $(t - 1)$ -uniform clique w.r.t. $\varphi_{1,t-1}$ and $D = [r]$, and let $D_1 := \{i : \varphi_{1,t-1}(\mathcal{K}_1) \cap C_i \neq \emptyset\}$. Now, for every $j \in \mathbb{N}$, let \mathcal{K}_{j+1} be a $\varphi_{j+1,t-1}$ -monochromatic, maximally-monochromatic clique w.r.t. $\varphi_{j+1,t-1}$ and D_j with $V(\mathcal{K}_{j+1}) \subseteq V(\mathcal{K}_j)$ and let $D_{j+1} = \{i : \varphi_{j+1,t-1}(\mathcal{K}_{j+1}) \cap C_i \neq \emptyset\}$. If there is some $i \in [r]$ such that $C_i \cap D_j \neq \emptyset$ for all $j \in \mathbb{N}$, then $\{\mathcal{K}_1, \mathcal{K}_2, \dots\}$ is a $(t - 1)$ -clique-chain with the desired property. Hence we may assume that there is no such i .

Thus, there exist j_1, \dots, j_r , such that $C_i \cap D_{j_i} = \emptyset$ but $C_i \cap D_{j_i-1} \neq \emptyset$ for every $i \in [r]$. Without loss of generality we may assume that $j_1 \leq \dots \leq j_r$. Let $X = V(\mathcal{K}_{j_r})$ and note that $V(\mathcal{K}_{j_i}) \supseteq X$ for every $i \in [r]$. Define $\Phi : \binom{X}{t-1} \rightarrow 2^{[r]}$ by

$$\Phi(f) = \{\varphi(e) : e \in E(\mathcal{K}) \text{ and } \{j_1, \dots, j_r\} \cup f \subseteq e\}.$$

Note that every $f \in \binom{X}{t-1}$ receives at least one colour, and that $\Phi(f) \subseteq \varphi_{j_i,t-1}(f)$ for every $f \in \binom{X}{t-1}$ and every $i \in [r]$. By Ramsey's theorem for hypergraphs there is a Φ -monochromatic infinite clique \mathcal{K}' in X . Therefore, there is some $\ell \in [r]$ such that $\Phi(\mathcal{K}') \cap C_\ell \neq \emptyset$ and consequently \mathcal{K}_{j_ℓ} is not maximally monochromatic. \square

We proceed now with the proof of Theorem 1.5.

Proof of Theorem 1.5 Let φ be the given r -colouring of $\mathcal{K} = \mathcal{K}_{\mathbb{N}}^{(k)}$. By Lemma 3.1, there is a $(t - 1)$ -clique-chain $\{\mathcal{K}_i : i \in \mathbb{N}\}$ that induces a clique-colouring χ using at most s colours (without loss of generality these colours are $1, \dots, s$). For $i \in [s]$, let $A_i \subseteq \mathbb{N}$ be the set of vertices of colour i according to χ .

By repeating the following process we will simultaneously build t -tight monochromatic Berge-paths P_1, \dots, P_r with core-vertex sequences $\{b_{i,1}, b_{i,2}, \dots\}$ for every $i \in [s]$. Let $b_{i,1} := \min A_i$ for every $i \in [s]$. In every step, we will add to each path t or $t - 1$ vertices making sure that for every $i \in [s]$, the last new vertex, say b_{i,n_i} , is in A_i , and that the other new vertices are in $V(\mathcal{K}_{b_{i,n_i}})$. Right after choosing the vertex $b_{i,j}$, we will choose a unique edge $e_{i,j} \in E(\mathcal{K})$ of colour i which contains the t consecutive vertices $b_{i,j-t+1}, \dots, b_{i,j}$ for every $j \geq t$ and $i \in [s]$. Let $X = \{b_{1,1}, \dots, b_{s,1}\}$ and let $Y = \emptyset$. We will use X to keep track of already used vertices and Y to keep track of already used edges.

For each $i \in [s]$ do the following.² Suppose the current path P_i ends in $b_{i,n} \in A_i$ for some $n \in \mathbb{N}$. We will now extend P_i by t or $t - 1$ vertices as follows. Let a be the

² To avoid unnecessary subscripts for 'local variables', we treat i as being fixed in the following.

smallest vertex in $A_i \setminus X$ (if $A_i \setminus X$ is empty, the path P_i is complete and we move to the next step). Add a to X and do the following for every $j = 1, \dots, t - 2$. Choose a vertex $b_{i,n+j} \in V(\mathcal{K}_a) \setminus (\bigcup Y)$ and add it to X (note that this is always possible since $V(\mathcal{K}_a)$ is infinite and Y is finite). Let $f_{i,n+j} = \{b_{i,n+j-t+1}, \dots, b_{i,n+j}\}$ be the set of the t consecutive vertices in the core of the Berge-path P_i ending at $b_{i,n+j}$. Note that $f_{i,n+j} \setminus \{b_{i,n}\} \in E(\mathcal{K}_{b_{i,n}})$ and thus $i \in \varphi_{b_{i,n},t-1}(f_{i,n+j} \setminus \{b_{i,n}\})$. Hence, by the definition of $\varphi_{b_{i,n},t-1}$, there exist $e_{i,n+j} \in E(\mathcal{K})$ with $f_{i,n+j} \subseteq e_{i,n+j}$ and $\varphi(e_{i,n+j}) = i$. Add $e_{i,n+j}$ to Y . Since $b_{i,n+j} \notin \bigcup Y$, we have $e_{i,n+j} \not\subseteq Y$ and we can therefore use $e_{i,n+j}$ as the edge for $f_{i,n+j}$ in our Berge-path.

Choosing the next vertex will be slightly more complicated (since a might be in some edge in Y). Let $b \in V(\mathcal{K}_a) \setminus \bigcup Y$ and let $f'_1 = \{b_{i,n}, \dots, b_{i,n+t-2}, b\}$ and $f'_2 = \{b_{i,n+1}, \dots, b_{i,n+t-2}, b, a\}$, and note that $f'_1 \setminus \{b_{i,n}\} \in E(\mathcal{K}_{b_{i,n}})$ and $f'_2 \setminus \{a\} \in E(\mathcal{K}_a)$. As before, $i \in \varphi_{b_{i,n},t-1}(f'_1 \setminus \{b_{i,n}\}) \cap \varphi_{a,t-1}(f'_2 \setminus \{a\})$ and thus there exist $e'_1, e'_2 \in E(\mathcal{K})$ with $f'_s \subseteq e'_s$ and $\varphi(e'_s) = i$ for both $s = 1, 2$. If $e'_1 \neq e'_2$, let $b_{i,n+t-1} := b$ and $b_{i,n+t} = a$ and let $e_{i,n+t-1} := e'_1$ and $e_{i,n+t} = e'_2$. Add $b_{i,n+t-1}$ to X and $e_{i,n+t-1}, e_{i,n+t}$ to Y . Note that $e_{i,n+t-1}$ and $e_{i,n+t}$ can be chosen as the edges for the t consecutive vertices of P_i ending in $b_{i,n+t-1}$ and $b_{i,n+t}$. If $e'_1 = e'_2$, let $b_{i,n+t-1} = a$ and $e_{i,n+t-1} = e'_1 = e'_2$, and add $e_{i,n+t-1}$ to Y . Note that the t consecutive vertices of P_i ending in $b_{i,n+t-1}$ are contained in $e_{i,n+t-1}$ and $e_{i,n+t-1} \not\subseteq Y$. Hence, $e_{i,n+t-1}$ can be chosen as the edge for the t consecutive vertices of P_i ending in $b_{i,n+t-1}$.

By construction, P_1, \dots, P_s are monochromatic t -tight Berge-paths whose cores are disjoint. Furthermore, since at the beginning of every step the smallest uncovered vertex a of A_i is chosen, we have $\bigcup_i V(P_i) = \mathbb{N}$. □

4 Further Remarks and Open Problems

Theorems 1.1–1.4 remain true when we consider cycles instead of paths, where an infinite cycle is a two-way infinite path. It is not clear to us however if one can replace paths by cycles in Theorem 1.5 when $2 < t < k$. Difficulties only arise when trying to close finite paths to cycles, hence we can replace paths by cycles if we allow finitely many vertices to be uncovered.

A natural question to ask is if similar results hold in the finite setting. Gyárfás, Lehel, Sárközy and Schelp [8] conjectured that every finite $(k - 1)$ -edge-coloured complete k -graph contains a monochromatic Hamiltonian Berge-cycle. Note that, in the infinite setting, this is a special case of Theorem 1.4. After partial results in [8,12,13,15], Omidi [16] announced a proof of this conjecture.

We believe that a generalisation of this to more colours, similar as in Theorem 1.4, is true as well.

Conjecture 4.1 *For all $s, k \in \mathbb{N}$ with $k \geq 2$, there is some $c = c(s, k) \in \mathbb{N}$ such that the following is true for all $n \in \mathbb{N}$. In every $r = s(k - 1)$ -edge-colouring of $\mathcal{K}_n^{(k)}$, there is a collection of at most s monochromatic t -tight Berge-cycles whose cores are disjoint and cover all but c vertices.*

For $k = 2$, this reduces to a conjecture of Pokrovskiy [17]. We further believe that this can be extended to t -tight Berge-cycles similarly to Theorem 1.5.

Conjecture 4.2 *For all $s, k, t \in \mathbb{N}$ with $k \geq t \geq 2$, there is some $c = c(s, k, t) \in \mathbb{N}$ such that the following is true for all $n \in \mathbb{N}$. In every $r = s(k - t + 1)$ -edge-colouring of $\mathcal{K}_n^{(k)}$, there is a collection of at most s monochromatic t -tight Berge-cycles whose cores are disjoint and cover all but c vertices.*

A simple modification of the construction in Sect. 2 shows that these conjectures are best possible (if true) apart from the finite leftover.

Recently we learned that Gerbner et al. [7, unpublished] made some progress towards these conjectures.

Acknowledgements The authors would like to thank Peter Allen and Jan van den Heuvel for their helpful suggestions on the manuscript. The first author was supported by CONICYT Doctoral Fellowship 21141116. Part of this research was done while the second and third author visited Universiad de Chile with the support of the Santander Travel Research Fund. The third author was supported by the grant of Russian Government N 075-15-2019-1926.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Allen, P.: Covering two-edge-coloured complete graphs with two disjoint monochromatic cycles. *Comb. Probab. Comput.* **17**(4), 471–486 (2008)
2. Ayl, J.: Sur l'existence de deux cycles supplémentaires unicolores, disjoints et de couleurs différentes dans un graphe complet bicolore, Ph.D. thesis, Université Joseph-Fourier-Grenoble I (1979)
3. Bessy, S., Thomassé, S.: Partitioning a graph into a cycle and an anticycle, a proof of Lehel's conjecture. *J. Comb. Theory Ser. B* **100**(2), 176–180 (2010)
4. Bustamante, S., Corsten, J., Frankl, N., Pokrovskiy, A., Skokan, J.: Partitioning edge-coloured hypergraphs into few monochromatic tight cycles. [arXiv:1903.04471](https://arxiv.org/abs/1903.04471) (2019)
5. Elekes, M., Soukup, D., Soukup, L., Szentmiklóssy, Z.: Decompositions of edge-colored infinite complete graphs into monochromatic paths. *Discrete Math.* **340**(8), 2053–2069 (2017)
6. Erdős, P., Gyárfás, A., Pyber, L.: Vertex coverings by monochromatic cycles and trees. *J. Comb. Theory Ser. B* **51**(1), 90–95 (1991)
7. Gerbner, D., Methuku, A., Omid, G., Vizer, M.: Monochromatic Berge cycles in uniform hypergraphs (2019) (**unpublished manuscript**)
8. Gyárfás, A., Lehel, J., Sárközy, G., Schelp, R.: Monochromatic Hamiltonian Berge-cycles in colored complete uniform hypergraphs. *J. Comb. Theory Ser. B* **98**(2), 342–358 (2008)
9. Gyárfás, A., Ruszinkó, M., Sárközy, G., Szemerédi, E.: An improved bound for the monochromatic cycle partition number. *J. Comb. Theory Ser. B* **96**(6), 855–873 (2006)
10. Gyárfás, A., Sárközy, G.: Monochromatic path and cycle partitions in hypergraphs. *Electron. J. Comb.* **20**(1), 18 (2013)
11. Gyárfás, A., Sárközy, G.: Monochromatic loose-cycle partitions in hypergraphs. *Electron. J. Comb.* **21**(2), 2–36 (2014)
12. Gyárfás, A., Sárközy, G., Szemerédi, E.: Long monochromatic Berge cycles in colored 4-uniform hypergraphs. *Graphs Comb.* **26**(1), 71–76 (2010)
13. Gyárfás, A., Sárközy, G., Szemerédi, E.: Monochromatic matchings in the shadow graph of almost complete hypergraphs. *Ann. Comb.* **14**(2), 245–249 (2010)

14. Łuczak, T., Rödl, V., Szemerédi, E.: Partitioning two-coloured complete graphs into two monochromatic cycles. *Comb. Probab. Comput.* **7**(4), 423–436 (1998)
15. Maherani, L., Omid, G.R.: Monochromatic Hamiltonian Berge-cycles in colored hypergraphs. *Discrete Math.* **340**(8), 2043–2052 (2017)
16. Omid, G.R.: A proof for a conjecture of Gyárfás, Lehel, Sárközy and Schelp on Berge-cycles. [arXiv:1404.3385](https://arxiv.org/abs/1404.3385) (2014)
17. Pokrovskiy, A.: Partitioning edge-coloured complete graphs into monochromatic cycles and paths. *J. Comb. Theory Ser. B* **106**, 70–97 (2014)
18. Rado, R.: Monochromatic paths in graphs. *Ann. Discrete Math.* **3**, 191–194 (1978)
19. Ramsey, F.P.: On a problem of formal logic. *Proc. Lond. Math. Soc.* **2**(1), 264–286 (1930)
20. Sárközy, G.: Improved monochromatic loose cycle partitions in hypergraphs. *Discrete Math.* **334**, 52–62 (2014)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.