

A maximum likelihood approach to combining forecasts

GILAT LEVY

Department of Economics, London School of Economics

RONNY RAZIN

Department of Economics, London School of Economics

We model an individual who wants to learn about a state of the world. The individual has a prior belief and has data that consist of multiple forecasts about the state of the world. Our key assumption is that the decision maker identifies explanations that could have generated this data and among these focuses on those that maximize the likelihood of observing the data. The decision maker then bases her final prediction about the state on one of these maximum likelihood explanations. We show that in all the maximum likelihood explanations, moderate forecasts are just statistical derivatives of extreme ones. Therefore, the decision maker will base her final prediction only on the information conveyed in the relatively extreme forecasts. We show that this approach to combining forecasts leads to a unique prediction, and a simple and dynamically consistent way to aggregate opinions.

KEYWORDS. Maximum likelihood, combining forecasts, misspecified models.

JEL CLASSIFICATION. D8.

1. INTRODUCTION

In many economic and political situations we find ourselves confronted with multiple opinions or forecasts about variables that are important for decision making. In financial markets, we are often exposed to multiple forecasts about particular stocks or investment possibilities, when we buy a new laptop, we might read multiple reviews either in news outlets or on social media platforms, and prior to election day, we are exposed to multiple polls or to opinions espoused by friends and colleagues. These different pieces of information might be important for our decision making about investments, what we buy, or who we vote for.

To aggregate forecasts, individuals need to take a view about how the forecasts are generated and how they are related to one another. Naturally, multiple forecasts may be correlated. This can arise when we aggregate advice from friends in a connected network, when election polling firms rely on the same data (as is often the case), or when forecasters in financial markets use similar data sources. Indeed, in the last two

Gilat Levy: g.levy1@lse.ac.uk

Ronny Razin: r.razin@lse.ac.uk

We thank seminar participants at Northwestern, ESSET 2018, and Bonn, and two anonymous referees, as well as Xitong Hui for her research assistance. This project received funding from the European Union's Horizon 2020 research and innovation programme under Grant 681579.

decades, online communication has introduced a more complicated web of information sources with potentially higher degrees of correlation across such forecasts. This complicates the problem of combining forecasts and puts into question traditional models that simply assume that individuals have a prior distribution about all relevant information structures generating these forecasts. One possible alternative assumption explored in the literature is that individuals use a simple heuristic, e.g., assume that forecasts are (conditionally) independent and aggregate them accordingly.¹

In this paper, we take a novel approach to this problem: We assume that individuals aggregate forecasts by looking for the most likely explanation for what they observe. That is, we consider a maximum likelihood theory of combining forecasts.² We model an individual who wants to learn about a state of the world (finite set of states). The individual has a prior belief and observes multiple forecasts about the state. These forecasts arrive sequentially, one in each period. We assume that the individual believes that each forecast was generated by a Bayesian forecaster but has no prior beliefs about the joint information structure generating the set of forecasts. Our key assumption is that she identifies joint information structures that generated the data with the highest likelihood and bases her final prediction on such explanations.

In our main result, we characterize the set of maximum likelihood (ML) explanations and ML predictions that the individual adopts. We show that the ML prediction is unique and depends only on a small set of forecasts, not more than the number of states. In particular, the individual who uses a maximum likelihood explanation to combine forecasts ignores forecasts that are relatively moderate and focuses attention only on a set of extreme forecasts.

The intuition for this result follows from two simple observations. First, to increase the likelihood of her explanation, the individual “looks” for correlation between the sources of information that she thinks generated the data. Correlated data sources are more likely to have generated a specific set of forecasts rather than independent sources. This implies that when interpreting the data through the lens of her explanation, she focuses attention on a small number of forecasts while ignoring others.

The second observation is that explaining extreme forecasts that are more distant from the prior involves a lower maximum likelihood. To give an extreme example, consider the highest likelihood of observing a forecast that puts a probability close to 1 that the state is high. To explain this forecast, we need to assume that the signals that generated it were very informative; thus, the likelihood of observing this forecast is close to the prior probability that the state is indeed high. Alternatively, to explain a forecast that is close to the prior, one can assume an uninformative signal; an uninformative signal can generate a posterior that is close to the prior with a probability close to 1. Therefore,

¹See, for example, De Marzo et al. (2003), Glaeser and Sunstein (2009), Levy and Razin (2015, 2018), and Golub and Jackson (2012). Ellis and Piccione (2017) and Levy and Razin (forthcoming) consider decision makers who misperceive correlation. Spiegel (2016) analyzes a model in which individuals misspecify the true causal Bayesian network. Arieli et al. (2019) use scoring rules and regret to evaluate forecast aggregation schemes.

²Such a procedure is in line with legal reasoning where plausible argumentation and balance of probabilities are suggested as normative approaches to the aggregation of information.

explaining the extreme views puts an upper bound on the likelihood of any explanation of the entire data. As we show, the individual can achieve this bound. She explains extreme views and treats moderate forecasts as noisy derivatives of extreme forecasts.

The above result has several behavioral implications for the dynamic evolution of the ML prediction. First, ML predictions can exhibit stagnation. When more moderate forecasts are added to her data set, the decision maker does not change her prediction. Second, the prior plays an important role; specifically, the notion of extreme views is measured with respect to the prior. As we show, in the long term, limit predictions can depend on the prior even when individuals are exposed to rich and large amounts of information.

In our model, the decision maker's misspecified view or narrative of the world is constantly changing with new observations. But does she have to change her "world view" drastically each time she observes a new forecast? We formulate a notion of time consistency across explanations under which older explanations are never discarded but continue to play a part in the new narrative the decision maker adopts. We show that the decision maker can always construct her ML explanations to be time-consistent. Intuitively, new forecasts are added to the previous explanation either by being correlated to previous extreme forecasts or the other way around. Thus, the decision maker's world view evolves to accommodate new forecasts without altering how previous ones are explained. Our formulation of time-consistency provides a novel normative criterion for such evolving misspecified models of the world.

A growing literature in economics studies how individuals combine forecasts in a non-Bayesian manner, with many papers focusing on correlation neglect. In contrast, in our analysis, decision makers entertain all possible correlation structures: The focus on maximizing the likelihood implies that decision makers actively "look" for correlation.³ While the correlation neglect assumption implies that individuals' beliefs become more extreme, in our model, a different form of "extremism" arises, as individuals base their prediction only on relatively extreme forecasts. Our result complements other explanations for this form of extremism; for example, several papers show how extreme views are more likely to get into the consideration set of decision makers (either as those who espouse them have greater incentive to do so, as in Osborne et al. 2000 and Levy and Razin 2013, or as they are simpler to communicate, as in Levy and Razin 2012).

In statistics and economics the ML approach was first formalized by Fisher (1912) for the purpose of parameter estimation,⁴ and by Robbins (1951), Good (1965), and Berger and Berliner (1986) for the purpose of forming posterior beliefs in the presence of ambiguity. Gilboa and Schmeidler (2003), Ortoleva (2012), and Suleymanov (2018) identify axioms that can rationalize maximum likelihood procedures while Gilboa and Schmeidler (2010) highlight the trade-off that can arise between maximizing likelihood and simplicity. In Epstein and Schneider (2007), who model learning with ambiguity, a decision maker weeds out explanations that yield a likelihood less than a fraction times

³In this sense our approach is related to small-sample biases such as the "hot hand" fallacy (see Rabin and Vayanos 2010), where individuals tend to over-infer correlations in the data rather than neglect them.

⁴Our analysis differs from the econometrics approach. In particular, our decision maker entertains a new information structure whenever new information arrives.

the maximum likelihood that can be achieved. Both [Ortoleva \(2012\)](#) and [Epstein and Schneider \(2007\)](#) use ML as a refinement criterion in extreme (zero or small) probability events, whereas we use ML as the main reasoning for decisions. Finally, [Salant and Cherry \(2020\)](#) suggest an equilibrium notion in which players best respond to a sample of their opponents' actions, where they make a statistical inference given this sample. One example they consider is for players to use ML to estimate the most likely parameter that generated the sample.

2. THE MODEL

A decision maker is forming a prediction about a state of the world $\omega \in \Omega$, where Ω is a finite set. She has a full support prior, $p \in \Delta(\Omega)$. At every period $t \in \{1, 2, \dots\}$, the decision maker observes a forecast: A forecast $q_t \in \Delta(\Omega)$ is a (full support) probability distribution over Ω . Let $\mathbf{q}^t \in (\Delta(\Omega))^t$ denote the vector of forecasts up to period t . At any period t , the decision maker's observed history is \mathbf{q}^t .

At every t , the decision maker combines these forecasts and the prior into a prediction, $\mu \in \Delta(\Omega)$, about the state. We assume that the decision maker thinks that at each period t , a single forecaster, with the same prior p , receives an informative signal and rationally derives q_t using Bayes' rule.⁵ To form a prediction, the decision maker considers *explanations*, which are Bayesian models that are consistent with the observations \mathbf{q}^t and the prior.

We now formally define what an explanation is. First, we define a joint information structure by a tuple $\mathcal{I} = (S, p, f(\mathbf{s}, \omega))$, such that $S = \times_{j=1}^t S_j$ is a finite set of signals, and for $\mathbf{s} \in S$, $f(\mathbf{s}, \omega)$ is a joint probability distribution over signals and states $\omega \in \Omega$. Given $f(\mathbf{s}, \omega)$, when they are well defined, we have $f(\mathbf{s}|\omega)$, $f(\omega|\mathbf{s})$ and $f(\mathbf{s})$ for all $\omega \in \Omega$. Moreover, $f(\mathbf{s}, \omega)$ also implies the marginal distributions over the realizations of signal s_j , $f_j(s_j|\omega)$.

An explanation of the data implies that any forecast q_j in \mathbf{q}^t was generated by a Bayesian forecaster who was exposed to a signal in S_j and knows the marginal distribution on signals in S_j , $f_j(s_j|\omega)$.

DEFINITION 1. An explanation of \mathbf{q}^t is $e = (\mathcal{I}^e, \mathbf{s}^e)$, where \mathcal{I}^e is a joint information structure and $\mathbf{s}^e = (s_j^e)_{j=1}^t \in S^e$ is a realization of signals such that for all $j \in \{1, 2, \dots, t\}$,

$$q_j(\omega) = \Pr^e(\omega|s_j^e) = \frac{p(\omega)f_j^e(s_j^e|\omega)}{\sum_{v \in \Omega} p(v)f_j^e(s_j^e|v)}.$$

In other words, the decision maker perceives some information structure \mathcal{I} and a particular realization of signals \mathbf{s} , such that all the forecasts q_j can be rationalized by Bayes' rule, assuming that each forecast q_j was based only on the signals generated by

⁵We can extend the analysis to the case of noncommon priors. The restrictions that forecasters use Bayes' rule and that the priors are known imply that there is no overfitting. If priors are both not known and not common, then the maximum likelihood of observing a vector of forecasts is 1.

$f_j(s_j|\omega)$. Note that this formulation is general in the sense that it allows for the possibility that s_j contains information observed by other forecasters $j' \neq j$.

Let $\mathcal{E}(\mathbf{q}^t)$ be the set of explanations as defined above. We assume that the decision maker uses the ML criterion to select which explanation in $\mathcal{E}(\mathbf{q}^t)$ to adopt to set her prediction. The likelihood of observing \mathbf{q}^t given an explanation $e \in \mathcal{E}(\mathbf{q}^t)$ is

$$L(\mathbf{q}^t|e) = \sum_{v \in \Omega} p(v) f^e(\mathbf{s}^e|v).$$

Below we show that $\arg \max_{e \in \mathcal{E}(\mathbf{q}^t)} L(\mathbf{q}^t|e)$ exists and, thus, we can define the ML prediction as follows.

ASSUMPTION 1 (ML prediction). At any period t , the decision maker’s prediction satisfies

$$\mu^{\text{ML}}(\omega|\mathbf{q}^t) = \frac{p(\omega) f^{\hat{e}}(\mathbf{s}^{\hat{e}}|\omega)}{\sum_{v \in \Omega} p(v) f^{\hat{e}}(\mathbf{s}^{\hat{e}}|v)} \quad \text{for some explanation } \hat{e} \in \arg \max_{e \in \mathcal{E}(\mathbf{q}^t)} L(\mathbf{q}^t|e).$$

Before proceeding to characterize ML explanations and predictions, we introduce a useful lemma that simplifies the analysis by reducing the set of explanations to those that have information structures with binary signals for each forecast.

LEMMA 1. Let $e = (I^e, s^e)$ be an explanation of \mathbf{q}^t . Then there exists an explanation $e' = (I^{e'}, s^{e'})$ of \mathbf{q}^t with $S^{e'} = \{s^*, s^{*-}\}^t$, and $L(\mathbf{q}^t|e) = L(\mathbf{q}^t|e')$.

Intuitively, the decision maker looks for an ex post rationalization, which implies that she has in mind a vector of signals attained by all forecasters. As a result, for any information structure that rationalizes the forecasts, we can construct an equivalent information structure that induces that vector of observed signals with the same probability and bunches all other (unobserved) signals together. This new information structure produces the observed forecasts with the same likelihood and, moreover, attains this with just two signals per forecaster.

3. THE ML PREDICTION

In this section, we characterize the ML prediction. We start with some helpful notation. Consider a history \mathbf{q}^t . For any forecast $j = 1, \dots, t$, let

$$\alpha_j(\omega) = \frac{q_j(\omega)}{p(\omega)}$$

and let

$$\bar{\alpha}_j = \max_{v \in \Omega} \alpha_j(v) \quad \text{and} \quad \omega^j \in \arg \max_{v \in \Omega} \alpha_j(v).$$

Thus, for each forecast j , ω^j can be described as the state that becomes most surprising given the prior and $\bar{\alpha}_j$ is the magnitude of this largest surprise. Next, note that

$$\frac{\alpha_j(\omega^j)}{\bar{\alpha}_j} = 1, \quad \text{whereas } \frac{\alpha_j(\omega)}{\bar{\alpha}_j} \leq 1 \text{ for any } \omega \neq \omega^j.$$

For example, if $|\Omega| = 3$ and the prior is uniform, then for a forecast $\{0.7, 0.2, 0.1\}$, the vector $\alpha_j(\omega)/\alpha_j(v)$ is $\{1, \frac{0.2}{0.7}, \frac{0.1}{0.7}\}$. As can be seen in the proof, $\alpha_j(\omega)/\alpha_j(v)$ is inherently related to the constraint that each forecaster uses Bayes' rule and specifically implies the probability that a forecast can be attained at state ω . Moreover, the most surprising state presents the upper bound of attaining a forecast, which is why the ratio $\alpha_j(\omega)/\bar{\alpha}_j$ is important.

With this notation, we can characterize the ML prediction of the decision maker.

PROPOSITION 1. *Given \mathbf{q}^t , the ML prediction of the decision maker is unique and equal to, for any $\omega \in \Omega$,*

$$\mu^{\text{ML}}(\omega|\mathbf{q}^t) = \frac{p(\omega) \min_{j=1,\dots,t} \left\{ \frac{\alpha_j(\omega)}{\bar{\alpha}_j} \right\}}{\sum_{v \in \Omega} p(v) \min_{j=1,\dots,t} \left\{ \frac{\alpha_j(v)}{\bar{\alpha}_j} \right\}}.$$

PROOF. Note that for any explanation e , for any $\omega \in \Omega$,

$$f^e(\mathbf{s}^e|\omega) \leq \min_{j=1,\dots,t} f_j^e(s_j^e|\omega). \tag{1}$$

Whereas an explanation is an information structure that rationalizes \mathbf{q}^t , we must have, for any $\omega, \omega' \in \Omega$,

$$\frac{q_j(\omega)}{q_j(\omega')} = \frac{p(\omega) f_j^e(s_j^e|\omega)}{p(\omega') f_j^e(s_j^e|\omega')}. \tag{2}$$

Choose some state v . Equation (2) implies that by setting $f_j^e(s_j^e|v)$ at some level, we pin down all values $f_j^e(s_j^e|\omega)$ for any $\omega \in \Omega$.

Using (1) and (2), we can write the upper bound for the likelihood of any explanation as

$$\begin{aligned} \sum_{\omega \in \Omega} p(\omega) f^e(\mathbf{s}^e|\omega) &\leq \sum_{\omega \in \Omega} p(\omega) \min_j \{ f_j^e(s_j^e|\omega) \} \\ &= p(v) \left[\sum_{\omega \in \Omega} \min_j \left\{ \frac{q_j(\omega)}{q_j(v)} f_j^e(s_j^e|v) \right\} \right]. \end{aligned}$$

To find the explanation that maximizes the likelihood of observing \mathbf{q}^t (the left-hand side above) we maximize the right-hand side and show that we can achieve a likelihood equal to the maximal upper bound.

First, to maximize the right-hand side, note that the problem is increasing in $f_j^e(s_j^e|v)$ for any $j \leq t$. By (2), for any $\omega \in \Omega$,

$$f_j^e(s_j^e|\omega) = \frac{q_j(\omega)}{q_j(v)} \frac{p(v)}{p(\omega)} f_j^e(s_j^e|v) \leq 1 \quad \Rightarrow \quad f_j^e(s_j^e|v) \leq \frac{p(\omega)}{q_j(\omega)} \frac{q_j(v)}{p(v)}$$

for any $\omega \in \Omega$. Recall, however, that ω^j maximizes $q_j(\omega)/p(\omega)$ and, hence, imposes the binding constraint on the upper bound on $f_j^e(s_j^e|v)$. Therefore, we set $f_j^e(s_j^e|v)$ at the upper bound:

$$f_j^e(s_j^e|v) = \frac{p(\omega^j)}{p(v)} \frac{q_j(v)}{q_j(\omega^j)} = \frac{\alpha_j(v)}{\bar{\alpha}_j}.$$

We can then use this and (2) to derive, for any other ω ,

$$f_j^e(s_j^e|\omega) = \frac{\alpha_j(\omega)}{\bar{\alpha}_j}.$$

Second, we construct our ML explanation, e^* , to attain the upper bound. For any j , let the set of signals be $S_j = \{s^*, s^{-*}\}$. Then set, for any ω ,

$$f_j^{e^*}(s^*|\omega) = \frac{\alpha_j(\omega)}{\bar{\alpha}_j}, \quad f_j^{e^*}(s^{-*}|\omega) = 1 - f_j^{e^*}(s^*|\omega).$$

By way of convention, order the signals so that s_j^* is “lower” than s_j^{-*} , and now set the cumulative marginal as $F_j^{e^*}(s^*|\omega) = f_j^{e^*}(s^*|\omega)$ and $F_j^{e^*}(s^{-*}|\omega) = 1$. As we have set the marginals, we can now set the joint distribution over signals. Note that given that we integrated the Bayes rule constraint for each forecast, which imposes how the probability that a forecast is attained is related across the states, we can focus only on the conditional joint distributions (that is, conditional on each state ω). In particular, the joint unconditional distribution over signals is determined first by the realization of the state of the world and then by the distribution over signals conditional on this state.

Fix some $\omega \in \Omega$. To attain the upper bound on the joint distribution over the observed signals, designated to be the vector \mathbf{s}^* , set

$$F^{e^*}(\mathbf{s}^*|\omega) = f^{e^*}(\mathbf{s}^*|\omega) = \min_j \{f_j^{e^*}(s^*|\omega)\} = \min_j \left\{ \frac{\alpha_j(\omega)}{\bar{\alpha}_j} \right\},$$

which implies that $F^{e^*}(\mathbf{s}|\omega) = \min_j F_j^{e^*}(s_j|\omega)$. To complete the explanation, continue by setting, for all \mathbf{s} ,

$$F^{e^*}(\mathbf{s}|\omega) = \min_j F_j^{e^*}(s_j|\omega).$$

The above equality is a proper distribution function, as given a set of marginals (here, $F_j^{e^*}(s_j|\omega)$ for all $j \leq t$), there is always a joint distribution function that attains the upper Frechet bound, which is the one defined above.⁶

⁶See Joe (1997).

We have constructed an explanation that consists of a set of conditional joint distribution functions and that achieves the upper bound. Note, moreover, that any explanation that achieves the upper bound must satisfy $f^{e^*}(\mathbf{s}^*|\omega) = \min_j\{\alpha_j(\omega)/\bar{\alpha}_j\}$ for some observed vector of signals \mathbf{s}^* . As a result, the ML prediction is unique and satisfies

$$\mu^{\text{ML}}(\omega|\mathbf{q}^t) = \frac{p(\omega)f^{e^*}(\mathbf{s}^*|\omega)}{\sum_{v \in \Omega} p(v)f^{e^*}(\mathbf{s}^*|v)} = \frac{p(\omega) \min_{j=1, \dots, t} \left\{ \frac{\alpha_j(\omega)}{\bar{\alpha}_j} \right\}}{\sum_{v \in \Omega} p(v) \min_{j=1, \dots, t} \left\{ \frac{\alpha_j(v)}{\bar{\alpha}_j} \right\}},$$

which is the expression in the proposition. □

From the above discussion it is clear that only a small set of forecasts would matter: those that minimize $\alpha_j(\omega)/\bar{\alpha}_j$ for some ω . Thus, at most $|\Omega|$ forecasts would be relevant for the final prediction, whereas other forecasts can be ignored. Moreover, the order in which these forecasts arrive before or at time t does not alter the prediction at time t . Note also the importance of the prior: The prior determines $\alpha_j(\omega)/\bar{\alpha}_j$, and, hence, affects which forecasts can be ignored and how those that are not ignored are combined into a prediction (we return to the role of the prior when we consider the limit predictions in Section 4.2).

In Section 3.1 below, we illustrate how the above result translates to a simple prediction rule in the binary state space, and we also construct an example of an ML explanation for $t = 2$. The ML explanation involves large degrees of correlation, and specifically it is moderate forecasts that are correlated to extreme ones and can, therefore, be ignored in the ML prediction. We show this feature more generally in Section 3.2.

3.1 Binary states: ML prediction and explanation

We now illustrate Proposition 1 (as well as its proof) in the binary state space, $\Omega = \{0, 1\}$. Assume without loss of generality a prior of $p(1) = p > \frac{1}{2}$. To simplify, with some abuse of notation, let q_j denote the probability that the state is 1 according to the forecast in period j . Note that in the binary space, $\omega^j = 1$ if $q > p$ and $\omega^j = 0$ if $q < p$.

Consider, for the sake of exposition, $t = 2$ and first assume that $q_1 > q_2 > p$. Thus, both forecasts “support” state 1 compared with the prior (so that $\omega^1 = \omega^2 = 1$).

Suppose, in line with Lemma 1, that each forecaster can receive two signal realizations, s^* and s^{-*} , which, given Bayesian updating, must then satisfy, for some $\beta_j \in (0, 1]$,

$$\begin{aligned} f^e(s_1 = s^*|\omega = 1) &= \beta_1, & f^e(s_1 = s^*|\omega = 0) &= \beta_1 \frac{p(1 - q_1)}{q_1(1 - p)} = \beta_1 \frac{\alpha_1(0)}{\bar{\alpha}_1} \\ f^e(s_2 = s^*|\omega = 1) &= \beta_2, & f^e(s_2 = s^*|\omega = 0) &= \beta_2 \frac{p(1 - q_2)}{q_2(1 - p)} = \beta_2 \frac{\alpha_2(0)}{\bar{\alpha}_2}. \end{aligned}$$

Maximizing β_1 and β_2 increases the marginal probability of attaining the s^* signals for both forecasters, as $q_j > p$, and so we can set $\beta_j = 1$. We continue then by setting the

joint probability over (s^*, s^*) :

$$f^e(s_1 = s^*, s_2 = s^* | \omega = 1) = 1 = f^e(s_1 = s^* | \omega = 1)$$

$$f^e(s_1 = s^*, s_2 = s^* | \omega = 0) = \min_j \frac{\alpha_j(0)}{\bar{\alpha}_j} = \frac{p(1 - q_1)}{q_1(1 - p)} = f^e(s_1 = s^* | \omega = 0).$$

Forecast 2 will be completely ignored as conditional on $s_1 = s^*$, the joint information structure constructed above necessitates $s_2 = s^*$ as well, and so the “observed” signal of the second forecaster is fully correlated with that of the first forecaster in each state. This readily implies the full ML joint information structure, depicted in Table 1, where each cell in each matrix denotes $f^e(s_1, s_2 | \omega)$.

As a result, the ML prediction satisfies

$$\mu = \frac{p}{p + (1 - p) \frac{p(1 - q_1)}{q_1(1 - p)}} = q_1,$$

implying that the more moderate forecast, q_2 , is completely ignored.

Assume instead that $q_2 < p < q_1$. To increase the likelihood of attaining q_2 —a forecast indicating that the state is more likely to be 0 compared with the prior—we need that in state 1 the second forecaster does not always observe s^* . Specifically, we would now have

$$f^e(s_1 = s^* | \omega = 1) = 1, \quad f^e(s_1 = s^* | \omega = 0) = \frac{p(1 - q_1)}{q_1(1 - p)}$$

$$f^e(s_2 = s^* | \omega = 1) = \frac{(1 - p)q_2}{p(1 - q_2)}, \quad f^e(s_2 = s^* | \omega = 0) = 1.$$

This implies the following joint ML information structure (Table 2), in which the “observed” signal of forecaster 1 is a sufficient statistic for the “observed” signal of forecaster 2 only in state 0, and the other way around in state 1.

As a result, the ML prediction now satisfies

$$\mu = \frac{p \frac{(1 - p)q_2}{p(1 - q_2)}}{p \frac{(1 - p)q_2}{p(1 - q_2)} + (1 - p) \frac{p(1 - q_1)}{q_1(1 - p)}} = \frac{q_2 q_1}{q_2 q_1 + \frac{p}{1 - p} (1 - q_1)(1 - q_2)}.$$

Corollary 1, derived from Proposition 1, shows how the above conclusion extends to any t .

COROLLARY 1. *Suppose that $\Omega = \{0, 1\}$. Let $q_t^{\max} = \max_{j \leq t} q_j$ and let $q_t^{\min} = \min_{j \leq t} q_j$. For any \mathbf{q}^t , the ML prediction is*

$$\mu^{\text{ML}}(1 | \mathbf{q}^t) = \begin{cases} q_t^{\max} & \text{if } q_t^{\min} \geq p \\ q_t^{\min} & \text{if } q_t^{\max} \leq p \\ \frac{q_t^{\min} q_t^{\max}}{q_t^{\min} q_t^{\max} + \frac{p}{(1 - p)} (1 - q_t^{\min})(1 - q_t^{\max})} & \text{otherwise.} \end{cases}$$

$\omega = 1$	$s_2 = s^*$	$s_2 = s^{-*}$	$\omega = 0$	$s_2 = s^*$	$s_2 = s^{-*}$
$s_1 = s^*$	1	0	$s_1 = s^*$	$\frac{p(1-q_1)}{q_1(1-p)}$	0
$s_1 = s^{-*}$	0	0	$s_1 = s^{-*}$	$\frac{p(1-q_2)}{q_2(1-p)} - \frac{p(1-q_1)}{q_1(1-p)}$	$1 - \frac{p(1-q_2)}{q_2(1-p)}$

TABLE 1. Joint ML information structure when $q_1 > q_2 > p$

$\omega = 1$	$s_2 = s^*$	$s_2 = s^{-*}$	$\omega = 0$	$s_2 = s^*$	$s_2 = s^{-*}$
$s_1 = s^*$	$\frac{(1-p)q_2}{p(1-q_2)}$	$1 - \frac{(1-p)q_2}{p(1-q_2)}$	$s_1 = s^*$	$\frac{p(1-q_1)}{q_1(1-p)}$	0
$s_1 = s^{-*}$	0	0	$s_1 = s^{-*}$	$1 - \frac{p(1-q_1)}{q_1(1-p)}$	0

TABLE 2. Joint ML information structure when $q_1 > p > q_2$

The implication of Proposition 1 to the binary state space is that at most two forecasts, the most extreme ones (on each side of the prior), do matter for the ML prediction. All other, more moderate, forecasts are ignored.

3.2 Ignoring moderate forecasts

In this section we characterize the set of forecasts that are ignored in the ML prediction. Formally, we say that forecast q_{t+1} is ignored if $\mu^{ML}(\omega|\mathbf{q}^{t+1}) = \mu^{ML}(\omega|\mathbf{q}^t)$.

Remember that for any forecast q_j , $\alpha_j(\omega)$ is the ratio of the probability of this state under the forecast divided by the prior probability. We have also introduced the notation of $\bar{\alpha}_j = \max_{v \in \Omega} \alpha_j(v)$ and $\omega^j \in \arg \max_{v \in \Omega} \alpha_j(v)$. Thus, for each forecast q_j , ω^j can be described as the state that becomes most surprising given the prior, and $\bar{\alpha}_j$ is the magnitude of this largest surprise.

Going back to the proof of Proposition 1, note that Bayesian updating bounds the probability that a forecast can be sent at each state ω . A forecast q_j , which is almost degenerate on state ω^j (hence, extreme) can be sent in any other state ω with only a very small probability, as we would have $\alpha_j(\omega)/\bar{\alpha}_j$; otherwise it could not indicate to a rational forecaster that the state is most likely to be ω^j . Thus, this forecast bounds the joint probability of obtaining all forecasts in all states $\omega \neq \omega^j$ and, therefore, matters for the ML prediction. Alternatively, a moderate forecast that agrees with the prior, $q_j(\omega) = p(\omega)$, can be sent in each state ω with probability $\alpha_j(\omega)/\bar{\alpha}_j = 1$. It then never imposes a bound on the joint probability distribution and does not affect the ML prediction. We now characterize the set of moderate forecasts, such as above, that are ignored in the final aggregation of forecasts. Let

$$I(\mathbf{q}^t) = \left\{ q_{t+1} \in \Delta(\Omega) \mid \forall v \in \Omega, \frac{\alpha_{t+1}(v)}{\bar{\alpha}_{t+1}} \geq \min_{j \leq t} \frac{\alpha_j(v)}{\bar{\alpha}_j} \text{ and } \exists \omega \in \Omega \text{ such that } \frac{\alpha_{t+1}(\omega)}{\bar{\alpha}_{t+1}} = \min_{j \leq t} \frac{\alpha_j(\omega)}{\bar{\alpha}_j} \right\}.$$

The set $I(\mathbf{q}^t)$ defines the boundary of the set of all forecasts that the decision maker is able to ignore at period $t + 1$. This boundary set contains forecasts that minimize

$\alpha_j(\omega)/\bar{\alpha}_j$ for some state ω , for $j \leq t$ and, hence, contains forecasts that have not been ignored up to and including period t . To gain intuition, let us consider the set $I(\mathbf{q}^t)$ in the case of binary states. Suppose that $\min_{j \leq t} q_j < p < \max_{j \leq t} q_j$. Assume without loss of generality that $\max_{j \leq t} q_j > 1 - \min_{j \leq t} q_j$ so that $\bar{\alpha}_j = \max_{j \leq t} q_j$. In this case, we have that

$$\min_{j \leq t} \frac{\alpha_j(1)}{\bar{\alpha}_j} = \frac{\min_{j \leq t} q_j}{\max_{j \leq t} q_j} \quad \text{and} \quad \min_{j \leq t} \frac{\alpha_j(0)}{\bar{\alpha}_j} = \frac{1 - \max_{j \leq t} q_j}{\max_{j \leq t} q_j}.$$

Therefore, if $q_{t+1} = \max_{j \leq t} q_j$, then

$$\frac{\alpha_{t+1}(1)}{\bar{\alpha}_{t+1}} = 1 > \frac{\min_{j \leq t} q_j}{\max_{j \leq t} q_j} \quad \text{and} \quad \frac{\alpha_{t+1}(0)}{\bar{\alpha}_{t+1}} = \frac{1 - \max_{j \leq t} q_j}{\max_{j \leq t} q_j} = \min_{j \leq t} \frac{\alpha_j(0)}{\bar{\alpha}_j}.$$

Similarly, when $q_{t+1} = \min_{j \leq t} q_j$, we have that

$$\frac{\alpha_{t+1}(1)}{\bar{\alpha}_{t+1}} = \frac{\min_{j \leq t} q_j}{\max_{j \leq t} q_j} = \min_{j \leq t} \frac{\alpha_j(1)}{\bar{\alpha}_j} \quad \text{and} \quad \frac{\alpha_{t+1}(0)}{\bar{\alpha}_{t+1}} = \frac{1 - \min_{j \leq t} q_j}{\max_{j \leq t} q_j} > \frac{1 - \max_{j \leq t} q_j}{\max_{j \leq t} q_j} = \min_{j \leq t} \frac{\alpha_j(0)}{\bar{\alpha}_j}.$$

For any $q_{t+1} \in (\min_{j \leq t} q_j, \max_{j \leq t} q_j)$, both inequalities are strict, and for any $q_{t+1} \notin [\min_{j \leq t} q_j, \max_{j \leq t} q_j]$, one of the inequalities holds in the opposite way. As a result, in this case, we have $I(\mathbf{q}^t) = \{\min_{j \leq t} q_j, \max_{j \leq t} q_j\}$.

Using the definition of the set $I(\mathbf{q}^t)$, we show the following result for the general state space.

LEMMA 2. *For any $q \in I(\mathbf{q}^t)$ and $\beta \in [0, 1]$, the forecast $q_{t+1} = \beta p + (1 - \beta)q$ is ignored.*

If we define moderation by the distance to the prior, Lemma 2 implies that all forecasts that are more moderate (weakly) than those in $I(\mathbf{q}^t)$ are ignored. In the case of binary states, when $\min_{j \leq t} q_j < p < \max_{j \leq t} q_j$, we saw above that $I(\mathbf{q}^t) = \{\min_{j \leq t} q_j, \max_{j \leq t} q_j\}$. Thus, all more moderate forecasts on $[\min_{j \leq t} q_j, p]$ and on $[p, \max_{j \leq t} q_j]$ are ignored. To see another example, consider a tertiary state space, $\Omega = \{0, 1, 2\}$. With equal prior, the forecast $\{0.7, 0.2, 0.1\}$ allows us to ignore all forecasts for which the vector $\alpha_j(\omega)/\alpha_j(\omega)$ is at least, element by element, $\{1, \frac{0.2}{0.7}, \frac{0.1}{0.7}\}$. The set of such ignored forecasts is depicted in Figure 1. As can be seen, it is all those forecasts that have the same mode, at $\omega = 0$, and lie on the line that connects each point on the closure to the prior.

Figure 2 shows the set of forecasts that are ignored when $\{0.7, 0.2, 0.1\}$ is observed and the set of forecasts that are ignored when $\{0.2, 0.1, 0.7\}$ is observed. These are the two light shaded areas in the figure. The set of forecasts that are ignored when both $\{0.7, 0.2, 0.1\}$ and $\{0.2, 0.1, 0.7\}$ are observed includes more forecasts than just the union of the above two sets. The additional forecasts that are ignored when both $\{0.7, 0.2, 0.1\}$ and $\{0.2, 0.1, 0.7\}$ are observed is given by the darker shaded area in the figure. Thus, the total set of forecasts that are ignored when both $\{0.7, 0.2, 0.1\}$ and $\{0.2, 0.1, 0.7\}$ are observed includes the union of the three shaded areas depicted in the figure.




FIGURE 1. Ignored forecasts when $(0.7, 0.2; 0.1)$ is observed.




FIGURE 2. Ignored forecasts with $(0.7, 0.2, 0.1)$ and $(0.7, 0.1, 0.2)$.

3.3 Time consistency of ML explanations

The ML decision maker potentially has a wrong model in mind, and with every new observation she adopts a different world view or narrative to interpret the accumulated data she has observed. However, if the procedure calls for a completely new model each period, we might question the sensibility of this method of aggregating data. Moreover, such a procedure implies higher computation costs, as each period a new model needs to be calculated.

Below we formalize a notion of consistency between explanations and show that the ML explanation can evolve in a time-consistent manner. Let e be an explanation of \mathbf{q}^t and let e' be an explanation of \mathbf{q}^{t+1} . We say that e' is time consistent with e if $S^{e'} = S^e \times S_{t+1}$ for some finite set S_{t+1} and for any $\mathbf{s} \in S^e, \omega \in \Omega$,

$$f^e(\mathbf{s}|\omega) = \sum_{(\mathbf{s}, s_{t+1}) \in S^{e'}} f^{e'}(\mathbf{s}, s_{t+1}|\omega).$$

In words, $f^e(\mathbf{s}|\omega)$, the probability of observing \mathbf{s} in explanation e , is equal to the expected probability of observing signal realizations \mathbf{s} in the first t periods under explanation e' . This implies that the explanation e for the first t observations is the projection of e' to

these t observation. As a result, if there is time consistency between e and e' , we can say that when moving from period t to period $t + 1$, the decision maker keeps, in a statistical sense, her explanation for the first t observations.

We can then show the following statement.

PROPOSITION 2. *For any explanation of e of \mathbf{q}^t , for any observation q_{t+1} , there exists an ML explanation e' of (\mathbf{q}^t, q_{t+1}) that is time consistent with e .*

Intuitively, for any state ω , there are two cases. (i) The new forecast is ignored, which means that the probability of observing it is higher than observing a previous one, and so we can fully correlate it with the previous vector of forecasts and peg the remaining probability of observing the new forecast onto the other, unobserved, realizations of the first t forecasts. (ii) Alternatively, old forecasts are ignored, so we fully correlate old forecasts to the new one and the remaining probability of observing the old forecasts is pegged onto other (unobserved) realizations of the $t + 1$ forecast. As a result, the decision maker does not need to come up with a completely new ML explanation each time and can be consistent in how she explains her data over time.

To get more intuition, recall the joint information structure described in the case of binary states, when $q_1 > q_2 > p$, which, for the observed signals, satisfied

$$f^e(s_1 = s^*, s_2 = s^* | \omega = 1) = 1, \quad f^e(s_1 = s^*, s_2 = s^* | \omega = 0) = \frac{p(1 - q_1)}{q_1(1 - p)}.$$

Assume that we now observe some q_3 where specifically we have $q_3 < p$, for which the ML information structure demands a marginal probability of

$$f^e(s_3 = s^* | \omega = 1) = \frac{(1 - p)q_3}{p(1 - q_3)}, f^e(s_3 = s^{-*} | \omega = 0) = 1.$$

Consider the following joint ML structure, where we combine the above marginal of the first two signals, s_1 and s_2 , with that of s_3 . First, we describe $f^e(s_1, s_2, s_3 = s^* | 1)$ and $f^e(s_1, s_2, s_3 = s^{-*} | 1)$ (see Table 3).

We now depict $f^e(s_1, s_2, s_3 = s^* | 0)$ and $f^e(s_1, s_2, s_3 = s^{-*} | 0)$ (see Table 4).

The marginal distribution over s_1, s_2 derived from the above structure is exactly as it is for the two signals generating $q_1 > q_2 > p$, as described in Section 3.1. Moreover, following from Proposition 1, the above structure is an ML information structure.

The forecast q_3 cannot be ignored and the signal generating it, $s_3 = s^*$, has the minimum probability of being attained in state 1 compared with the probability of attaining $s_1 = s_2 = s^*$. As a result, these signals are fully correlated to $s_3 = s^*$ in state 1 and the remaining probability of attaining $s_1 = s_2 = s^*$ in this state is pegged onto the (non)observation of $s_3 = s^{-*}$. In state 0, the joint probability of attaining $s_1 = s_2 = s^*$ is lower than that of $s_3 = s^*$, which is 1. As a result, $s_3 = s^*$ arises whenever $s_1 = s_2 = s^*$ arises, but also arises for other configurations of these signals. The proof in the Appendix uses similar arguments to construct a time-consistent joint distribution between the marginal of the first t signals and the additional signal arising in the $t + 1$ th period.

$\omega = 1, s_3 = s^*$	$s_2 = s^*$	$s_2 = s^{-*}$	$\omega = 1, s_3 = s^{-*}$	$s_2 = s^*$	$s_2 = s^{-*}$
$s_1 = s^*$	$\frac{(1-p)q_3}{p(1-q_3)}$	0	$s_1 = s^*$	$1 - \frac{(1-p)q_3}{p(1-q_3)}$	0
$s_1 = s^{-*}$	0	0	$s_1 = s^{-*}$	0	0

TABLE 3. Joint ML information structure when $q_1 > q_2 > p > q_3$ and $w = 1$

$\omega = 0, s_3 = s^*$	$s_2 = s^*$	$s_2 = s^{-*}$	$\omega = 0, s_3 = s^{-*}$	$s_2 = s^*$	$s_2 = s^{-*}$
$s_1 = s^*$	$\frac{p(1-q_1)}{q_1(1-p)}$	$\frac{p(1-q_2)}{q_2(1-p)} - \frac{p(1-q_1)}{q_1(1-p)}$	$s_1 = s^*$	0	0
$s_1 = s^{-*}$	0	$1 - \frac{p(1-q_2)}{q_2(1-p)}$	$s_1 = s^{-*}$	0	0

TABLE 4. Joint ML information structure when $q_1 > q_2 > p > q_3$ and $w = 0$

4. EXTENSIONS

We consider two extensions. In the first one, we analyze a model in which the decision maker does not observe forecasts, but only the history of her past ML predictions. The second extension looks at limit predictions and highlights the importance of the prior belief in the ML process. To simplify the exposition, we henceforth focus on a binary state space $\Omega = \{0, 1\}$. Thus, a forecast q_j simply denotes the probability that the state is 1.

4.1 Source amnesia

We have so far assumed that the decision maker remembers the history of forecasts she has seen. We now consider an extension to a decision maker who does not remember forecasts but does remember her past predictions. This form of memory is termed *source amnesia* in the psychological literature.⁷

Specifically, in our previous model, at time t , the decision maker observes $\mathbf{q}^t = (q_1, q_2, \dots, q_t)$. We now assume that at time t , a decision maker observes q_t , and $\boldsymbol{\mu}^{t-1} = (\mu_1, \mu_2, \dots, \mu_{t-1})$, where μ_j is the ML prediction at period j .

Assume that the prior is uniform and suppose without loss of generality that $q_1 > \frac{1}{2}$. Let us consider period 3, where the decision maker observes q_3, μ_2 , and $\mu_1 = q_1$. She knows that for whichever q_2 she imagines, it has to be that $\mu_2 = \mu^{\text{ML}}(\cdot | \mathbf{q}^2)$. By [Corollary 1](#), if $\mu_2 \geq q_1$, then $\mu_2 = q_2$. If $\mu_2 = q_1$, then $q_1 > q_2 > \frac{1}{2}$ and it is not important for her to know the exact value of q_2 . If $\mu_2 < q_1$, then

$$q_2 = \frac{\mu^2(1 - q_1)}{q_1 + \mu^2(1 - 2q_1)}.$$

Thus, in period 3 she either knows q_2 or knows that it should be ignored. It follows then that at period t , for each $j < t$, she either knows q_j or knows that it could be ignored. We therefore have the following proposition.

⁷See, for example, [Schacter et al. \(1984\)](#).

PROPOSITION 3. *A decision maker who has source amnesia has the same ML prediction as a decision maker who remembers the history of forecasts.*

REMARK 1 (Social learning). Social learning is typically modelled as an environment in which an individual at period t observes the sequence of the actions or predictions of her predecessors, as well as her own forecast. Thus, as above, adopting social learning (with observed beliefs) to our setup, she observes $\boldsymbol{\mu}^{t-1} = \{\mu_1, \mu_2, \dots, \mu_{t-1}\}$ as well as q_t . If we assume that every decision maker believes that every one before her also uses the ML procedure and that this is common knowledge, then the problem becomes identical to the single agent problem. Given this, all the results above hold. Specifically, when all previous predictions are observed, then the sequence of predictions is identical to the one that arises if all forecasts were shared instead.

4.2 The persistence of the prior

In this section we explore the convergence properties of the ML predictions. The key observation that we highlight is the importance of the prior in the limit ML predictions. Specifically, we show that even when exposed to a large data set, the individual’s limit belief might still depend on her initial prior.

To focus our discussion, assume a true information generating process by which signals are drawn in an independent and identically distributed (iid) manner each period. In particular, assume that the information structure induces a distribution F over forecasts in $(0, 1)$, with $a = \inf(\text{Support } F)$ and $b = \sup(\text{Support } F)$. Assume that the prior p satisfies $p \in (a, b)$ and $0 \leq a < b \leq 1$, so that the posteriors are informative. When the individual observes forecasts drawn from this process, where would her predictions converge?

PROPOSITION 4. (i) *Suppose that $a > 0$ and $b < 1$. The prediction of the ML decision maker satisfies, for any $\varepsilon > 0$,*

$$\lim_{t \rightarrow \infty} \Pr \left[\mu^{\text{ML}}(1|\mathbf{q}^t) \in \left(\frac{1}{1 + \frac{p}{1-p} \frac{(1-a)(1-b)}{b} \frac{1}{a}} - \varepsilon, \frac{1}{1 + \frac{p}{1-p} \frac{(1-a)(1-b)}{b} \frac{1}{a}} + \varepsilon \right) \right] = 1.$$

(ii) *Suppose that $a = 0$ and $b = 1$. There exist distributions F for which, for any $\varepsilon, \delta > 0$,*

$$\lim_{t \rightarrow \infty} \Pr[\mu^{\text{ML}}(1|\mathbf{q}^t) \in (p - \varepsilon, p + \varepsilon)] > \delta.$$

The ML decision maker updates based on just the extreme forecasts she has observed along the sequence, on each side of the prior. To see (i), note that in the limit, by the law of large numbers, these extreme forecasts converge to be a and b . Therefore, by **Proposition 1**, the ML prediction converges to the expression in the proposition. As

$$0 < \frac{(1-a)(1-b)}{b} \frac{1}{a} < \infty,$$

the limit prediction still depends on the prior.

What happens when $[a, b] = [0, 1]$, that is, when we have a rich signal structure that can induce posteriors of unlimited accuracy? For case (ii), we need to look at the limit statistical properties of \tilde{q}_t^{\max} and \tilde{q}_t^{\min} , as they determine whether the two posteriors converge. Specifically, consider a distribution over posteriors conditional on the state $\omega = 1$. Given that the state is 1, the probability of observing a posterior close to 0 must be close to 0. This is because such a posterior is based on Bayesian updating and so for a forecaster to believe that the state is likely to be 0, it must be that this posterior is sent in state 1 only with a small probability. This implies that the distribution over posteriors given $\omega = 1$ must have a “thin tail” around 0. But this distribution can also have a similar thin tail around posteriors that are close to 1. Extreme value theory tells us that we can construct distributions of \tilde{q}_t^{\max} and \tilde{q}_t^{\min} to accord, in the limit, with the Gumbel distribution.⁸ This allows us to construct information structures for which there is a strictly positive probability that the limit of the ratio

$$\frac{1 - \tilde{q}_t^{\max}}{\tilde{q}_t^{\min}}$$

is bounded and equal to 1, resulting in the ML belief converging to the prior p .

REMARK 2 (Divergence). Consider now two ML decision makers with different priors, where each assumes that the forecasts she observes are generated according to an information structure incorporating her own prior. The above result implies that there is a strictly positive probability of divergence between the two ML predictions they generate, even after being exposed to the same rich and large set of observations. This differs from the convergence properties of Bayesian decision makers with heterogeneous priors. For example, in a recent paper [Acemoglu et al. \(2016\)](#) show that decision makers with heterogeneous priors, who agree about the information process generating their signals, converge to have the same beliefs.⁹

5. CONCLUSION

There is some recent empirical and experimental evidence showing that decision makers tend to neglect correlation in some environments (e.g., [Ortoleva and Snowberg 2015](#), [Enke and Zimmerman 2019](#), [Kallir and Sonsino 2009](#), and [Eyster and Weizsacker 2011](#)), while in other contexts they overestimate correlation: Consistent with the maximum likelihood approach, we assumed in this paper, [De Filippis et al. \(2017\)](#) and [Hossain and Okui \(2020\)](#) derive results in which subjects sometimes overestimate the level of correlation.¹⁰ It would be interesting to understand empirically when it is more likely for

⁸For example, see [Fisher and Tippett \(1928\)](#).

⁹See Section 3.1 in their paper, where they use this result as a benchmark to show, in the main part of the paper, that small disagreement about the information generating process can imply large disagreements about where the process converges.

¹⁰Specifically, [De Filippis et al. \(2017\)](#) consider a class of updating rules that generalize the ML updating rule, and their experimental evidence supports the rule with a biased critical value (rather than an unbiased critical value).

decision makers to become aware of or, alternatively, excessively consider, correlation in their observed forecasts. One possibility is that observing many repeated forecasts increases the suspicion of the decision maker that these are correlated. One interesting example for such reasoning is the Talmudic Sanhedrin court law that requires that if judges are unanimous in conviction, the defendant should be set free, while if only a majority convict, this majority verdict pertains. Glatt (2013) offers a maximum likelihood rationalization of this rule: unanimity among many judges most likely is a result of strong correlation between the judges and, therefore, demands caution. Gunn et al. (2016) also discuss this interpretation in other legal scenarios.

APPENDIX

PROOF OF LEMMA 1. Assume that $e = (I^e, \mathbf{s}^e)$ is an explanation of \mathbf{q}^t . We construct a new explanation of \mathbf{q}^t with $S_j = \{s^*, s^{-*}\}$ for any j , which maintains the same likelihood of observing \mathbf{q}^t as e does.

Specifically, construct the new explanation $e' = (I^{e'}, \mathbf{s}^{e'})$ as follows. Let $\mathbf{s}^{e'} = (s^*, s^*, \dots, s^*)$. Let, $\forall \omega \in \Omega$,

$$(*) \quad f_j^{e'}(s^*|\omega) = f_j^e(s_j^e|\omega) \quad \text{for all } j \leq t; \quad f^{e'}(s^*, s^*, \dots, s^*|\omega) = f^e(\mathbf{s}^e|\omega)$$

and $\forall \mathbf{s} \neq \mathbf{s}^*, \mathbf{s} \in \{s^*, s^{-*}\}^t$,

$$f^{e'}(\mathbf{s}|\omega) = \sum_{\substack{\mathbf{s}' \in S^e \text{ s.t.} \\ \text{if } s_j = s^* \text{ then } s'_j = s_j^e \\ \text{otherwise } s'_j \neq s_j^e}} f^e(\mathbf{s}'|\omega).$$

Note that by (*), e' is an explanation of \mathbf{q}^t , as the marginal distributions of the observed signals are maintained. Moreover, by (*), it is an explanation that maintains the same likelihood of observing \mathbf{q}^t as e induces. □

PROOF OF LEMMA 2. We start by showing that a forecast $q_h = \beta p + (1 - \beta)q_l$ can be ignored if $q_l \in I(\mathbf{q}^t)$ for any $\beta \in (0, 1]$. To see this, note first that $\omega^h = \omega^l$: this arises as

$$\begin{aligned} \arg \max_{\omega} \frac{q_h(\omega)}{p(\omega)} &= \arg \max_{\omega} \frac{\beta p(\omega) + (1 - \beta)q_l(\omega)}{p(\omega)} \\ &= \arg \max_{\omega} \left(\beta + (1 - \beta) \frac{q_l(\omega)}{p(\omega)} \right) = \arg \max_{\omega} \frac{q_l(\omega)}{p(\omega)}. \end{aligned}$$

Then note that for any ω ,

$$\frac{\alpha_h(\omega)}{\bar{\alpha}_h} = \frac{\frac{q_h(\omega)}{p(\omega)}}{\frac{q_h(\omega^l)}{p(\omega^l)}} = \frac{\frac{\beta p(\omega) + (1 - \beta)q_l(\omega)}{p(\omega)}}{\frac{\beta p(\omega^l) + (1 - \beta)q_l(\omega^l)}{p(\omega^l)}} = \frac{\beta + (1 - \beta) \frac{q_l(\omega)}{p(\omega)}}{\beta + (1 - \beta) \frac{q_l(\omega^l)}{p(\omega^l)}} \geq \frac{\frac{q_l(\omega)}{p(\omega)}}{\frac{q_l(\omega^l)}{p(\omega^l)}} = \frac{\alpha_l(\omega)}{\bar{\alpha}_l}.$$

Now note that all forecasts in $q_{t+1} \in I(\mathbf{q}^t)$ can be ignored by Proposition 1. As a result of this and the above results, all forecasts $q_{t+1} = \beta p + (1 - \beta)q$ for some $q \in I(\mathbf{q}^t)$ and $\beta \in [0, 1]$ can be ignored. \square

PROOF OF PROPOSITION 2. Suppose that $e_t^* = (\mathcal{I}^{e_t^*}, \mathbf{s}^{e_t^*})$ is an ML explanation of \mathbf{q}^t . Recall that by Proposition 1, any ML explanation requires that

$$f^e(\mathbf{s}^e | \omega) = \min_{j \in \{1, 2, \dots, t\}} \frac{\alpha_j(\omega)}{\bar{\alpha}_j}.$$

We now construct a new ML explanation, e_{t+1}^* for \mathbf{q}^{t+1} , which is consistent with e_t^* . To construct $e_{t+1}^* = (\mathcal{I}^{e_{t+1}^*}, \mathbf{s}^{e_{t+1}^*})$, we set (i) $S^{e_{t+1}^*} = \times_{j=1}^{t+1} S_j^{e_{t+1}^*}$, with $\times_{j=1}^t S_j^{e_{t+1}^*} = S^{e_t^*}$, $S_{t+1}^{e_{t+1}^*} = \{s_{t+1}^{e_{t+1}^*}, s_{t+1}^{-e_{t+1}^*}\}$, and $\mathbf{s}^{e_{t+1}^*} = (s_1^{e_{t+1}^*}, \dots, s_t^{e_{t+1}^*}, s_{t+1}^{e_{t+1}^*})$ such that $(s_1^{e_{t+1}^*}, \dots, s_t^{e_{t+1}^*}) = \mathbf{s}^{e_t^*}$. (ii) We set

$$f_j^{e_{t+1}^*}(s_j^{e_{t+1}^*} | \omega) = \frac{\alpha_j(\omega)}{\bar{\alpha}_j} \quad \text{for all } j \leq t + 1.$$

(iii) The marginal of $f^{e_{t+1}^*}(\cdot | \omega)$ on $S^{e_t^*}$ to equal $f^{e_t^*}(\cdot | \omega)$.

We now set two cumulative marginals: on the first t signals and on the $t + 1$ th signal. Order signals in $S_j^{e_{t+1}^*}$ as a normalization so that $s_j^{e_{t+1}^*}$ is the smallest in $S_j^{e_{t+1}^*}$. This and (i)–(iii) imply that we have

$$\begin{aligned} F_{t+1}^{e_{t+1}^*}(s_{t+1}^{e_{t+1}^*} | \omega) &= f_{t+1}^{e_{t+1}^*}(s_{t+1}^{e_{t+1}^*} | \omega), & F_{t+1}^{e_{t+1}^*}(s_{t+1}^{-e_{t+1}^*} | \omega) &= 1 \\ F^{e_{t+1}^*}(\mathbf{s}^{e_t^*} | \omega) &= f^{e_{t+1}^*}(\mathbf{s}^{e_t^*} | \omega) = f^{e_t^*}(\mathbf{s}^{e_t^*} | \omega), \end{aligned}$$

where the remainder of $F^{e_{t+1}^*}(\mathbf{s}^t | \omega)$ on $\mathbf{s}^t \in S^{e_t^*}$, $\mathbf{s}^t \neq \mathbf{s}^{e_t^*}$, can be completed using (iii).

We conclude the construction by setting the joint distribution on $S^{e_{t+1}^*}$, and combining the marginals $F^{e_{t+1}^*}(\mathbf{s}^t | \omega)$ and $F_{t+1}^{e_{t+1}^*}(s_{t+1} | \omega)$. We first consider $\mathbf{s}^{e_{t+1}^*}$. For each state ω , and $\mathbf{s}^{e_{t+1}^*}$, we set

$$\begin{aligned} (**) \quad F^{e_{t+1}^*}(\mathbf{s}^{e_{t+1}^*} | \omega) &= f^{e_{t+1}^*}(\mathbf{s}^{e_{t+1}^*} | \omega) \\ &= \min\{f_{t+1}^{e_{t+1}^*}(s_{t+1}^{e_{t+1}^*} | \omega), f^{e_{t+1}^*}(\mathbf{s}^{e_t^*} | \omega)\} \\ &= \min\{F_{t+1}^{e_{t+1}^*}(s_{t+1}^{e_{t+1}^*} | \omega), F^{e_{t+1}^*}(\mathbf{s}^{e_t^*} | \omega)\}. \end{aligned}$$

Next, as we did in Proposition 1, for all other $\mathbf{s} = \{\mathbf{s}^t, s_{t+1}\} \neq \mathbf{s}^{e_{t+1}^*}$, we set

$$F^{e_{t+1}^*}(\mathbf{s} | \omega) = \min\{F_{t+1}^{e_{t+1}^*}(s_{t+1} | \omega), F^{e_t^*}(\mathbf{s}^t | \omega)\}.$$

This is a proper cumulative distribution function (cdf), as there is always a joint information structure that achieves the upper Frechet bound.

We have therefore constructed an explanation e_{t+1}^* that (a) explains the data by (ii), (b) maximizes the likelihood of the data, as by (**), (ii), and (iii),

$$f^{e_{t+1}^*}(\mathbf{s}^{e_{t+1}^*}|\omega) = \min_{j \in \{1, 2, \dots, t+1\}} \frac{\alpha_j(\omega)}{\bar{\alpha}_j},$$

and (c) is time-consistent by construction, following (iii). □

The proof of [Proposition 3](#) is provided in the text.

PROOF OF PROPOSITION 4. (i) This is a corollary of [Proposition 1](#) and the law of large numbers. (ii) In what follows, we denote a forecast about the state by $q \in [0, 1]$, interpreted as the probability of state 1, where p is the prior probability that the state is 1. We construct a true signal generating process by choosing the cumulative distributions over posteriors it generates, $F(q|1)$ in state 1 and $F(q|0)$ in state 0, with corresponding continuous densities $f(q|1)$ and $f(q|0)$.

Let us focus on state $\omega = 0$ and, hence, on $F(q|0)$. We consider a distribution that has symmetric tails so that $F(q|0) = 1 - F(1 - q|0)$ for all $q > \hat{q}$ for some $\hat{q} > 0.5$. To approximate the limit distribution of the extreme posteriors, we note that these are the extreme values of $F(q|0)$. We can, therefore, use the extreme value limit results in [Fisher and Tippett \(1928\)](#) and [Leadbetter et al. \(1983\)](#). In particular, we construct $F(q|0)$ to satisfy

$$\exists \gamma(q) > 0 \text{ such that } \lim_{q \rightarrow 1} \frac{1 - F(q + x\gamma(q)|0)}{1 - F(q|0)} = e^{-x} \text{ for any } x \in R.$$

By [Leadbetter et al. \(1983\)](#), for distributions that satisfy the above condition, there exist sequences $\{a_n\}, \{b_n\}$ such that $a_n = \gamma(F^{-1}(1 - 1/n)) \rightarrow 0$ and $b_n = F^{-1}(1 - 1/n) \rightarrow 1$ so that we can use the Gumbel distribution to approximate the distribution of the maximal posterior of $F(q|0)$, $\max V_n$:

$$\frac{\max V_n - b_n}{a_n} \sim \exp\left\{-\exp\left(-\frac{\max V_n - b_n}{a_n}\right)\right\}.$$

So if we choose two cutoffs $0 < \alpha_n < \beta_n < 1$ such that

$$\frac{\beta_n - b_n}{a_n} \rightarrow \beta, \quad \frac{\alpha_n - b_n}{a_n} \rightarrow \alpha$$

and $\beta > \alpha$, we have that the probability $\max V_n \in [\alpha_n, \beta_n]$ converges to $\exp\{-\exp(\beta)\} - \exp\{-\exp(\alpha)\} > 0$. Similarly, by symmetry, for the minimum value $\min V_n$, we have that the probability $\min V_n \in [1 - \beta_n, 1 - \alpha_n]$ converges to $\exp\{-\exp(\beta)\} - \exp\{-\exp(\alpha)\}$.

Note that as

$$\frac{1 - F(q + x\gamma(q)|0)}{1 - F(q|0)} \rightarrow_{q \rightarrow 1} e^{-x}$$

for any $x \in R$, this implies that for large enough q , $q + x\gamma(q) \leq 1$, and so we have

$$\gamma(q) < \frac{1 - q}{x}$$

for any $x \in R$. As a result, we have that

$$\frac{1 - \beta_n}{1 - \alpha_n} \simeq \frac{1 - b_n - \beta a_n}{1 - b_n - \alpha a_n} = \frac{\frac{1 - b_n}{a_n} - \beta}{\frac{1 - b_n}{a_n} - \alpha} \rightarrow 1,$$

as for large enough q ,

$$\frac{1 - b_n}{a_n} = \frac{1 - F^{-1}\left(1 - \frac{1}{n}\right)}{\gamma\left(F^{-1}\left(1 - \frac{1}{n}\right)\right)} > x \frac{1 - F^{-1}\left(1 - \frac{1}{n}\right)}{1 - F^{-1}\left(1 - \frac{1}{n}\right)} = x$$

for any $x \in R$.

Therefore, with probability $(\exp\{-\exp(\beta)\} - \exp\{-\exp(\alpha)\})^2$, we have that

$$1 \leftarrow \frac{1 - \beta_n}{1 - \alpha_n} < \frac{1 - \max V_n}{\min V_n} < \frac{1 - \alpha_n}{1 - \beta_n} \rightarrow 1.$$

Moreover, note that

$$\frac{1 - \min V_n}{\max V_n} \rightarrow 1.$$

Therefore, there is a strictly positive probability that the prior will matter for the limit prediction; in other words, with probability $(\exp\{-\exp(\beta)\} - \exp\{-\exp(\alpha)\})^2$,

$$\mu = \frac{1}{1 + \frac{1-p}{p} \frac{1 - \max V_n}{\min V_n} \frac{1 - \min V_n}{\max V_n}} \rightarrow_{n \rightarrow \infty} P.$$

Finally, we need to check that the above result can indeed be constructed as a distribution over posteriors. A distribution over posteriors arising from Bayesian updating implies a joint restriction on $f(q|1)$ and $f(q|0)$ so that for any q ,¹¹

$$q = \frac{pf(q|\omega = 1)}{pf(q|\omega = 1) + (1 - p)f(q|\omega = 0)} \Rightarrow f(q|1) = f(q|0) \frac{(1 - p)}{p} \frac{q}{1 - q}.$$

When considering the tails of $f(q|1)$ and $f(q|0)$, for example, the above means that when $q \rightarrow 0$, assuming that $f(q|0) < \infty$, then $f(q|1) \rightarrow 0$ at a certain rate. Similarly,

$$f(q|0) = f(q|1) \frac{p}{1 - p} \frac{1 - q}{q}$$

so that, assuming that $f(q|1) < \infty$ as $q \rightarrow 1$, then $f(q|0) \rightarrow 0$ at a certain rate. However, given that our approximation implies that $f(q|0)$ drops to zero around the tails very quickly, specifically, as $f(q|0) \rightarrow_{q \rightarrow 1} 0$ faster than $q/(1 - q) \rightarrow_{q \rightarrow 1} \infty$, then we can indeed construct $f(q|1)$ as a probability distribution so that they can jointly satisfy the above Bayesian restriction. □

¹¹These conditions in turn imply Bayesian plausibility, i.e., that $\int_0^1 q(pf(q|1) + (1 - p)f(q|0)) dq = p$.

REFERENCES

- Acemoglu, Daron, Victor Chernozhukov, and Muhamet Yildiz (2016), “Fragility of asymptotic agreement under Bayesian learning.” *Theoretical Economics*, 11, 187–225. [64]
- Arieli, Itai, Jakob Babichenko, and Rann Smorodinsky (2019), “Robust forecast aggregation.” *PNAS*, 115, E12135–E12143. [50]
- Berger, James and L. Mark Berliner (1986), “Robust Bayes and empirical Bayes analysis with ϵ -contaminated priors.” *Annals of Statistics*, 14, 461–486. [51]
- De Filippis, Roberta, Philippe Jehiel, and Antonio Guarino (2017), “Updating ambiguous beliefs in a social learning experiment.” Unpublished Manuscript. [64]
- De Marzo, Peter, Dimitri Vayanos, and Jeffrey Zwiebel (2003), “Persuasion bias, social influence, and unidimensional opinions.” *Quarterly Journal of Economics*, 118, 909–968. [50]
- Ellis, Andrew and Michele Piccione (2017), “Correlation misperception in choice.” *American Economic Review*, 107, 1264–1292. [50]
- Enke, Benjamin and Florian Zimmerman (2019), “Correlation neglect in belief formation.” *Review of Economics Studies*, 86, 313–332. [64]
- Epstein, Larry G. and Martin Schneider (2007), “Learning under ambiguity.” *Review of Economic Studies*, 74, 1275–1303. [51, 52]
- Eyster, Erik and Georg Weizsacker (2011), *Correlation Neglect in Financial Decision-Making*. Discussion Papers of DIW Berlin 1104, SSRN 1735339. [64]
- Fisher, Ronald A. (1912), “On an absolute criterion for fitting frequency curves.” *Messenger of Mathematics*, 41, 155–160. [51]
- Fisher, Ronald A., and Leonard H.C., Tippett, (1928), “Limiting forms of the frequency-distribution of the largest and smallest member of a sample.” *Proceedings of the Cambridge Philosophical Society*, 24, 180–190. [64, 67]
- Gilboa, Itzhak and David Schmeidler (2003), “Inductive inference: An axiomatic approach.” *Econometrica*, 71, 1–26. [51]
- Gilboa, Itzhak and David Schmeidler (2010), “Simplicity and likelihood: An axiomatic approach.” *Journal of Economic Theory*, 145, 1757–1775. [51]
- Glaeser, Edward and Cass R. Sunstein (2009), “Extremism and social learning.” *Journal of Legal Analysis*, 1, 263–324. [50]
- Glatt, Ephraim (2013), “The unanimous verdict according to the talmud: Ancient law providing insight into modern legal theory.” *Pace International Law Review Online Companion*, 3, 316–335. [65]
- Golub, Benjamin and Matthew Jackson (2012), “How homophily affects the speed of learning and best-response dynamics.” *Quarterly Journal of Economics*, 127, 1287–1338. [50]

Good, Irving (1965), *The Estimation of Probabilities: An Essay on Modern Bayesian Methods*. MIT Press, Cambridge, Massachusetts. [51]

Gunn, Lachlan J., Francois Chapeau-Blondeau, Mark D. McDonnell, Bruce R. Davis, Andrew Allison, and Derek Abbott (2016), “Too good to be true: When overwhelming evidence fails to convince.” *Proceedings of the Royal Society A*, 472, 1–15. [65]

Hossain, Tangim and Ryo Okui (2020), “Belief formation under signal correlation.” Unpublished paper, SSRN 3218152. [64]

Joe, Harry (1997), *Multivariate Models and Dependence Concepts*. Chapman & Hall, London. [55]

Kallir, Ido and Doron Sonsino (2009), “The neglect of correlation in allocation decisions.” *Southern Economic Journal*, 75, 1045–1066. [64]

Leadbetter, M. Ross, Georg Lindgren, and Holger Rootze (1983), *Extremes and Related Properties of Random Sequences and Processes*. Springer Verlag, New York. [67]

Levy, Gilat and Ronny Razin (forthcoming), “Combining forecasts in the presence of ambiguity over correlation structures.” *Journal of Economic Theory*, doi: 10.1016/j.jet.2020.105075. [50]

Levy, Gilat and Ronny Razin (2012), “When do simple policies win?” *Economic Theory*, 49, 621–637. [51]

Levy, Gilat and Ronny Razin (2013), “Dynamic legislative decision making when interest groups control the agenda.” *Journal of Economic Theory*, 148, 1862–1890. [51]

Levy, Gilat and Ronny Razin (2015), “Correlation neglect, voting behaviour and information aggregation.” *Journal of Economic Theory*, 105, 1634–1645. [50]

Levy, Gilat and Ronny Razin (2018), “Information diffusion in networks with the Bayesian peer influence heuristic.” *Games and Economic Behaviour*, 109, 262–270. [50]

Ortoleva, Pietro (2012), “Modeling the change of paradigm: Non-Bayesian reactions to unexpected news.” *American Economic Review*, 102, 2410–2436. [51, 52]

Ortoleva, Pietro and Erik Snowberg (2015), “Overconfidence in political behaviour.” *American Economic Review*, 105, 504–535. [64]

Osborne, Martin J., Jeffrey S. Rosenthal, and Matthew A. Turner (2000), “Meetings with costly participation.” *American Economic Review*, 90, 927–943. [51]

Rabin, Matthew and Dimitri Vayanos (2010), “The gambler’s and hot-hand fallacies: Theory and applications.” *Review of Economic Studies*, 77, 730–778. [51]

Robbins, Herbert (1951), “Asymptotically subminimax solutions of compound statistical decision problems.” In *Proceedings of the Second Berkeley Symposium on Mathematical Statistics and Probability*, Vol. 1, 131–148, Berkeley. [51]

Salant, Yuval and Josh Cherry (2020), “Statistical inference in games.” *Unpublished Manuscript*, 88, 1725–1752. [52]

Schacter, L. Daniel, Joanne L. Harbluk, and Donald R. McLachlan (1984), “Retrieval without recollection: An experimental analysis of source amnesia.” *Journal of Verbal Learning and Verbal Behaviour*, 23, 593–611. [62]

Spiegler, Ran (2016), “Bayesian networks and boundedly rational expectations.” *Quarterly Journal of Economics*, 131, 1243–1290. [50]

Suleymanov, Elchin (2018). “Robust maximum likelihood updating.” Unpublished paper. [51]

Co-editor Ran Spiegler handled this manuscript.

Manuscript received 16 September, 2019; final version accepted 22 April, 2020; available online 24 April, 2020.