



Female Genital Mutilation/Cutting across Africa: Dynamics of Change and Socioeconomic Variation

Ewa Batyra, Ben Wilson, Ernestina Coast and Valeria Cetorelli



Stockholm
University

Female Genital Mutilation/Cutting across Africa: Dynamics of Change and Socioeconomic Variation

Ewa Batyra ¹, Ben Wilson ^{2,3*}, Ernestina Coast ⁴, Valeria Cetorelli ⁵

1: Population Studies Center, University of Pennsylvania, 2: Department of Sociology, Stockholm University, 3: Department of Methodology, LSE, 4: Department of International Development, LSE, 5: United Nations Relief and Works Agency

* Corresponding author – ben.wilson@sociology.su.se

Abstract

Background: The majority of women who undergo female genital mutilation/cutting (FGM/C) live in Africa. Although the UN Sustainable Development Goals call for intensified efforts to accelerate the abandonment of FGM/C, little is known about where in Africa the declines in prevalence have been fastest and whether changes in prevalence differ by women's socioeconomic status.

Methods: We use data from Demographic and Health Surveys and Multiple Indicator Cluster Surveys for 23 African countries, collected between 2005 and 2015, and covering 293,170 women. We reconstruct long-term trends in FGM/C prevalence spanning 35 years. We compute absolute and relative rates of change in FGM/C prevalence and differentials in prevalence by women's education and urban-rural residence. We examine whether socioeconomic differences in FGM/C are converging or diverging.

Findings: FGM/C prevalence has declined fastest (in relative terms) in countries with lower initial prevalence, and more slowly in countries with higher initial prevalence. Although better-educated women and those living in urban areas tend to have lower prevalence, in some countries the opposite pattern is observed. Socioeconomic differentials in FGM/C have grown in the majority of countries, particularly in countries with moderate-to-higher overall prevalence.

Conclusions: The documented relationship between absolute and relative FGM/C prevalence rates suggests that in settings with higher initial prevalence, FGM/C practice is likely to be more entrenched and to change more slowly. There is substantial variation between countries in socioeconomic differentials in prevalence and their changes over time. As countries change from higher to lower overall prevalence, socioeconomic inequalities in FGM/C are increasing.

Keywords: FGM/C; Sustainable Development Goals; Africa; socioeconomic inequality; cross-national comparison; cohort dynamics; absolute trends; relative trends; DHS; MICS

The views expressed in this paper are those of the authors and do not necessarily reflect the views of the United Nations.

Stockholm Research Reports in Demography 2020:25

ISSN 2002-617X

© Ewa Batyra, Ben Wilson, Ernestina Coast and Valeria Cetorelli



This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Introduction

More than 200 million women and girls alive today are estimated to have undergone female genital mutilation/cutting (FGM/C), most of whom live in Africa.¹ In 2015, the United Nations General Assembly agreed a series of Sustainable Development Goals (SDGs), including a specific target to eliminate FGM/C by 2030.² FGM/C is defined as “*all procedures involving partial or total removal of the external female genitalia or other injury to the female genital organs for non-medical reasons*” and is classified by severity into four categories.³ Despite socio-cultural variations in FGM/C practices, it is widely recognised as a violation of human rights and harmful to health of women and girls.⁴ The SDGs call for intensified national efforts to accelerate FGM/C elimination, nonetheless, there is limited evidence of change across countries.⁵ Many studies have compared FGM/C prevalence in different African countries^{6–14}, but only a few studies have considered change in FGM/C prevalence rates over time.^{15–20} Little is known about the dynamics of change in FGM/C prevalence between and within African countries.

First, there is little evidence about the rate of change in practice of FGM/C, including research that identifies where declines in prevalence have been fastest. A few studies have compared long-run trends in adult prevalence rates for African countries where FGM/C is concentrated^{17,19,20}, but none of these have estimated both absolute and relative changes in prevalence.

Second, although cross-sectional studies have shown that FGM/C rates differ according to women’s socioeconomic status^{6,7,18,8–15}, there is little cross-national evidence about whether socioeconomic differentials in FGM/C are changing. Prevalence is declining for most types of FGM/C, including substitution of more severe with less severe types.¹⁷ For a limited number of countries, there is evidence that absolute declines in prevalence rates vary considerably by intra-country region, including the Central African Republic and Liberia.¹⁵ However, there is no systematic cross-country comparison of regional trends. Similarly, there is a lack of cross-national evidence to understand whether FGM/C prevalence is changing at different rates by socio-economic factors.

These are important gaps in research. By analysing the dynamics of FGM/C prevalence, we can identify and begin to understand inequalities and – given evidence of declines – the extent to which sub-groups are leading the transition toward low (or lower) prevalence. Accurate estimates of FGM/C dynamics are also vital if evidence-based policies are to be designed and evaluated^{21–23}, as well as in order to ensure the accuracy of projections of future FGM/C prevalence, including assessments of whether the SDG target is likely to be met.

This study compares and contrasts long-term trends in FGM/C prevalence across 23 African countries. It extends understanding by estimating absolute and relative long-term cohort trends in FGM/C prevalence – by education and rural/urban residence - using harmonised data that are comparable across

all 23 countries. We use a completed cohort approach based on estimation of nationally-representative prevalence rates for birth cohorts of adults who are no longer at risk of FGM/C, over 35 years. This approach offers more stable estimates of prevalence by excluding those who may be at future risk of FGM/C and greater insight into underlying socioeconomic dynamics of behaviours and experiences that are more comparable across cohorts. Comparative studies of FGM/C prevalence trends using a cohort perspective^{15,17,19,20} neither consider relative change nor rural/urban or educational dynamics.

Our approach allows us to demonstrate how the dynamics of FGM/C in Africa are changing. We identify countries where FGM/C prevalence is static, rising, or falling, as well as where it has been changing fastest in relative and absolute terms. By exploring the relationship between absolute and relative changes in FGM/C prevalence rates we link our empirical findings to theories that aim to explain FGM/C decline (or lack thereof), specifically the Theory of Social Convention.²⁴ By including the degree of socioeconomic heterogeneity in FGM/C dynamics, we additionally show which groups are leading the decline in FGM/C and provide much-needed evidence for evaluating future changes in FGM/C prevalence.

Methods

Our method included three stages: (1) data harmonisation, (2) estimating trends in absolute and relative FGM/C prevalence for each country by birth cohort, and (3) analysing educational and urban/rural differentials in these trends.

Data Harmonisation

For each country, we either use data from the Demographic and Health Survey (DHS) or the Multiple Indicator Cluster Survey (MICS) (table 1). We focus on countries which had at least one survey including an FGM/C module and where FGM/C practice is non-negligible (national prevalence of at least 5%). Three countries (Cameroon, Nigeria, and Uganda) were excluded from the analysis due to a very small number of women undergoing FGM/C reflecting national prevalence of 1-2%.

Both DHS and MICS are representative of national populations, have similar designs, and are comparable. We harmonised all 23 national surveys to ensure that questions and response codes are similar, and that the data are of sufficient quality for analysis. We use the same approach for each country, thereby facilitating a direct comparison within and between countries over time. During this harmonisation, we corresponded with DHS/MICS support teams and completed quality assurance checks of our derived variables, including those for age and birth cohort, which are essential for our time series estimates validity. Our analyses account for the design of each survey using appropriate survey weights and the 'SVY' command in Stata version 14.²⁵

Trends in absolute and relative FGM/C prevalence by country and birth cohort

We restrict our analysis to women aged 15-49 in order to generate comparable time-series estimates of completed FGM/C prevalence rates. Of those who undergo FGM/C, in all countries in our study, almost all women did so before age 15. Those under 15 at the time of each survey are judged to be still at risk of FGM/C and are excluded from analysis. All the FGM/C modules include a question: “Have you yourself ever been circumcised?” which we use to calculate prevalence, i.e. the percentage of women in a given cohort who report having undergone FGM/C. The potential limitation of such self-reported data is that women may be unwilling to disclose having undergone FGM/C due to the sensitivity of the subject or its illegality in some countries¹⁵, or they may be unaware that they have been cut, especially if that happened at a young age.²⁶ Despite these limitations, such self-reported data are the only source of information of FGM/C that can be used for comprehensive comparative analyses. Table S1 in the Supplementary material shows that missing information about women’s FGM/C status is below 5% in all countries except for Chad (35%) and Kenya (53%). There are no substantial differences in the level of missing values by education level or place of residence in Chad and Kenya. We retain these two countries in our analysis, but the results should be interpreted keeping in mind high levels of missing information on FGM/C status.

We reconstruct historic trends in prevalence using data on year of birth, grouped into five-year birth cohorts, using one cross-sectional survey for each country. The range of cohorts that can be included depends upon the date of each national survey. Since the majority of countries conducted their latest survey around 2010, we use surveys conducted just before or after 2010, even if a more recent survey is available. This is in order to generate trends that are comparable between countries. For most countries, we are able to estimate FGM/C prevalence for women born between 1965-69 and 1995-99. However, the upper end of this range is more restricted for some countries, most notably Burkina Faso, Central African Republic, Djibouti, Eritrea, Ethiopia and Somalia (table 1). We conduct additional analysis using three-year, instead of five-year, birth cohorts to make sure that our results are not sensitive to alternative ways of grouping cohorts. We show these rates in the Supplementary material (table S4).


We then calculate an additional time series (for each country) that enables a cross-national comparison of relative change in FGM/C over time. This is done by indexing aggregate prevalence rates for each cohort to the national prevalence for women born 1965-69 and allows us to examine similarities and differences in the rate of change in FGM/C prevalence between countries.

Analysing educational and urban/rural differentials

Factors influencing the practice of FGM/C are dynamic, heterogeneous and context-specific.¹⁵ To explore the determinants of FGM/C, drivers of change, and dynamics of inequality, we focus on two socioeconomic indicators standardised across all surveys: education and place of residence. These two factors are known to be associated with FGM/C practices^{6-14,18} but are not the only factors. We considered, but were unable, to calculate consistent and comparable trends by other factors including ethnicity and religion. In part due to the unavailability of questions about other factors for all countries, and in part due to the difficulty of cross-national harmonisation of these variables (and their categories).

We calculate the difference in absolute prevalence for: (a) women who have no education versus those who have some education and (b) women who live in rural areas versus those who live in urban areas. Having some education is defined as having completed any years of schooling. We use the classification of urban-rural areas as defined in a given survey for each country, which is based on each country's urban-rural definition at the time of the survey.²⁷ Both education and residence variables are recorded at the time of the survey, and do not capture an individual's characteristics at the time of FGM/C. While this fact has no implications for the analysis of educational differences, it is a potential limitation for the study of urban-rural differences because migration between rural and urban areas may differ by FGM/C status. We are not aware of evidence that indicates whether this is the case, but we acknowledge this limitation and later discuss its possible influence on our results.

Figure 1: Indexed time series of trends in FGM/C prevalence by birth cohort (relative change in FGM/C prevalence). Countries are sorted according to the national FGM/C prevalence rate for 1965-69 (youngest) cohort, which is shown in the parentheses.




Results

In general, there has been a decline in the prevalence of FGM/C, however there is stark variation across countries in long-run trends (figure 1). For women born in 1965-69, prevalence varies from 5% in Ghana (the lowest in our study) to 99% in Guinea (the highest). In some countries, such as Somalia and Mali, there has been no decline in prevalence, and in Gambia there appears to have been a small increase. In Sierra Leone the rate has fallen by 27% for women born thirty years later, reflected in a change in the indexed FGM/C prevalence from 1 to 0.73 between 1965-69 and 1995-99. There is no obvious pattern by region, for example, the neighbouring countries of Gambia, Senegal and Mauritania all exhibit very different trends. Gambia is the only country in our study where FGM/C prevalence increased over the 30-year period; prevalence fell steadily in Mauritania by around 2-3% per year, whereas prevalence in Senegal increased for cohorts born 1970-89, but declined rapidly thereafter.

Alongside Sierra Leone, several other countries in the top two quartiles of prevalence (top two rows of figure 1) exhibit material declines in prevalence of more than 10%: Sudan (11%), Eritrea (15%), Ethiopia (21%), and Burkina Faso (30%). Nonetheless, the pace of decline in these countries is generally slower as compared with countries of below average prevalence (bottom two rows of figure 1). Most of these lower prevalence countries have seen a reduction of more than 30% in FGM/C prevalence across three decades, with the exception of Chad, Senegal and Guinea-Bissau (reductions of 20%, 16% and 8% respectively). Different countries show the 'largest' reduction, depending on whether the reduction is calculated in absolute or relative terms. When comparing those born 1965-69 with the most recent cohorts, the largest absolute declines are for Kenya (28 percentage points), Burkina Faso (27), Sierra Leone (27), Nigeria (23), and Ethiopia (18) (absolute FGM/C prevalence rates for total population are shown in Supplementary material table S2). These are countries with substantial relative declines as well (71%, 30%, 27%, 60% and 21%, respectively). Nonetheless, the largest relative declines over the same period occurred in Tanzania, Benin, Togo and Ghana (declines of 74%, 88%, 83% and 78% respectively).

Figure 2: Relationship between initial FGM/C prevalence (1965-69 cohort) and the rate of change



Note: The percentage change is the change in FGM/C prevalence rates for those born 1965-69 (youngest cohort) as compared with the prevalence rates for the youngest cohort available for each country.


Another pattern that emerges from our analysis of long-term trends is the relationship between ‘initial’ absolute FGM/C prevalence, for those born 1965-69 (index =100), and the relative change in prevalence for subsequent cohorts [$R^2=0.63$]. Countries with the highest prevalence for those born in 1965-69 also exhibit a much smaller relative change in prevalence for cohorts born over the next three decades (figure 2). The largest relative declines are for countries where FGM/C was already a minority practice for the 1965-9 birth cohort.

There is considerable cross-national variation in the magnitude of educational and urban-rural differences in FGM/C prevalence. Figures 3 and 4 show absolute FGM/C prevalence rates by socioeconomic characteristics; countries are ranked by aggregate FGM/C prevalence rate (relative FGM/C prevalence rates by education level and place of residence are shown in the Supplementary material table S2). For women born 1965-69, those with no education have a much higher prevalence in Guinea-Bissau, Gambia, Côte d'Ivoire and Kenya (differentials of more than 15 percentage points as compared with those who have any education) (figure 3). In some countries (Sudan, Nigeria) women with some education from these cohorts have a higher prevalence than women with no education, and the difference is larger than 10 percentage points. This shows that education is not a good predictor of FGM/C cross-nationally, although it is notable that educational differentials among women born in 1965-69 are largest in countries with moderate levels of overall prevalence, and smallest in countries with the highest prevalence.

The heterogeneity that is evident for women born in 1965-69 is amplified when examining how differentials have changed over time. Not only do the trends in differentials move in different directions for different countries, including in divergent ways for educational differentials, but there is also sizable variation in the changing magnitude of differentials. For example, in Central African Republic, the difference in prevalence between women with no education and some education increases from 4 percentage points for the 1965-69 cohort to 17 percentage points for the 1990-95 cohort. By contrast, in Côte d'Ivoire where there is also a sizable differential by education, it is much more stable over time. In Ghana the same educational differential changes in the opposite direction, decreasing from 11 to 2 percentage points.

These socioeconomic differentials can be interpreted as a measure of inequality in FGM/C practices. To this extent, countries can be categorised as becoming more or less equal if differentials become smaller or larger (in absolute terms). Countries trending towards convergence include Gambia, Nigeria, Benin, Togo and Ghana. However, these countries appear to be in the minority as compared with those where inequality is widening, such as Sierra Leone, Burkina Faso, Eritrea, Ethiopia, Guinea-Bissau, Kenya or Central African Republic. Increases in educational differentials occurred in countries that had moderate-to-high levels of initial overall prevalence (i.e. among the 1965-69 cohort). Consequently, for more recent cohorts, educational differentials are largest in countries with moderate-to-high prevalence. It is only among countries with the lowest initial prevalence rates – such as Benin, Togo and Ghana – that educational differentials have consistently reduced.


Figure 3: Absolute FGM/C prevalence rates by education level, percentage point difference in FGM/C prevalence between women with no education and some education, oldest and youngest cohorts



Note: Countries are ranked according to the national FGM/C prevalence rates of the 1965-69 cohorts. The oldest and youngest cohorts as listed in table 1.

The variation in magnitude (and direction) of rural/urban differentials is less pronounced than for education (figure 4). In the majority of countries, women who lived in urban areas at the time of the survey have lower FGM/C prevalence than rural women, although in many cases the difference is small or negligible. For the oldest cohort, there is a much lower urban prevalence (more than 10 percentage points) in Guinea-Bissau, Mauritania and Tanzania. However, the opposite is true in Nigeria, Chad, Sudan, Côte d'Ivoire and Eritrea, where urban prevalence rates are higher (with differentials ranging from 2 to 18 percentage points). As with education, urban/rural differentials tend to be largest in countries with moderate levels of overall prevalence. For the latest cohorts, differentials remain largest in countries with moderate prevalence, but also show an increase in countries with higher overall levels of prevalence (Sierra Leone, Egypt, Eritrea, Sudan, and Burkina Faso). In drawing these conclusions, it is important to note that unlike education, place of residence is recorded at the time of interview, such that urban/rural differentials also reflect patterns of (rural to urban) migration over time. Nevertheless, we frequently observe that differentials between rural and urban areas are increasing.

Figure 4: Absolute FGM/C prevalence rates by place of residence, percentage point difference in FGM/C prevalence between women in rural areas and urban areas, oldest and youngest cohorts



Note: Countries are ranked according to the national FGM/C prevalence rates of the 1965-69 cohorts. The oldest and youngest cohorts as listed in table 1.

Discussion

While FGM/C practices vary substantially across and within countries, a comparison of national trends can illuminate where, to what extent, and how rapidly change is occurring. Our results focus on the dynamics and heterogeneity of FGM/C within and between countries, and show the extent to which the practice is changing across Africa.

The analysis of relative changes in prevalence rates provides a novel perspective of looking at the evolution of the practice of FGM/C and allows us to identify where it has been changing fastest. Despite considerable variation across countries, we uncover a clear relationship between absolute and relative changes in FGM/C prevalence. Relative declines in cohort-specific prevalence rates have been faster in countries that began with low absolute levels of prevalence. These findings provide macro-level evidence in support of the Theory of Social Convention, as applied to FGM/C.²⁴ In countries with lower initial prevalence rates, the mechanisms of social convention – such as marriageability or peer

conventions that support the practice of FGM/C - are less prevalent and less likely to be reinforced because a minority of the population practices FGM/C.²⁸ In settings with higher initial prevalence, these social conventions are more likely to be entrenched and less likely to change.

These macro-level patterns are not deterministic: high prevalence does not mean that high prevalence will persist. With the exception of a few countries such as Somalia, Mali and Gambia, even in the highest prevalence countries FGM/C prevalence has started to decline, albeit at a slow pace. However, we also document that, as in the cases of Senegal and Guinea-Bissau, prevalence is not necessarily decreasing monotonically: initial declines in prevalence do not always lead to subsequent declines.

We show substantial variation between countries in their socioeconomic differentials in FGM/C prevalence. Theories of diffusion suggest that certain socioeconomic groups – such as those with higher education – are more likely to adopt new (non-traditional) behaviours.²⁹ Yet, we show that FGM/C differentials are not in the same direction across countries, and women with some education (or living in urban areas) can have higher average prevalence rates. As noted elsewhere, it is important to realize that different communities practice FGM/C in different social contexts, and that each context presents specific challenges for reformers.³⁰ Interventions towards abandonment of FGM/C must be designed to acknowledge and accommodate the heterogeneity we document, not least with respect to generalisability.

Finally, we show how cross-national variation in socioeconomic differentials change over time. In the past, differentials were smallest in countries where the majority of the population practice FGM/C, and were largest in countries with moderate prevalence levels. However, socioeconomic inequalities in FGM/C have decreased in the lowest prevalence countries and increased in countries with moderate-to-high levels of prevalence. We speculate that these changes are, at least in part, driven by an underlying FGM/C transition, as countries change from high- to low-prevalence – or from FGM/C being a majority to a minority practice. Once a minority of the population practices FGM/C, then it appears that socioeconomic differences in the practice (begin to) disappear, at least at the national-level. As well as adding knowledge about the dynamics of change in FGM/C, these findings can inform research to develop and enhance theories of change, as well as in order to project future levels of FGM/C prevalence.

Overall, this comparative study generates future research agendas by raising questions about why trends and patterns in FGM/C prevalence converge or diverge across countries. Our evidence and analyses enable program directors, policy-makers, researchers, and members of civil society to better understand changes in FGM/C crucial for considering plausible future pathways.

Bibliography

1. UNICEF. Female Genital Mutilation/Cutting: A Global Concern. New York: UNICEF; 2016.
2. UN. Transforming Our World: The 2030 Agenda for Sustainable Development. New York, United Nations; 2015.
3. WHO. Female Genital Mutilation [Internet]. Fact Sheet. 2020 [cited 2020 Feb 16]. Available from: <https://www.who.int/news-room/fact-sheets/detail/female-genital-mutilation>
4. Kimani S, Muteshi J, Njue C. Health Impacts Of Female Genital Mutilation/Cutting: A Synthesis Of The Evidence. Evidence to End FGM/C Programme: Research to Help Girls and Women Thrive. New York: Population Council; 2016.
5. Yoder PS, Wang S, Johansen E. Estimates of female genital mutilation/cutting in 27 African countries and Yemen. *Stud Fam Plann*. 2013 Jun;44(2):189–204.
6. Slinger T, Snow RC, Oronsaye F, Okonofua F, Wacker J. Female genital cutting in southern urban and peri-urban Nigeria: Self-reported validity, social determinants and secular decline. *Trop Med Int Heal*. 2002 Jan;7(1):91–100.
7. El-Gibaly O, Ibrahim B, Mensch BS, Clark WH. The decline of female circumcision in Egypt: evidence and interpretation. *Soc Sci Med*. 2002;54(2):205–20.
8. Hayford SR. Conformity and change: Community effects on female genital cutting in Kenya. *J Health Soc Behav*. 2005;46(June):121–40.
9. Klouman E, Manongi R, Klepp KI. Self-reported and observed female genital cutting in rural Tanzania: Associated demographic factors, HIV and sexually transmitted infections. *Trop Med Int Heal*. 2005 Jan;10(1):105–15.
10. Satti A, Elmusharaf S, Bedri H, Idris T, Hashim MSK, Suliman GI, et al. Prevalence and determinants of the practice of genital mutilation of girls in Khartoum, Sudan. *Ann Trop Paediatr*. 2006;26:303–10.
11. Ouldzeidoune N, Keating J, Bertrand J, Rice J. A Description of Female Genital Mutilation and Force-Feeding Practices in Mauritania: Implications for the Protection of Child Rights and Health. *PLoS One*. 2013;8(4):1–9.
12. Chikhungu LC, Madise NJ. Trends and protective factors of female genital mutilation in Burkina Faso: 1999 to 2010. *Int J Equity Health*. 2015;14:1–10.
13. Sipsma HL, Chen PG, Ofori-Atta A, Ilozumba UO, Karfo K, Bradley EH. Female genital cutting: Current practices and beliefs in western Africa. *Bull World Health Organ*. 2012;90(2):120–7.
14. Yount KM. Like Mother, like Daughter? Female Genital Cutting in Minia, Egypt. *J Health Soc Behav*. 2002;43(3):336–58.
15. UNICEF. Female Genital Mutilation/Cutting: A Statistical Overview and Exploration of the Dynamics of Change. UNICEF, New York; 2013.
16. Kandala NB, Ezejimofor MC, Uthman OA, Komba P. Secular trends in the prevalence of female genital mutilation/cutting among girls: a systematic analysis. *BMJ Glob Heal*. 2018;3(5):1–7.

17. Koski A, Heymann J. Thirty-year trends in the prevalence and severity of female genital mutilation: A comparison of 22 countries. *BMJ Glob Heal.* 2017;2:1–8.
18. Andro A, Lesclingand M. Female genital mutilation. Overview and current knowledge. *Population (Paris).* 2016;71(2):224–311.
19. Engelsma B, Mackie G, Marrella B. Unprogrammed abandonment of female genital mutilation cutting. *World Dev.* 2020;129.
20. Koski A, Heymann J. Changes in support for the continuation of female genital mutilation/cutting and religious views on the practice in 19 countries. *Glob Public Health.* 2019;14(5):696–708.
21. Muthumbi J, Svanemyr J, Scolaro E, Temmerman M, Say L. Female Genital Mutilation: A Literature Review of the Current Status of Legislation and Policies in 27 African Countries and Yemen. *Afr J Reprod Health.* 2015;19(3):32–40.
22. Shell-Duncan B, Wander K, Hernlund Y, Moreau A. Legislating change? Responses to criminalizing female genital cutting in Senegal. *Law Soc Rev.* 2013;47(4):803–35.
23. Cetorelli V, Wilson B, Batyra E, Coast E. Female Genital Mutilation/Cutting in Mali and Mauritania: Understanding Trends and Evaluating Policies. *Stud Fam Plann.* 2020;51(1):51–69.
24. Shell-Duncan B, Wander K, Hernlund Y, Moreau A. Dynamics of change in the practice of female genital cutting in Senegambia: Testing predictions of social convention theory. *Soc Sci Med.* 2011;73(8):1275–83.
25. StataCorp. *Stata Statistical Software: Release 14.* College Station, TX: StataCorp LP. College Station, TX: StataCorp LP: College Station, TX: StataCorp LP; 2015.
26. Jackson EF, Akweongo P, Sakeah E, Hodgson A, Asuru R, Phillips JF. Inconsistent reporting of female genital cutting status in northern Ghana: Explanatory factors and analytical consequences. *Stud Fam Plann.* 2003;34(3):200–10.
27. DHS. Demographic and Health Survey Forum [Internet]. 2020. [cited 2020 Feb 16]. Available from: <https://userforum.dhsprogram.com/index.php?t=msg&th=55&start=0&>
28. Mackie G. Ending Footbinding and Infibulation: A Convention Account. *Am Sociol Rev.* 1996;61(6):999–1017.
29. Bongaarts J. Completing the fertility transition in the developing world: The role of educational differences and fertility preferences. *Popul Stud (NY).* 2003;57(3):321–35.
30. Hernlund Y, Shell-Duncan B. Transcultural Positions: Negotiating Rights and Culture. In: Hernlund Y, Shell-Duncan B, editors. *Transcultural Bodies: Female Genital Cutting in Global Context.* New Brunswick, NJ: Rutgers University Press; 2007. p. 1–45.

Table 1: Countries, data sources and sample sizes

	Country Abbr.	Survey Type	Survey Year	Oldest Cohort	Youngest Cohort	Sample Size
Benin	BJ	DHS	2011/12	1965-69	1995-97	16,152
Burkina Faso	BF	DHS	2010	1965-69	1990-94	15,430
Central African Republic	CF	MICS	2010	1965-69	1990-94	10,562
Chad	TD	DHS	2014/15	1965-69	1995-99	11,402
Cote d'Ivoire	CI	DHS	2011/12	1965-69	1995-97	9,708
Djibouti	DJ	MICS	2006	1965-69	1990-91	5,471
Egypt	EG	DHS	2014	1965-69	1995-99	21,441
Eritrea	ER	DHS	2002	1965-69	1985-87	6,659
Ethiopia	ET	DHS	2005	1965-69	1985-89	11,367
Gambia	GM	DHS	2013	1965-69	1995-98	10,060
Ghana	GH	MICS	2011	1965-69	1995-96	9,992
Guinea	GN	DHS	2012	1965-69	1995-97	8,852
Guinea-Bissau	GW	MICS	2014	1965-69	1995-99	10,193
Kenya	KE	DHS	2014	1965-69	1995-99	14,682
Mali	ML	DHS	2012/13	1965-69	1995-97	10,259
Mauritania	MR	MICS	2015	1965-69	1995-99	13,612
Nigeria	NG	DHS	2013	1965-69	1995-98	35,983
Senegal	SN	DHS	2014	1965-69	1995-99	8,453
Sierra Leone	SL	DHS	2013	1965-69	1995-98	16,371
Somalia	SO	MICS	2006	1965-69	1990-91	6,241
Sudan	SD	MICS	2014	1965-69	1995-99	18,292
Tanzania	TZ	DHS	2015/16	1965-69	1995-99	12,619
Togo	TG	DHS	2013/14	1965-69	1995-99	9,369
<i>Total</i>						<i>293,170</i>

Note: DHS- Demographic and Health Survey, MICS- Multiple Indicator Cluster Survey

Supplementary material

Table S1: Per cent missing values for information about women's FGM/C status, by country, for total population and according to women's socioeconomic characteristics

	total population	some education	no education	urban	rural
Benin	0.0	0.0	0.0	0.0	0.0
Burkina Faso	0.1	0.0	0.1	0.0	0.1
Central African Republic	0.2	0.3	0.2	0.3	0.2
Chad	35.1	34.6	35.4	36.5	34.7
Cote d'Ivoire	0.1	0.1	0.1	0.1	0.1
Djibouti	0.3	0.1	0.4	0.3	0.4
Egypt	0.0	0.0	0.0	0.1	0.0
Eritrea	0.0	0.1	0.0	0.1	0.0
Ethiopia	3.1	3.9	2.5	5.1	2.2
Gambia	0.3	0.3	0.2	0.3	0.2
Ghana	0.0	0.0	0.0	0.0	0.0
Guinea	0.1	0.1	0.1	0.2	0.0
Guinea-Bissau	0.0	0.0	0.0	0.0	0.0
Kenya	52.6	52.6	52.7	52.9	52.4
Mali	0.0	0.0	0.0	0.0	0.0
Mauritania	1.1	1.3	0.7	1.2	1.1
Nigeria	4.5	3.4	6.5	4.9	4.2
Senegal	0.0	0.0	0.0	0.0	0.0
Sierra Leone	0.1	0.1	0.1	0.1	0.1
Somalia	0.1	0.2	0.1	0.3	0.1
Sudan	0.1	0.1	0.1	0.0	0.1
Tanzania	0.0	0.0	0.0	0.0	0.0
Togo	0.1	0.1	0.1	0.1	0.1

Table S2: FGM/C prevalence for total population for the oldest and youngest cohorts (absolute rates) and FGM/C prevalence for the youngest cohort indexed at 1 according to 1965-69 rate (relative rates) by education level and place of residence

	total population, oldest cohort (absolute rate)	total population, youngest cohort (absolute rate)	some education, youngest cohort, (relative rate)	no education, youngest cohort, (relative rate)	urban, youngest cohort, (relative rate)	rural, youngest cohort, (relative rate)
Guinea	0.99	0.94	0.94	0.97	0.97	0.93
Somalia	0.99	0.98	0.98	1.01	1.00	0.99
Sierra Leone	0.98	0.71	0.74	0.83	0.64	0.79
Egypt	0.95	0.88	0.96	0.81	0.86	0.91
Djibouti	0.93	0.88	0.92	0.99	0.94	0.95
Eritrea	0.93	0.79	0.83	1.00	0.77	0.91
Sudan	0.92	0.82	0.86	0.75	0.80	0.93
Mali	0.92	0.90	0.99	0.98	1.00	0.98
Burkina Faso	0.89	0.62	0.61	0.78	0.63	0.74
Ethiopia	0.84	0.66	0.70	0.87	0.74	0.80
Gambia	0.75	0.77	1.37	0.93	1.06	1.00
Mauritania	0.73	0.64	0.92	0.93	0.82	0.93
Guinea-Bissau	0.46	0.42	1.25	1.49	0.98	0.93
Cote d'Ivoire	0.45	0.29	0.59	0.86	0.64	0.64
Chad	0.40	0.32	0.75	0.96	0.70	0.83
Kenya	0.40	0.12	0.26	1.53	0.27	0.30
Nigeria	0.38	0.15	0.34	0.48	0.36	0.46
C.A. Republic	0.31	0.20	0.48	0.92	0.43	0.75
Senegal	0.25	0.21	1.03	0.69	0.75	0.90
Tanzania	0.20	0.05	0.25	0.53	0.24	0.28
Benin	0.11	0.01	0.11	0.23	0.07	0.15
Togo	0.10	0.02	0.26	0.30	0.16	0.18
Ghana	0.05	0.01	0.53	0.25	0.09	0.26

Note: Countries are ranked according to the national FGM/C prevalence rates of the 1965-69 cohorts. The oldest and youngest cohorts as listed in table 1. Relative rates by education level and place of residence for the oldest cohort are equal to 1.

Table S3: Per cent of women with some education and living in urban areas, oldest and youngest cohorts (%)

Country /Cohort	% Some Education		% Urban	
	Oldest	Youngest	Oldest	Youngest
Benin	28.8	73.9	45.5	49.1
Burkina Faso	13.5	48.5	22.5	25.2
Central African Republic	54.6	65.3	39.1	40.7
Chad	20.8	55.2	20.3	26.4
Cote d'Ivoire	36.3	60.7	45.6	59.3
Djibouti	29.2	77.2	96.8	97.5
Egypt	55.9	90.6	42.4	17.7
Eritrea	34.9	84.1	43.4	47.0
Ethiopia	19.6	54.4	14.0	21.3
Gambia	23.8	76.0	51.7	52.6
Ghana	69.4	97.2	55.8	48.6
Guinea	14.3	60.0	27.8	40.3
Guinea-Bissau	32.8	85.3	45.0	53.1
Kenya	87.7	97.6	29.8	30.3
Mali	13.7	48.9	19.1	31.9
Mauritania	60.1	86.4	49.7	49.0
Nigeria	52.8	71.7	41.9	41.5
Senegal	31.2	69.9	54.8	50.0
Sierra Leone	20.1	81.7	30.4	40.3
Somalia	31.2	57.0	32.2	45.3
Sudan	46.9	85.9	34.9	32.5
Tanzania	81.4	93.5	28.7	37.9
Togo	47.6	89.4	35.1	44.5

Note: The oldest and youngest cohorts as listed in table 1.

Table S4: FGM/C prevalence by 3-year cohorts, total population (T), by education level (SE, NE) and place of residence (U, R), absolute rates (A) for the oldest (O) and the youngest (Y) cohorts, relative rates (R) for the youngest cohort

	Oldest cohort (O)	Youngest cohort (Y)	T-O-A	T-Y-A	T-Y-R	SE-O-A	SE-Y-A	SE-Y-R	NE-O-A	NE-Y-A	NE-Y-R	U-O-A	U-Y-A	U-Y-R	R-O-A	R-Y-A	R-Y-R
Guinea	1965-67	1995-97	1.00	0.94	0.94	0.98	0.92	0.93	1.00	0.97	0.97	0.98	0.95	0.96	1.00	0.93	0.93
Somalia	1965-67	1989-91	0.98	0.97	0.99	0.98	0.95	0.97	0.98	0.98	1.01	0.96	0.96	0.99	0.99	0.97	0.98
Sierra Leone	1965-67	1995-97	0.98	0.74	0.75	0.91	0.71	0.78	0.99	0.84	0.85	0.93	0.63	0.68	0.99	0.81	0.82
Egypt	1965-67	1995-97	0.95	0.89	0.93	0.93	0.90	0.97	0.98	0.80	0.82	0.91	0.79	0.87	0.99	0.91	0.93
Sudan	1965-67	1995-97	0.94	0.82	0.88	0.98	0.85	0.87	0.89	0.66	0.74	0.95	0.77	0.81	0.93	0.85	0.92
Eritrea	1965-67	1983-85	0.93	0.79	0.84	0.91	0.77	0.84	0.95	0.86	0.90	0.92	0.74	0.80	0.94	0.82	0.87
Djibouti	1965-67	1989-91	0.93	0.89	0.95	0.95	0.88	0.92	0.92	0.91	0.99	0.93	0.89	0.96	0.96	0.90	0.94
Mali	1965-67	1995-97	0.92	0.90	0.98	0.91	0.89	0.98	0.92	0.91	0.99	0.90	0.90	0.99	0.92	0.90	0.98
Burkina Faso	1965-67	1992-94	0.88	0.58	0.66	0.82	0.47	0.57	0.89	0.67	0.76	0.87	0.52	0.60	0.88	0.61	0.69
Ethiopia	1965-67	1986-88	0.83	0.65	0.78	0.87	0.62	0.71	0.83	0.70	0.84	0.87	0.61	0.70	0.83	0.67	0.81
Gambia	1965-67	1995-97	0.74	0.77	1.05	0.52	0.78	1.51	0.80	0.74	0.92	0.68	0.74	1.08	0.79	0.81	1.02
Mauritania	1965-67	1995-97	0.74	0.64	0.86	0.68	0.62	0.92	0.80	0.70	0.87	0.60	0.49	0.82	0.87	0.78	0.90
Cote d'Ivoire	1965-67	1995-97	0.46	0.29	0.64	0.37	0.18	0.49	0.50	0.46	0.92	0.52	0.30	0.58	0.40	0.28	0.70
Guinea-Bissau	1965-67	1995-97	0.42	0.43	1.01	0.28	0.36	1.29	0.49	0.80	1.63	0.33	0.37	1.14	0.50	0.50	1.00
Kenya	1965-67	1995-97	0.40	0.12	0.29	0.38	0.10	0.25	0.49	0.89	1.81	0.31	0.09	0.30	0.43	0.13	0.29
Nigeria	1965-67	1995-97	0.37	0.16	0.42	0.48	0.17	0.35	0.26	0.12	0.47	0.48	0.17	0.35	0.30	0.14	0.49
Chad	1965-67	1995-97	0.35	0.33	0.94	0.28	0.25	0.90	0.37	0.43	1.14	0.36	0.32	0.89	0.35	0.34	0.96
C.A. Republic	1965-67	1992-94	0.33	0.18	0.54	0.31	0.13	0.41	0.35	0.29	0.82	0.26	0.10	0.40	0.37	0.23	0.61
Senegal	1965-67	1995-97	0.24	0.21	0.88	0.20	0.22	1.11	0.26	0.19	0.73	0.23	0.18	0.80	0.27	0.25	0.92
Tanzania	1965-67	1995-97	0.19	0.05	0.29	0.14	0.04	0.31	0.34	0.17	0.51	0.05	0.02	0.33	0.24	0.08	0.32
Benin	1965-67	1995-97	0.12	0.01	0.10	0.05	0.01	0.14	0.15	0.03	0.19	0.09	0.01	0.06	0.14	0.02	0.13
Togo	1965-67	1995-97	0.10	0.02	0.21	0.06	0.02	0.31	0.13	0.05	0.34	0.10	0.02	0.20	0.09	0.02	0.22
Ghana	1965-67	1992-94	0.05	0.02	0.37	0.02	0.01	0.63	0.12	0.12	1.04	0.02	0.01	0.59	0.08	0.02	0.28

Note: O-oldest cohort, Y-youngest cohort, T-total population, SE-some education, NE-no education, U-urban, R-rural, A-absolute rate, R-relative rate.

Countries are ranked according to the national FGM/C prevalence rates of the 1965-67 cohorts. Relative rates for the oldest cohort are equal to 1.

Acknowledgements

The authors are grateful for financial support from STICERD (LSE).

Stockholm Research Reports in Demography
Stockholm University,
106 91 Stockholm,
Sweden
www.su.se | info@su.se | ISSN 2002-617X



Stockholm
University

Demography Unit