



# DISCOUNTED OPTIMAL STOPPING ZERO-SUM GAMES IN DIFFUSION TYPE MODELS WITH MAXIMA AND MINIMA

PAVEL V. GAPEEV,\* *London School of Economics and Political Science*

## Abstract

We present a closed-form solution to a discounted optimal stopping zero-sum game in a model based on a generalised geometric Brownian motion with coefficients depending on its running maximum and minimum processes. The optimal stopping times forming a Nash equilibrium are shown to be the first times at which the original process hits certain boundaries depending on the running values of the associated maximum and minimum processes. The proof is based on the reduction of the original game to the equivalent coupled free-boundary problem and the solution of the latter problem by means of the smooth-fit and normal-reflection conditions. We show that the optimal stopping boundaries are partially determined as either unique solutions to the appropriate system of arithmetic equations or unique solutions to the appropriate first-order nonlinear ordinary differential equations. The results obtained are related to the valuation of the perpetual lookback game options with floating strikes in the appropriate diffusion-type extension of the Black–Merton–Scholes model.

*Keywords:* Three-dimensional optimal stopping zero-sum game; Brownian motion; running maximum and minimum processes; free-boundary problem; instantaneous stopping and smooth fit; normal reflection; a change-of-variable formula with local time on surfaces; perpetual American lookback game options

2020 Mathematics Subject Classification: Primary 60G40; 34B40; 91G20  
Secondary 60J60; 60J65; 91B70

## 1. Formulation of the problem

For a precise formulation of the problem, let us consider a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  with a standard Brownian motion  $B = (B_t)_{t \geq 0}$ . Suppose that the process  $X = (X_t)_{t \geq 0}$  is given by

$$X_t = x \exp \left( \int_0^t \left( r - \delta(S_u, Q_u) - \frac{\sigma^2(S_u, Q_u)}{2} \right) du + \int_0^t \sigma(S_u, Q_u) dB_u \right), \quad (1.1)$$

so that it solves the stochastic differential equation

$$dX_t = (r - \delta(S_t, Q_t)) X_t dt + \sigma(S_t, Q_t) X_t dB_t \quad (X_0 = x), \quad (1.2)$$

where  $r > 0$  is a given constant,  $\delta(s, q) > 0$  and  $\sigma(s, q) > 0$  are continuously differentiable bounded functions on  $[0, \infty]^2$ , and  $x > 0$  is fixed. We further assume that the function  $\delta(s, q)$

Received 10 February 2021; accepted 7 June 2024.

\* Postal address: London School of Economics, Department of Mathematics, Houghton Street, London WC2A 2AE, United Kingdom. Email address: [p.v.gapeev@lse.ac.uk](mailto:p.v.gapeev@lse.ac.uk)

© The Author(s), 2024. Published by Cambridge University Press on behalf of Applied Probability Trust. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

is increasing in both variables  $s$  and  $q$  on  $[0, \infty]^2$ . Here,  $S = (S_t)_{t \geq 0}$  and  $Q = (Q_t)_{t \geq 0}$  are the associated with the *running maximum* and *minimum* processes of  $X$ , respectively defined by

$$S_t = s \vee \max_{0 \leq u \leq t} X_u \quad \text{and} \quad Q_t = q \wedge \min_{0 \leq u \leq t} X_u \tag{1.3}$$

for arbitrary  $0 < q \leq x \leq s$ . Since the functions  $\delta(s, q)$  and  $\sigma(s, q)$  are assumed to be bounded on  $[0, \infty]^2$ , it follows from [48, Chapter IV, Theorem 4.8] that there exists a (pathwise) unique strong solution to the stochastic differential equation in (1.2). It can be assumed that the process  $X$  describes the price of a risky asset on a financial market, where  $r$  is the riskless interest rate,  $\delta(s, q)$  is the dividend rate paid to the asset holders, and  $\sigma(s, q)$  is the volatility rate.

The main aim of this paper is to present closed-form solutions to a discounted optimal stopping zero-sum game with the value

$$V_* = \sup_{\tau} \inf_{\zeta} \mathbb{E} \left[ e^{-r\tau} F(X_\tau, S_\tau) I(\tau < \zeta) + e^{-r\zeta} G(X_\zeta, S_\zeta, Q_\zeta) I(\zeta \leq \tau) \right], \tag{1.4}$$

where we set

$$F(x, s) = s - Kx \quad \text{and} \quad G(x, s, q) = (s - Kx) \vee (q - Lx) \equiv \max\{s - Kx, q - Lx\} \tag{1.5}$$

for all  $0 < q \leq x \leq s$  and some given constants  $0 < L < K < L + 1$ , while  $I(\cdot)$  denotes the indicator function. Suppose that the supremum and infimum in (1.4) are taken over all stopping times  $\tau$  and  $\zeta$  of the process  $X$ , and the expectation there is taken with respect to the risk-neutral probability measure  $\mathbb{P}$ . In that case, the value of (1.4) can be interpreted as the rational (or no-arbitrage) price of a perpetual lookback game (or Israeli) option with the floating strikes  $KX$  and  $LX$  in the diffusion-type extension of the Black–Merton–Scholes model considered in Gapeev and Rodosthenous [30]–[32]. Such game-type contingent claims were introduced by Kifer [41] and further studied by Kyprianou [45], Kühn and Kyprianou [44], Kallsen and Kühn [39], Baurdoux and Kyprianou [3]–[5], Gapeev and Kühn [25], Ekström and Villeneuve [15], Ekström and Peskir [14], Peskir [55]–[56], and Baurdoux *et al.* [6], among others. We also refer to Shiryaev [68, Chapter VIII, Section 2a], Peskir and Shiryaev [59, Chapter VII, Section 25], and Detemple [12] for extensive overviews of the solutions to the American option pricing problems as well as other related results on optimal stopping problems in financial mathematics.

The study of discounted optimal stopping problems for certain reward functionals depending on the running maxima and minima of continuous Markov (diffusion-type) processes was initiated by Shepp and Shiryaev [64] and further developed by Pedersen [51], Guo and Shepp [36], Peskir [53], Gapeev [19]–[20], Guo and Zervos [37], Peskir [57]–[58], Glover *et al.* [33], Gapeev and Rodosthenous [30]–[32], Kitapbayev [42], Rodosthenous and Zervos [63], Gapeev *et al.* [23], Gapeev and Li [27]–[28], Gapeev and Al Motairi [29], Gapeev *et al.* [24], and Gapeev [22], among others. The main feature in the analysis of such optimal stopping problems was that the normal-reflection conditions hold for the value functions at the diagonal planes of the state spaces of the multi-dimensional continuous Markov processes having the initial processes and the running extrema as their components. It was shown, by using the maximality principle established by Peskir [52] for solutions of optimal stopping problems for maxima of the original diffusion processes, which is equivalent to the superharmonic characterisation of the value functions, that the optimal stopping boundaries for the original processes represent functions of the running values of the associated maxima processes and are characterised by

the appropriate extremal solutions of certain first-order nonlinear ordinary differential equations. In this paper, we continue these developments and study the problem of (1.4) related to the pricing of the floating-strike lookback game options as the associated optimal stopping zero-sum game of (2.3) for a three-dimensional (continuous) Markov diffusion-type process which has the underlying risky asset price  $X$  as well as its running maximum  $S$  and minimum  $Q$  as their state-space components.

Note that the resulting problems turn out to be necessarily three-dimensional, in the sense that they cannot be reduced to optimal stopping problems for Markov processes of lower dimensions. It is shown that the optimal exercise times forming a Nash equilibrium are the first times at which the original process exits certain two-sided regions restricted by stochastic boundaries depending on the running values of the associated maximum and minimum processes. We apply the smooth-fit and normal-reflection conditions for the value functions to determine the optimal stopping boundaries as either unique solutions to the appropriate system of arithmetic equations or unique solutions to the appropriate first-order nonlinear ordinary differential equations. Optimal stopping problems with the one-sided continuation regions in similar models based on the original diffusion-type processes with coefficients depending on the running maximum and the running maximum drawdown were considered in Gapeev and Rodosthenous [30]–[32]. Other optimal stopping problems in models with spectrally negative Lévy processes and their running maxima were studied by Asmussen *et al.* [1], Avram *et al.* [2], Ott [50], and Kyrianiou and Ott [46], among others.

The dependence of the local drift and diffusion coefficients on the past dynamics of observable diffusion-type processes through certain processes playing the role of sufficient statistics is often used in financial practice and is well studied in the related literature. For instance, an increase of the running maximum or decrease of the running minimum of a risky asset price normally causes a structural change in the local drift representing its expected return and dividend policy. It also triggers changes in the diffusion coefficient representing the volatility rate of an asset price with a higher impact under either a maximum increase or a minimum decrease, rather than either a minimum increase or maximum decrease, respectively. Such sufficient statistics transparently exhibit the risk levels of the assets and therefore usually influence the decisions taken by market participants. The demand for option pricing in models with stochastic interest rates and volatility initiated the development and subsequent calibration of these models, based on diffusion-type processes with tractable path-dependent coefficients, which were realised by Henry-Labordère [38] and Ren *et al.* [61], among others (see also [31] for further discussion of diffusion-type models for prices of financial assets with coefficients depending on the running maxima and minima as well as the maxima drawdowns and maxima drawups).

This paper is organised as follows. In Section 2, we formulate the optimal stopping zero-sum game for a necessarily three-dimensional continuous Markov process, which has the underlying asset price and the running values of its maximum and minimum as the state-space components. The resulting optimal stopping game is reduced to the equivalent coupled free-boundary problem for the value function which satisfies the smooth-fit conditions at the stopping boundaries and the normal-reflection conditions at the edges of the state space of the three-dimensional process. In Section 3, we obtain closed-form expressions for the candidate value functions, and we derive the appropriate arithmetic equations and first-order nonlinear ordinary differential equations for the candidate stopping boundaries as solutions to the associated free-boundary problems. We specify the starting conditions for the solutions to the first-order nonlinear ordinary differential equations and provide a recursive algorithm to

determine the value functions and the optimal stopping boundaries, along with their lines of intersection with the edges of the three-dimensional state space. In Section 4, by applying change-of-variable formula with local time on surfaces from Peskir [54], we verify that the resulting solution to the free-boundary problem provides the expressions for the value function and the optimal stopping boundaries for the underlying asset price process in the original problem. In Section 5, we give closed-form solutions to some auxiliary optimal stopping problems in the same model, which give the appropriate bounds for the value functions and optimal stopping boundaries for the original game. We apply the maximality principle from Peskir [52] to the framework of the three-dimensional optimal stopping problem under consideration to show that the optimal stopping boundaries provide the extremal solutions of the associated first-order nonlinear ordinary differential equations (see also [30, 32, 58] for optimal stopping problems in other related three-dimensional models). The main results of the paper are stated in Theorems 1 and 2.

## 2. The optimal stopping game and free-boundary problem

In this section, we introduce the setting and notation for the three-dimensional optimal stopping zero-sum game associated with the value of (1.4), which is related to the pricing of the perpetual floating-strike lookback game options. We specify the structure of the optimal stopping times forming a Nash equilibrium and formulate the equivalent free-boundary problem.

### 2.1. The three-dimensional optimal stopping zero-sum game

Suppose that an investor writes a perpetual lookback game option and sells the contract to another investor at time 0. The holder of the option can then exercise the contract at some random time  $\tau$ , which they can choose, by collecting the amount of the running maximum  $S$  and paying the floating strike  $KX$  to the writer, for some  $K > 0$ . At the same time, the writer of the option can either recall the contract at some random time  $\zeta$ , which they choose, by paying the amount of the running minimum  $Q$  to the holder and collecting the floating strike  $LX$ , for some  $L > 0$ , when  $Q - LX > S - KX$  holds, or agree with the holder on the payment of  $S$  and the collection of  $KX$ . Consequently, the holder of the option looks for the exercise time  $\tau_*$  maximising the expected total payoff received from the writer, while, at the same time, the writer of the contract looks for the recall time  $\zeta_*$  minimising the expected total payoff sent to the holder. In other words, the perpetual lookback game option pricing problem seeks to determine the pair of stopping times  $\tau_*$  and  $\zeta_*$  of the process  $X$  that corresponds to a *saddle point* for the total expected reward functional given by

$$\mathbb{J}(\tau, \zeta) = \mathbb{E}\left[e^{-r\tau} F(X_\tau, S_\tau) I(\tau < \zeta) + e^{-r\zeta} G(X_\zeta, S_\zeta, Q_\zeta) I(\zeta \leq \tau)\right], \quad (2.1)$$

with the functions  $F(x, s)$  and  $G(x, s, q)$  defined in (1.5), for some  $0 < L < K < L + 1$  fixed, which means that the inequalities

$$\mathbb{J}(\tau, \zeta_*) \leq \mathbb{J}(\tau_*, \zeta_*) \leq \mathbb{J}(\tau_*, \zeta) \quad (2.2)$$

should hold, for any exercise and recall times  $\tau$  and  $\zeta$ . Such a pair  $\tau_*$  and  $\zeta_*$  satisfying the inequalities of (2.2) with the functional defined in (2.1) is called a *Nash equilibrium* in the optimal stopping zero-sum game of (1.4) (see e.g. [8, 14, 55] for a precise definition of this notion).

It thus follows from the results of [39, 41] that the rational (or no-arbitrage) price of the game contingent claim described above coincides with the value function  $V_*(x, s, q)$  of the optimal stopping zero-sum game for the (time-homogeneous strong) Markov process  $(X, S, Q) = (X_t, S_t, Q_t)_{t \geq 0}$  of the form

$$V_*(x, s, q) = \sup_{\tau} \inf_{\zeta} \mathbb{E}_{x,s,q} [e^{-r\tau} F(X_\tau, S_\tau) I(\tau < \zeta) + e^{-r\zeta} G(X_\zeta, S_\zeta, Q_\zeta) I(\zeta \leq \tau)], \quad (2.3)$$

where the supremum and infimum are taken over all stopping times  $\tau$  and  $\zeta$  with respect to the natural filtration  $(\mathcal{F}_t)_{t \geq 0}$  of the process  $X$ , and the functions  $F(x, s)$  and  $G(x, s, q)$  are given by (1.5), for some  $0 < L < K < L + 1$  fixed. Here, we denote by  $\mathbb{E}_{x,s,q}$  the expectation with respect to the probability measure  $\mathbb{P}$  under the assumption that the three-dimensional (strong Markov) process  $(X, S, Q)$  defined in (1.1)–(1.2) and (1.3) starts at  $(x, s, q) \in E$ , and by  $E = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < q \leq x \leq s\}$  the state space of the process  $(X, S, Q)$ . It therefore follows from the results of [11, Theorem 4.1], based on the solutions of the associated (doubly) reflected backward stochastic differential equations, that the game-type optimal stopping problem of (2.3) has a value. The existence of the associated *Stackelberg equilibria* in various optimal stopping games is proved in the results of [47, 55–56, 69–70], among others. We further establish the existence and describe the structure of the stopping times  $\tau_*$  and  $\zeta_*$  forming a *Nash equilibrium* of the optimal stopping zero-sum game of (2.3).

### 2.2. The structure of the optimal stopping times

Let us first determine the structure of the stopping times forming a Nash equilibrium in the optimal stopping game of (2.3).

(i) By means of the results of general optimal stopping theory for Markov processes (see e.g. [59, Chapter I, Section 2.2]) and the results of the general theory of optimal stopping games (see e.g. [7–8, 11, 16–17, 43, 47, 69], among others), we obtain from the structure of the reward functional that the stopping times forming a Nash equilibrium in the optimal stopping game of (2.3) exist and are given by

$$\tau_* = \inf \{t \geq 0 \mid V_*(X_t, S_t, Q_t) = F(X_t, S_t)\} \quad (2.4)$$

and

$$\zeta_* = \inf \{t \geq 0 \mid V_*(X_t, S_t, Q_t) = G(X_t, S_t, Q_t)\}, \quad (2.5)$$

so that the associated continuation and stopping regions have the forms

$$C_* = \{(x, s, q) \in E \mid F(x, s) < V_*(x, s, q) < G(x, s, q)\} \quad (2.6)$$

and

$$D_* = \{(x, s, q) \in E \mid \text{either } V_*(x, s, q) = F(x, s) \text{ or } V_*(x, s, q) = G(x, s, q)\} \quad (2.7)$$

respectively, where the functions  $F(x, s)$  and  $G(x, s, q)$  are given by (1.5), for some  $0 < L < K < L + 1$  fixed. It can be seen from the results of Theorem 1 below that the value function  $V_*(x, s, q)$  is continuous, so that the set  $C_*$  is open and the set  $D_*$  is closed.

It follows from the structure of the payoff functions  $F(x, s)$  and  $G(x, s, q)$  given by (1.5) that the inequality  $F(X_t, S_t) < G(X_t, S_t, Q_t)$  holds when  $X_t > (S_t - Q_t)/(K - L)$ , while the equality  $F(X_t, S_t) = G(X_t, S_t, Q_t)$  holds when  $X_t \leq (S_t - Q_t)/(K - L)$ , for any  $t \geq 0$ . Moreover, by virtue

of the structure of the processes  $S$  and  $Q$  in (1.3), we may conclude that the inequalities  $S_t - KX_t \geq (1 - K)X_t \geq Q_t - LX_t$  hold when  $X_t \geq Q_t/(L - K + 1)$ , because of the assumption  $0 < L < K < L + 1$ , so that the equality  $F(X_t, S_t) = G(X_t, S_t, Q_t)$  is also satisfied in that case, for any  $t \geq 0$ . Observe that the latter condition in particular holds in the case  $0 < L < K \leq 1$ , which is more restrictive but allows us to keep both the payoffs  $F(X_t, S_t)$  and  $G(X_t, S_t, Q_t)$  positive, for all  $t \geq 0$ , which also represents an important feature from the point of view of callable financial contracts. Note that, in the case  $L \geq K$ , the inequality  $S_t - KX_t < Q_t - LX_t$  cannot be satisfied for any  $t \geq 0$ , which implies that the solution of the optimal stopping zero-sum game in (2.3) is trivial in that case, for each  $0 < q < s$  fixed.

Summarising the arguments above, we realise that we are considering the three-dimensional continuous strong Markov process  $(X, S, Q)$  defined in (1.1)–(1.2) and (1.3) within the discounted optimal stopping zero-sum game of (2.3) with the continuous payoff functions  $F(x, s)$  and  $G(x, s, q)$  from (1.5) such that the equality  $F(x, s) = G(x, s, q)$  holds, when  $x \leq (s - q)/(K - L)$  as well as  $x \geq q/(L - K + 1)$ , for all  $(x, s, q) \in E$ , under the assumption  $0 < L < K < L + 1$ . Observe that the arguments of the proofs from [14, Theorem 2.1] and [55, Theorem 2.1] (also applicable for continuous strong Markov time-space processes with constant killing rates) can be naturally extended to the case of the discounted optimal stopping game of (2.3) for the process  $(X, S, Q)$  with the second and third components changing only at the diagonals  $d_1 = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < q \leq x = s\}$  and  $d_2 = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < q = x \leq s\}$ , respectively, of the state space  $E$  (see also [56, Theorem 3.1] for the corresponding result in a model based on a standard Brownian motion in the interval  $[0, 1]$  absorbed at either 0 or 1). Note that the natural analogues of the conditions of [14, Formula (2.1)] and [55, Formulae (2.9) and (2.12)] are clearly satisfied for the discounted payoffs  $e^{-rt}F(X_t, S_t)$  and  $e^{-rt}G(X_t, S_t, Q_t)$  from (1.5) (see e.g. [64, Formula (2.16)]). Hence, by applying the resulting extensions mentioned above, we may conclude that the continuation region  $C_*$  in (2.6) should belong to the set

$$E' = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < a'(s, q) \vee q \leq x \leq s \wedge b'(s, q)\} \tag{2.8}$$

with

$$a'(s, q) = (s - q)/(K - L) \quad \text{and} \quad b'(s, q) = q/(L - K + 1) \tag{2.9}$$

for  $0 < q < s$ , which represents all points  $(x, s, q)$  from the state space  $E$  of the process  $(X, S, Q)$  for which the solution of the original problem of (2.3) may be nontrivial, while the complement  $E \setminus E'$  surely belongs to the stopping region  $D_*$  in (2.7), under the assumption  $0 < L < K < L + 1$ . Therefore, the property  $\tau_* \wedge \zeta_* \leq \theta$  ( $\mathbb{P}_{x,s,q}$ -almost surely (a.s.)) holds, for  $\tau_*$  and  $\zeta_*$  from (2.4) and (2.5) and for any point  $(x, s, q) \in E'$ , under the assumption  $0 < L < K < L + 1$ , where we put

$$\theta = \inf \{t \geq 0 \mid \text{either } X_t \leq a'(S_t, Q_t) \text{ or } X_t \geq b'(S_t, Q_t)\}, \tag{2.10}$$

which is a stopping time of the process  $(X, S, Q)$ .

(ii) We now describe the structure of the continuation and stopping regions  $C_*$  and  $D_*$  from (2.6)–(2.7). For this purpose, by means of standard applications of Itô’s formula (see e.g. [48, Theorem 4.4] or [62, Chapter IV, Theorem 3.3]) to the processes  $e^{-rt}(S_t - KX_t)$  and  $e^{-rt}(Q_t - LX_t)$ , we obtain the representations

$$e^{-rt}(S_t - KX_t) = s - Kx + \int_0^t e^{-ru} (K \delta(S_u, Q_u) X_u - r S_u) du + \int_0^t e^{-ru} dS_u + N_t^1 \tag{2.11}$$

and

$$e^{-rt} (Q_t - L X_t) = q - Lx + \int_0^t e^{-ru} (L \delta(S_u, Q_u) X_u - r Q_u) du + \int_0^t e^{-ru} dQ_u + N_t^2 \tag{2.12}$$

for all  $t \geq 0$ . Here, the processes  $N^i = (N_t^i)_{t \geq 0}$ , for  $i = 1, 2$ , defined by

$$N_t^1 = -K \int_0^t e^{-ru} \sigma(S_u, Q_u) X_u dB_u \quad \text{and} \quad N_t^2 = -L \int_0^t e^{-ru} \sigma(S_u, Q_u) X_u dB_u \tag{2.13}$$

are continuous uniformly integrable martingales under the probability measure  $\mathbb{P}_{x,s,q}$ . Then, inserting  $\tau \wedge \zeta$  in place of  $t$  and applying Doob's optional sampling theorem (see e.g. [48, Chapter III, Theorem 3.6] and [62, Chapter II, Theorem 3.2]) to the expressions in (2.11) and (2.12), we get that the equalities

$$\begin{aligned} & \mathbb{E}_{x,s,q} [e^{-r\tau} (S_\tau - K X_\tau) I(\tau < \zeta) + e^{-r\zeta} (Q_\zeta - L X_\zeta) I(\zeta \leq \tau)] \\ &= \mathbb{E}_{x,s,q} [e^{-r(\tau \wedge \zeta)} (S_{\tau \wedge \zeta} - K X_{\tau \wedge \zeta}) - e^{-r\zeta} (S_\zeta - K X_\zeta - Q_\zeta + L X_\zeta) I(\zeta \leq \tau)] \\ &= s - Kx - \mathbb{E}_{x,s,q} [e^{-r\zeta} (S_\zeta - K X_\zeta - Q_\zeta + L X_\zeta) I(\zeta \leq \tau)] \\ & \quad + \mathbb{E}_{x,s,q} \left[ \int_0^{\tau \wedge \zeta} e^{-ru} (K \delta(S_u, Q_u) X_u - r S_u) du + \int_0^{\tau \wedge \zeta} e^{-ru} dS_u \right] \end{aligned} \tag{2.14}$$

and

$$\begin{aligned} & \mathbb{E}_{x,s,q} [e^{-r\tau} (S_\tau - K X_\tau) I(\tau < \zeta) + e^{-r\zeta} (Q_\zeta - L X_\zeta) I(\zeta \leq \tau)] \\ &= \mathbb{E}_{x,s,q} [e^{-r(\tau \wedge \zeta)} (Q_{\tau \wedge \zeta} - L X_{\tau \wedge \zeta}) + e^{-r\tau} (S_\tau - K X_\tau - Q_\tau + L X_\tau) I(\tau < \zeta)] \\ &= q - Lx + \mathbb{E}_{x,s,q} [e^{-r\tau} (S_\tau - K X_\tau - Q_\tau + L X_\tau) I(\tau < \zeta)] \\ & \quad + \mathbb{E}_{x,s,q} \left[ \int_0^{\tau \wedge \zeta} e^{-ru} (L \delta(S_u, Q_u) X_u - r Q_u) du + \int_0^{\tau \wedge \zeta} e^{-ru} dQ_u \right] \end{aligned} \tag{2.15}$$

hold, for any stopping times  $\tau$  and  $\zeta$  such that  $\tau \wedge \zeta \leq \theta$  ( $\mathbb{P}_{x,s,q}$ -a.s.) holds with  $\theta$  defined in (2.10), and for any starting point  $(x, s, q) \in E'$  of  $(X, S, Q)$ . Hence, it follows from the expressions in (2.14) and (2.15) and the structure of the optimal stopping times in (2.4) and (2.5) that the value function of the optimal stopping game in (2.3) admits the representations

$$\begin{aligned} V_*(x, s, q) &= s - Kx - \mathbb{E}_{x,s,q} [e^{-r\zeta_*} (S_{\zeta_*} - K X_{\zeta_*} - Q_{\zeta_*} + L X_{\zeta_*}) I(\zeta_* \leq \tau_*)] \\ & \quad + \mathbb{E}_{x,s,q} \left[ \int_0^{\tau_* \wedge \zeta_*} e^{-ru} (K \delta(S_u, Q_u) X_u - r S_u) du + \int_0^{\tau_* \wedge \zeta_*} e^{-ru} dS_u \right] \end{aligned} \tag{2.16}$$

and

$$\begin{aligned} V_*(x, s, q) &= q - Lx + \mathbb{E}_{x,s,q} [e^{-r\tau_*} (S_{\tau_*} - K X_{\tau_*} - Q_{\tau_*} + L X_{\tau_*}) I(\tau_* < \zeta_*)] \\ & \quad + \mathbb{E}_{x,s,q} \left[ \int_0^{\tau_* \wedge \zeta_*} e^{-ru} (L \delta(S_u, Q_u) X_u - r Q_u) du + \int_0^{\tau_* \wedge \zeta_*} e^{-ru} dQ_u \right] \end{aligned} \tag{2.17}$$

for the optimal stopping times  $\tau_*$  and  $\zeta_*$  forming a Nash equilibrium in (2.3), because the property  $\tau_* \wedge \zeta_* \leq \theta$  ( $\mathbb{P}_{x,s,q}$ -a.s.) holds with  $\theta$  as defined in (2.10), for any  $(x, s, q) \in E'$ .

Here and subsequently, we denote by  $\tau_* = \tau_*(x, s, q)$  and  $\zeta_* = \zeta_*(x, s, q)$  the optimal stopping times forming a Nash equilibrium in (2.3) for the starting point  $(x, s, q) \in E'$  of

the process  $(X, S, Q)$ , where  $E'$  is defined in (2.8) above. Thus, on the one hand, it follows from the structure of the integrand in the first integral of (2.16) and the fact that the second integral there increases whenever the process  $(X, S, Q)$  is located at the diagonal  $d_1 = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < q \leq x = s\}$  that it should not be optimal for the option holder (maximiser of the expected reward) to exercise the contract earlier than the option writer recalls it, when the inequalities  $a'(S_t, Q_t) \vee rS_t / (K\delta(S_t, Q_t)) < X_t \leq S_t \wedge b'(S_t, Q_t)$  hold, for any  $t \geq 0$ . Moreover, it follows from the structure of the integrand in first integral of (2.17) and the fact that the second integral there decreases whenever the process  $(X, S, Q)$  is located at the diagonal  $d_2 = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < q = x \leq s\}$  that it should not be optimal for the option writer (minimiser of the expected reward) to recall the contract earlier than the option holder exercises it, when the inequalities  $a'(S_t, Q_t) \vee Q_t \leq X_t < rQ_t / (L\delta(S_t, Q_t)) \wedge b'(S_t, Q_t)$  hold, for any  $t \geq 0$ . Since both participants of the contract are acting simultaneously, these facts yield that the set

$$C' = \{(x, s, q) \in E' \mid a'(s, q) \vee q \vee \bar{a}(s, q) < x < \underline{b}(s, q) \wedge s \wedge b'(s, q)\} \tag{2.18}$$

with

$$\bar{a}(s, q) = rs / (K\delta(s, q)) \quad \text{and} \quad \underline{b}(s, q) = rq / (L\delta(s, q)) \tag{2.19}$$

for  $0 < q < s$ , which may be nonempty because of the assumption that  $0 < L < K < L + 1$  holds, represents a part of the continuation region  $C_*$  in (2.6).

(iii) Let us finally prove the connectivity of the left-hand and right-hand parts of the stopping region  $D_*$  from (2.7) which are located outside the region  $C'$  from (2.18). For this purpose, we first recall from the arguments of Part (i) above that all the points  $(x', s, q)$  and  $(x'', s, q)$  of the set  $E \setminus E'$ , with  $E'$  as defined in (2.8), such that  $0 < x' < a'(s, q)$  and  $x'' > b'(s, q)$  belong to the stopping region  $D_*$ , for each  $0 < q < s$  fixed, under the assumption  $0 < L < K < L + 1$ . We also recall that the process  $X$  admits the explicit expression of (1.1) and provides a (pathwise) unique strong solution of the stochastic differential equation in (1.2) with the processes  $S$  and  $Q$  given by (1.3), so that the solutions starting from the different points  $x > 0$  do not intersect each other over the entire infinite time interval. For ease of presentation, in the rest of this section we indicate by  $(X^{(x)}, S^{(s,x)}, Q^{(q,x)})$  the dependence of the process  $(X, S, Q)$  defined in (1.1)–(1.2) and (1.3) on its starting point  $(x, s, q) \in E'$ .

Observe that, if we take some  $(x, s, q) \in D_*$  such that  $a'(s, q) < x < \bar{a}(s, q) \wedge s \wedge b'(s, q)$  holds, for each  $0 < q < s$  fixed, then the arguments of Part (ii) above would imply that it is not optimal for the option writer to recall the contract earlier than the holder exercises it, so that the value in (2.3) would admit the representation

$$\begin{aligned} &V_*(x', s, q) - (s - Kx') \tag{2.20} \\ &= \mathbb{E} \left[ \int_0^{\tau'_*} e^{-ru} (K\delta(S_u^{(s,x')}, Q_u^{(q,x')}) X_u^{(x')} - rS_u^{(s,x')}) du + \int_0^{\tau'_*} e^{-ru} dS_u^{(s,x')} \right], \end{aligned}$$

where  $\tau'_* = \tau_*(x', s, q)$  denotes the optimal exercise time for the holder in the problem of (2.3) under the assumption that the process  $(X, S, Q)$  starts at  $(x', s, q)$ , for all  $a'(s, q) \leq x' < x$  and each  $0 < q < s$  fixed. Hence, because of the assumption that the function  $\delta(s, q)$  is increasing in both the variables  $s$  and  $q$  on  $[0, \infty]^2$  and the fact that the process  $(X, S, Q)$  started at  $(x', s, q)$  reaches the point  $(x, s, q')$  for some  $0 < q' \leq q$  before hitting the upper diagonal  $d_1 = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < q \leq x = s\}$ , we see that the integrand in the first integral in the right-hand side of (2.20) (and thus the resulting total expected reward functional there) is



increasing in  $x'$  for  $a'(s, q) \leq x' < x$ , for each  $0 < q < s$  fixed. Thus, we may conclude that the inequalities

$$V_*(x', s, q) - (s - Kx') \leq V_*(x, s, q) - (s - Kx) = 0 \tag{2.21}$$

hold, so that the point  $(x', s, q)$  such that  $a'(s, q) \leq x' < x < \bar{a}(s, q) \wedge s \wedge b'(s, q)$  also belongs to the left-hand part of the stopping region  $D_*$  in (2.7).

Similarly, if we take some  $(x, s, q) \in D_*$  such that  $a'(s, q) \vee q \vee \underline{b}(s, q) < x < b'(s, q)$  holds, for each  $0 < q < s$  fixed, then the arguments of Part (ii) above would imply that it is not optimal for the option holder to exercise the contract earlier than the writer recalls it, so that the value in (2.3) would admit the representation

$$\begin{aligned} & V_*(x'', s, q) - (q - Lx'') \tag{2.22} \\ &= \mathbb{E} \left[ \int_0^{\zeta'_*} e^{-ru} (L \delta(S_u^{(s,x'')}, Q_u^{(q,x'')}) X_u^{(x'')} - r Q_u^{(q,x'')}) du + \int_0^{\zeta'_*} e^{-ru} dQ_u^{(q,x'')} \right], \end{aligned}$$

where  $\zeta'_* = \zeta_*(x'', s, q)$  denotes the optimal exercise time for the writer in the problem of (2.3) under the assumption that the process  $(X, S, Q)$  starts at  $(x'', s, q)$ , for all  $x < x'' \leq b'(s, q)$  and each  $0 < q < s$  fixed. Hence, by the assumption that the function  $\delta(s, q)$  is increasing in both the variables  $s$  and  $q$  on  $[0, \infty]^2$  and the fact that the process  $(X, S, Q)$  started at  $(x'', s, q)$  reaches the point  $(x, s', q)$  for some  $0 < s \leq s'$  before hitting the lower diagonal  $d_2 = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < q = x \leq s\}$ , we see that the integrand in the first integral in the right-hand side of (2.22) (and thus the resulting total expected reward functional there) is increasing in  $x''$  for  $x < x'' \leq b'(s, q)$ , for each  $0 < q < s$  fixed. Thus, we may conclude that the inequalities

$$V_*(x'', s, q) - (q - Lx'') \geq V_*(x, s, q) - (q - Lx) = 0 \tag{2.23}$$

hold, so that the point  $(x'', s, q)$  such that  $a'(s, q) \vee q \vee \underline{b}(s, q) < x < x'' \leq b'(s, q)$  also belongs to the right-hand part of the stopping region  $D_*$  in (2.7).

Combining these arguments with the facts deduced in Parts (i)–(ii) above and noting the comments in [13, Subsection 3.3] and [52, Subsection 3.3], we conclude that there exist functions  $a_*(s, q)$  and  $b_*(s, q)$  satisfying the inequalities  $a'(s, q) \leq a_*(s, q) \leq \bar{a}(s, q) \wedge s \wedge b'(s, q)$  and  $a'(s, q) \vee q \vee \underline{b}(s, q) \leq b_*(s, q) \leq b'(s, q)$ , for  $0 < q < s$ , such that the continuation region  $C_*$  in (2.6) has the form

$$C_* = \{(x, s, q) \in E' \mid a_*(s, q) < x < b_*(s, q)\}, \tag{2.24}$$

while the stopping region  $D_*$  in (2.7) is given by

$$D_* = \{(x, s, q) \in E \mid \text{either } x \leq a_*(s, q) \text{ or } x \geq b_*(s, q)\}. \tag{2.25}$$

(iv) Finally, in order to determine upper and lower bounds for the value function in (2.3) and optimal stopping boundaries in (2.24) and (2.25), we consider the optimal stopping problems with the value functions  $\bar{V}(x, s, q)$  and  $\underline{V}(x, s, q)$  from (5.1). It is shown in Section 5 below that the functions  $\bar{V}(x, s, q)$  and  $\underline{V}(x, s, q)$  admit the explicit expressions in (5.11) and (5.12), while the associated optimal stopping times  $\bar{\tau}$  and  $\underline{\zeta}$  are given by (5.2), where the boundaries  $\underline{a}(s, q)$  and  $\bar{b}(s, q)$  are determined as the maximal and minimal solutions of the first-order nonlinear ordinary differential equations in (3.24) and (3.27) staying below or above the diagonals  $d_1 = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < q \leq x = s\}$  and  $d_2 = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < q = x \leq s\}$ , respectively. If we suppose that either the inequality  $a_*(s, q) < \underline{a}(s, q)$  or the inequality  $b_*(s, q) > \bar{b}(s, q)$  holds, then,




FIGURE 1. A computer drawing of the optimal exercise boundaries  $a_*(s, q)$ ,  $b_*(s, q)$ , and  $\underline{a}(s, q)$ , for each  $q > 0$  fixed.




FIGURE 2. A computer drawing of the optimal exercise boundaries  $a_*(s, q)$ ,  $b_*(s, q)$ , and  $\bar{b}(s, q)$ , for each  $s > 0$  fixed.

for each given and fixed  $x > 0$  such that either  $x \in (a_*(s, q), \underline{a}(s, q))$  or  $x \in (\bar{b}(s, q), b_*(s, q))$ , we would have either  $V_*(x, s, q) > s - Kx = \bar{V}(x, s, q)$  or  $V_*(x, s, q) < q - Lx = \underline{V}(x, s, q)$ , respectively, contradicting the obvious fact that the inequalities  $V_*(x, s, q) \leq \bar{V}(x, s, q)$  and  $V_*(x, s, q) \geq \underline{V}(x, s, q)$  hold for all  $(x, s, q) \in E'$ . Thus, we may conclude that the inequalities  $a_*(s, q) \geq \underline{a}(s, q)$  and  $b_*(s, q) \leq \bar{b}(s, q)$  should be satisfied for all  $0 < q < s$  (see Figures 1 and 2 for computer drawings of the optimal stopping boundaries  $a_*(s, q)$  and  $b_*(s, q)$ , as well as the estimates  $\underline{a}(s, q)$  and  $\bar{b}(s, q)$ ).

### 2.3. The three-dimensional coupled free-boundary problem

By means of standard arguments based on an application of Itô’s formula, it can be shown that the infinitesimal operator  $\mathbb{L}$  of the process  $(X, S, Q)$  has the form

$$\mathbb{L} = (r - \delta(s, q))x \partial_x + \frac{\sigma^2(s, q)x^2}{2} \partial_{xx} \quad \text{in } 0 < q < x < s, \tag{2.26}$$

$$\partial_s = 0 \quad \text{at } 0 < q \leq x = s, \quad \text{and} \quad \partial_q = 0 \quad \text{at } 0 < q = x \leq s \tag{2.27}$$

(see e.g. [52, Subsection 3.1]). In order to find analytic expressions for the unknown value function  $V_*(x, s, q)$  from (2.3) and the unknown boundaries  $a_*(s, q)$  and  $b_*(s, q)$  from (2.24)–(2.25), let us build on the results of the general theory of optimal stopping problems for Markov processes (see e.g. [59, Chapter IV, Section 8]). We can reduce the optimal stopping game of (2.3) to the equivalent coupled free-boundary problem for  $V(x, s, q)$  with  $a(s, q)$  and  $b(s, q)$  given by

$$(\mathbb{L}V - rV)(x, s, q) = 0 \quad \text{for } q \vee a(s, q) < x < b(s, q) \wedge s, \tag{2.28}$$

$$V(x, s, q)|_{x=a(s,q)+} = s - K a(s, q), \quad V(x, s, q)|_{x=b(s,q)-} = q - L b(s, q), \tag{2.29}$$

$$V(x, s, q) = s - Kx \quad \text{for } x < a(s, q), \quad V(x, s, q) = q - Lx \quad \text{for } x > b(s, q), \tag{2.30}$$

$$s - Kx < V(x, s, q) < q - Lx \quad \text{for } a(s, q) < x < b(s, q), \tag{2.31}$$

$$(\mathbb{L}V - rV)(x, s, q) < 0 \quad \text{for } x < a(s, q), \quad (\mathbb{L}V - rV)(x, s, q) > 0 \quad \text{for } x > b(s, q), \tag{2.32}$$

where the *instantaneous-stopping* conditions in (2.29) are respectively satisfied when either  $a(s, q) \geq a'(s, q) \vee q$  or  $b(s, q) \leq s \wedge b'(s, q)$  holds, for each  $0 < q < s$ . Moreover, we further assume that the *smooth-fit* conditions

$$\partial_x V(x, s, q)|_{x=a(s,q)+} = -K, \quad \partial_x V(x, s, q)|_{x=b(s,q)-} = -L \tag{2.33}$$

are satisfied, when either  $a(s, q) > a'(s, q) \vee q$  or  $b(s, q) < s \wedge b'(s, q)$  holds, while the *normal-reflection* conditions

$$\partial_s V(x, s, q)|_{x=s-} = 0, \quad \partial_q V(x, s, q)|_{x=q+} = 0 \tag{2.34}$$

are satisfied, when either  $b'(s, q) \geq b(s, q) > s$  or  $a'(s, q) \leq a(s, q) < q$  holds, for each  $0 < q < s$ . On the one hand, when either the inequality  $a(s, q) > a'(s, q) \vee q$  or the inequality  $b(s, q) < s \wedge b'(s, q)$  holds, for some  $0 < q < s$ , the continuous process  $X$  can cross the left-hand boundary  $a(S, Q)$  before hitting the lower diagonal  $d_2$  or cross the right-hand boundary  $b(S, Q)$  before hitting the upper diagonal  $d_1$ , so that we can assume that the left-hand or the right-hand smooth-fit conditions of (2.33) are satisfied for the candidate value function  $V(x, s, q)$  at  $a(s, q)$  or  $b(s, q)$ , respectively. On the other hand, when either the inequalities  $b'(s, q) \geq b(s, q) > s$  or the inequalities  $a'(s, q) \leq a(s, q) < q$  hold, for some  $0 < q < s$ , the process  $X$  can hit the upper diagonal  $d_1 = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < q \leq x = s\}$  before crossing the right-hand boundary  $b(S, Q)$  or hit the lower diagonal  $d_2 = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < q = x \leq s\}$  before crossing the left-hand boundary  $a(S, Q)$ , so that we can assume that either the left-hand or the right-hand normal-reflection conditions of (2.34) are satisfied for  $V(x, s, q)$  at  $d_1$  or  $d_2$ , respectively. These properties are verified in the proof of Theorem 1 below, while the inequalities in (2.32) follow directly from the arguments of Parts (i)–(ii) of Subsection 2.2 above.

### 3. Solutions to the coupled free-boundary problem

In this section, we obtain closed-form expressions for the value function  $V_*(x, s, q)$  in (2.3) associated with the perpetual floating-strike lookback game option and derive first-order non-linear ordinary differential equations for the optimal stopping boundaries  $a_*(s, q)$  and  $b_*(s, q)$  from (2.24)–(2.25) forming a solution to the free-boundary problem in (2.28)–(2.32) with (2.33) and (2.34).

#### 3.1. The candidate value function

We first observe that the general solution of the second-order ordinary differential equation in (2.28) has the form

$$V(x, s, q) = C_1(s, q) x^{\gamma_1(s, q)} + C_2(s, q) x^{\gamma_2(s, q)}, \tag{3.1}$$

where  $C_i(s, q)$ , for  $i = 1, 2$ , are some continuously differentiable functions and  $\gamma_i(s, q)$ , for  $i = 1, 2$ , are given by

$$\gamma_i(s, q) = \frac{1}{2} - \frac{r - \delta(s, q)}{\sigma^2(s, q)} - (-1)^i \sqrt{\left(\frac{1}{2} - \frac{r - \delta(s, q)}{\sigma^2(s, q)}\right)^2 + \frac{2r}{\sigma^2(s, q)}}, \tag{3.2}$$

so that  $\gamma_2(s, q) < 0 < 1 < \gamma_1(s, q)$  holds for all  $0 < q < s$ . Then, by applying the instantaneous-stopping conditions from (2.29) to the function in (3.1), we get that the equalities

$$C_1(s, q) a^{\gamma_1(s, q)}(s, q) + C_2(s, q) a^{\gamma_2(s, q)}(s, q) = s - K a(s, q), \tag{3.3}$$

$$C_1(s, q) b^{\gamma_1(s, q)}(s, q) + C_2(s, q) b^{\gamma_2(s, q)}(s, q) = q - L b(s, q) \tag{3.4}$$

are satisfied when  $a(s, q) \geq a'(s, q) \vee q$  and  $b(s, q) \leq s \wedge b'(s, q)$  respectively hold, for each  $0 < q < s$ . Hence, by using the smooth-fit conditions from (2.33), we obtain that the equalities

$$C_1(s, q) \gamma_1(s, q) a^{\gamma_1(s, q)}(s, q) + C_2(s, q) \gamma_2(s, q) a^{\gamma_2(s, q)}(s, q) = -K a(s, q), \tag{3.5}$$

$$C_1(s, q) \gamma_1(s, q) b^{\gamma_1(s, q)}(s, q) + C_2(s, q) \gamma_2(s, q) b^{\gamma_2(s, q)}(s, q) = -L b(s, q) \tag{3.6}$$

are satisfied when  $a(s, q) > a'(s, q) \vee q$  and  $b(s, q) < s \wedge b'(s, q)$  respectively hold, for each  $0 < q < s$ . Thus, by applying the normal-reflection conditions from (2.34) to the function in (3.1), we obtain that the equalities

$$\sum_{i=1}^2 \left( \partial_s C_i(s, q) s^{\gamma_i(s, q)} + C_i(s, q) \partial_s \gamma_i(s, q) s^{\gamma_i(s, q)} \ln s \right) = 0, \tag{3.7}$$

$$\sum_{i=1}^2 \left( \partial_q C_i(s, q) q^{\gamma_i(s, q)} + C_i(s, q) \partial_q \gamma_i(s, q) q^{\gamma_i(s, q)} \ln q \right) = 0 \tag{3.8}$$

are satisfied when  $b'(s, q) \geq b(s, q) > s$  and  $a'(s, q) \leq a(s, q) < q$  respectively hold, for each  $0 < q < s$ . Here, the partial derivatives  $\partial_s \gamma_i(s, q)$  and  $\partial_q \gamma_i(s, q)$  take the form

$$\partial_s \gamma_i(s, q) = \varphi(s, q) - (-1)^i \frac{\varphi(s, q)(\gamma_1(s, q) + \gamma_2(s, q))\sigma^3(s, q) - 4r\partial_s \sigma(s, q)}{\sigma^2(s, q)\sqrt{(\gamma_1(s, q) + \gamma_2(s, q))^2\sigma^2(s, q) + 8r}}, \tag{3.9}$$

$$\partial_q \gamma_i(s, q) = \psi(s, q) - (-1)^i \frac{\psi(s, q)(\gamma_1(s, q) + \gamma_2(s, q))\sigma^3(s, q) - 4r\partial_q \sigma(s, q)}{\sigma^2(s, q)\sqrt{(\gamma_1(s, q) + \gamma_2(s, q))^2\sigma^2(s, q) + 8r}} \tag{3.10}$$

for  $i = 1, 2$ , and the functions  $\varphi(s, q)$  and  $\psi(s, q)$  are defined by

$$\varphi(s, q) = \frac{\sigma(s, q)\partial_s \delta(s, q) + 2(r - \delta(s, q))\partial_s \sigma(s, q)}{\sigma^3(s, q)}, \tag{3.11}$$

$$\psi(s, q) = \frac{\sigma(s, q)\partial_q \delta(s, q) + 2(r - \delta(s, q))\partial_q \sigma(s, q)}{\sigma^3(s, q)} \tag{3.12}$$

for  $0 < q < s$ .

Now, by solving the system of equations in (3.3)–(3.4), we obtain that the function in (3.1) admits the representation

$$V(x, s, q; a(s, q), b(s, q)) = \sum_{i=1}^2 C_i(s, q; a(s, q), b(s, q)) x^{\gamma_i(s, q)} \tag{3.13}$$

for  $a'(s, q) \vee q \leq a(s, q) < x < b(s, q) \leq s \wedge b'(s, q)$ , where

$$C_i(s, q; a(s, q), b(s, q)) = \frac{(s - Ka(s, q))b^{\gamma_{3-i}(s, q)}(s, q) - (q - Lb(s, q))a^{\gamma_{3-i}(s, q)}(s, q)}{a^{\gamma_i(s, q)}(s, q)b^{\gamma_{3-i}(s, q)}(s, q) - b^{\gamma_i(s, q)}(s, q)a^{\gamma_{3-i}(s, q)}(s, q)} \tag{3.14}$$

when  $a'(s, q) \vee q \leq a(s, q) < b(s, q) \leq s \wedge b'(s, q)$  holds, for every  $i = 1, 2$ . Then, by solving the system of equations in (3.3) and (3.5), we obtain that the function in (3.1) admits the representation

$$V(x, s, q; a(s, q)) = C_1(s, q; a(s, q)) x^{\gamma_1(s, q)} + C_2(s, q; a(s, q)) x^{\gamma_2(s, q)} \tag{3.15}$$

for  $a'(s, q) \vee q \leq a(s, q) < x \leq s < b(s, q) \leq b'(s, q)$ , where

$$C_i(s, q; a(s, q)) = \frac{\gamma_{3-i}(s, q)(s - Ka(s, q)) + Ka(s, q)}{(\gamma_{3-i}(s, q) - \gamma_i(s, q))a^{\gamma_i(s, q)}(s, q)} \tag{3.16}$$

when  $a'(s, q) \vee q \leq a(s, q) < s < b(s, q) \leq b'(s, q)$  holds, for every  $i = 1, 2$ . Also, by solving the system of equations in (3.4) and (3.6), we obtain that the function in (3.1) admits the representation

$$V(x, s, q; b(s, q)) = C_1(s, q; b(s, q)) x^{\gamma_1(s, q)} + C_2(s, q; b(s, q)) x^{\gamma_2(s, q)} \tag{3.17}$$

for  $a'(s, q) \leq a(s, q) < q < b(s, q) \leq s \wedge b'(s, q)$ , where

$$C_i(s, q; b(s, q)) = \frac{\gamma_{3-i}(s, q)(q - Lb(s, q)) + Lb(s, q)}{(\gamma_{3-i}(s, q) - \gamma_i(s, q))b^{\gamma_i(s, q)}(s, q)} \tag{3.18}$$

when  $a'(s, q) \leq a(s, q) < q < b(s, q) \leq s \wedge b'(s, q)$  holds, for every  $i = 1, 2$ .

Finally, by means of straightforward computations, it can be deduced from the expression in (3.13) that the first-order and second-order partial derivatives  $\partial_x V(x, s, q; a(s, q), b(s, q))$  and  $\partial_{xx} V(x, s, q; a(s, q), b(s, q))$  of the function  $V(x, s, q; a(s, q), b(s, q))$  take the forms

$$\partial_x V(x, s, q; a(s, q), b(s, q)) = \sum_{i=1}^2 C_i(s, q; a(s, q), b(s, q)) \gamma_i(s, q) x^{\gamma_i(s, q)-1} \tag{3.19}$$

and

$$\partial_{xx}V(x, s, q; a(s, q), b(s, q)) = \sum_{i=1}^2 C_i(s, q; a(s, q), b(s, q)) \gamma_i(s, q)(\gamma_i(s, q) - 1) x^{\gamma_i(s, q)-2} \tag{3.20}$$

for  $a'(s, q) \vee q \leq a(s, q) < x < b(s, q) \leq s \wedge b'(s, q)$ . Note that the same first-order and second-order partial derivatives of the functions  $V(x, s, q; a(s, q))$ , for  $a'(s, q) \vee q \leq a(s, q) < x < b(s, q) \leq b'(s, q)$ , and  $V(x, s, q; b(s, q))$ , for  $a'(s, q) \leq a(s, q) < q < b(s, q) \leq s \wedge b'(s, q)$ , from (3.15) with (3.16) and (3.17) with (3.18) are computed similarly.

### 3.2. The candidate stopping boundaries

We now apply the conditions of (3.5)–(3.6) to the functions  $C_i(s, q; a(s, q), b(s, q))$ , for  $i = 1, 2$ , in (3.14) to obtain the equalities

$$\frac{\gamma_i(s, q)(s - Ka(s, q)) + Ka(s, q)}{\gamma_i(s, q)(q - Lb(s, q)) + Lb(s, q)} = \left(\frac{a(s, q)}{b(s, q)}\right)^{\gamma_3-i(s, q)} \tag{3.21}$$

for  $a'(s, q) \vee q \leq a(s, q) < b(s, q) \leq s \wedge b'(s, q)$  and  $i = 1, 2$ . Then we set  $b(s, q) = b'(s, q)$  and apply the condition of (3.5) to the same functions to obtain the equality

$$\sum_{i=1}^2 \frac{(s - Ka(s, q))b'^{\gamma_3-i(s, q)}(s, q) - (q - Lb'(s, q))a'^{\gamma_3-i(s, q)}(s, q)}{a'^{\gamma_i(s, q)}(s, q)b'^{\gamma_3-i(s, q)}(s, q) - b'^{\gamma_i(s, q)}(s, q)a'^{\gamma_3-i(s, q)}(s, q)} \gamma_i(s, q) a'^{\gamma_i(s, q)}(s, q) \tag{3.22}$$

$$= -K a(s, q)$$

for  $a'(s, q) \vee q \leq a(s, q) < s \wedge b'(s, q) < b(s, q)$ . Also, we set  $a(s, q) = a'(s, q)$  and apply the condition of (3.6) to the same functions to obtain the equality

$$\sum_{i=1}^2 \frac{(s - Ka'(s, q))b'^{\gamma_3-i(s, q)}(s, q) - (q - Lb(s, q))a'^{\gamma_3-i(s, q)}(s, q)}{a'^{\gamma_i(s, q)}(s, q)b'^{\gamma_3-i(s, q)}(s, q) - b'^{\gamma_i(s, q)}(s, q)a'^{\gamma_3-i(s, q)}(s, q)} \gamma_i(s, q) b'^{\gamma_i(s, q)}(s, q) \tag{3.23}$$

$$= -L b(s, q)$$

for  $a(s, q) < a'(s, q) \vee q < b(s, q) \leq s \wedge b'(s, q)$ .

The existence and uniqueness of solutions of the system of arithmetic equations in (3.21) as well as of the equations in (3.22) and (3.23) on the admissible intervals follow from the arguments of Subsection 3.4 below. Observe that the system of arithmetic equations in (3.21) and the equation in (3.23) satisfy the conditions of the classical (two-dimensional) implicit function theorem, so that the resulting solutions  $a_*(s, q)$  and  $b_*(s, q)$  turn out to be continuously differentiable. Furthermore, assuming that the candidate boundary functions  $a(s, q)$  and  $b(s, q)$  are continuously differentiable, we apply the condition of (3.7) to the functions  $C_i(s, q; a(s, q))$ , for  $i = 1, 2$ , in (3.16) to conclude that the candidate boundary  $a(s, q)$  satisfies the ordinary differential equation

$$\partial_s a(s, q) = \sum_{j=1}^2 \frac{C_j(s, q; a(s, q)) \partial_s \gamma_j(s, q) s^{\gamma_j(s, q)} \ln s + \Psi_{1,j}(a(s, q), s, q) (s/a(s, q))^{\gamma_j(s, q)}}{\Phi_1(a(s, q), s, q) ((s/a(s, q))^{\gamma_1(s, q)} - (s/a(s, q))^{\gamma_2(s, q)})} \tag{3.24}$$

for  $a'(s, q) \vee q \leq a(s, q) < s < b(s, q) \leq b'(s, q)$ , where we set

$$\Phi_1(x, s, q) = \frac{(\gamma_1(s, q) + \gamma_2(s, q))K + \gamma_1(s, q)\gamma_2(s, q)(s - Kx)/x}{\gamma_2(s, q) - \gamma_1(s, q)} \tag{3.25}$$

and

$$\Psi_{1,i}(x, s, q) = \frac{\partial_s \gamma_{3-i}(s, q)(s - Kx) + \gamma_{3-i}(s, q)}{\gamma_{3-i}(s, q) - \gamma_i(s, q)} - \frac{(Kx + \gamma_{3-i}(s, q)(s - Kx))(\partial_s \gamma_i(s, q) \ln x + \partial_s \ln(\gamma_{3-i}(s, q) - \gamma_i(s, q)))}{\gamma_{3-i}(s, q) - \gamma_i(s, q)} \tag{3.26}$$

for all  $0 < q \leq x \leq s$  and every  $i = 1, 2$ . We also apply the condition of (3.8) to the functions  $C_i(s, q; b(s, q))$ , for  $i = 1, 2$ , in (3.18) to conclude that the candidate boundary  $b(s, q)$  satisfies the ordinary differential equation

$$\partial_q b(s, q) = \sum_{i=1}^2 \frac{C_i(s, q; b(s, q)) \partial_q \gamma_i(s, q) q^{\gamma_i(s, q)} \ln q + \Psi_{2,i}(b(s, q), s, q)(q/b(s, q))^{\gamma_i(s, q)}}{\Phi_2(b(s, q), s, q)((q/b(s, q))^{\gamma_1(s, q)} - (q/b(s, q))^{\gamma_2(s, q)})} \tag{3.27}$$

for  $a'(s, q) \leq a(s, q) < q < b(s, q) \leq s \wedge b'(s, q)$ , where we set

$$\Phi_2(x, s, q) = \frac{(\gamma_1(s, q) + \gamma_2(s, q))L + \gamma_1(s, q)\gamma_2(s, q)(q - Lx)/x}{\gamma_2(s, q) - \gamma_1(s, q)} \tag{3.28}$$

and

$$\Psi_{2,i}(x, s, q) = \frac{\partial_q \gamma_{3-i}(s, q)(q - Lx) + \gamma_{3-i}(s, q)}{\gamma_{3-i}(s, q) - \gamma_i(s, q)} - \frac{(Lx + \gamma_{3-i}(s, q)(q - Lx))(\partial_q \gamma_i(s, q) \ln x + \partial_q \ln(\gamma_{3-i}(s, q) - \gamma_i(s, q)))}{\gamma_{3-i}(s, q) - \gamma_i(s, q)} \tag{3.29}$$

for all  $0 < q \leq x \leq s$  and every  $i = 1, 2$ . Note that, by virtue of the assumptions on the coefficients  $\delta(s, q) > 0$  and  $\sigma(s, q) > 0$  of the diffusion-type process  $X$  from (1.1)–(1.2) and (1.3), the right-hand sides of the expressions in (3.24) with (3.25)–(3.26) and in (3.27) with (3.28)–(3.29) are (locally) continuous in  $(s, q, a(s, q))$  and  $(s, q, b(s, q))$  and (locally) Lipschitz in  $a(s, q)$  and  $b(s, q)$ , for each  $0 < q < s$  fixed. Thus, by the classical results on the existence and uniqueness of solutions for first-order nonlinear ordinary differential equations, the equations in (3.24) and (3.27) admit (locally) unique solutions, which can be constructed by means of Picard’s method of successive approximations (see Subsection 5.3 below for further constructions and references).

### 3.3. The structure of the continuation region

In order to specify the optimal exercise boundaries for the floating-strike lookback game options, let us consider the functions  $a_*(s, q)$  and  $b_*(s, q)$ , which provide a unique solution to the system of arithmetic equations in (3.21) such that  $a'(s, q) \leq a_*(s, q) < b_*(s, q) \leq b'(s, q)$ , or the function  $a_*(s, q) < b_*(s, q) = b'(s, q)$  represents the largest root of the equation in (3.22), or the function  $b_*(s, q) > a_*(s, q) = a'(s, q)$  represents the largest root of the equation in (3.23), or otherwise the properties  $a_*(s, q) = a'(s, q)$  and  $b_*(s, q) = b'(s, q)$  hold, for each  $0 < q < s$  fixed.

On the one hand, we can set  $s_0^*(q) = \infty$  and define the functions  $s_{2k-1}^*(q) = \sup\{s < s_{2k-2}^*(q) \mid b_*(s, q) > s\}$  and  $s_{2k}^*(q) = \sup\{s < s_{2k-1}^*(q) \mid b_*(s, q) < s\}$ , whenever they exist, and put  $s_{2k-1}^*(q) = s_{2k}^*(q) = 0$  otherwise, so that the inequalities  $0 \leq s_{2k-1}^*(q) \leq s_{2k-2}^*(q) \leq \infty$  hold, for all  $k \in \mathbb{N}$ , and each  $q > 0$  fixed. In other words, the boundary  $b_*(s, q)$  exits the region  $E$  from the side of the diagonal  $d_1 = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < q \leq x = s\}$  passing through the points  $(s_{2k-1}^*(q), s_{2k-1}^*(q), q)$  and comes back to  $E$  from the side of  $d_1$  passing through the points  $(s_{2k}^*(q), s_{2k}^*(q), q)$ , for  $k \in \mathbb{N}$ , for each  $q > 0$  fixed. Hence, the candidate value function  $V(x, s, q; a_*(s, q), b_*(s, q))$  admits the representation of (3.13) with (3.14) and the candidate stopping boundaries  $a_*(s, q)$  and  $b_*(s, q)$  solve the system of arithmetic equations in (3.21) in the regions

$$\tilde{R}_{2k-1}(a_*, b_*) = \{(x, s, q) \in E' \mid s_{2k-1}^*(q) \leq s < s_{2k-2}^*(q)\}, \tag{3.30}$$

while the candidate value function  $V(x, s, q; a_*(s, q))$  admits the representation of (3.15) with (3.16) and the candidate stopping boundary  $a_*(s, q)$  either solves the first-order nonlinear ordinary differential equation in the regions

$$\tilde{R}_{2k}(a_*) = \{(x, s, q) \in E' \mid s_{2k}^*(q) \leq s < s_{2k-1}^*(q)\} \tag{3.31}$$

both representing subsets of the continuation region  $C_*$  in (2.24), or coincides with  $a'$  ( $s, q$ ), for each  $q > 0$  fixed, and every  $k \in \mathbb{N}$ . Furthermore, we observe that the process  $(X, S, Q)$  can enter the region  $\tilde{R}_{2k-1}(a_*, b_*)$  in (3.31) from the region  $\tilde{R}_{2k}(a_*)$  in (3.30) only through the point  $(s_{2k-1}^*(q), s_{2k-1}^*(q), q)$ , for any  $k \in \mathbb{N}$ , by hitting the plane  $d_1 = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < q \leq x = s\}$ , that is, by increasing its second component  $S$ . Therefore, the candidate value function should satisfy the instantaneous-stopping and smooth-fit conditions at the points  $(s_{2k-1}^*(q), s_{2k-1}^*(q), q)$ , which are expressed by the equalities

$$\sum_{i=1}^2 C_i(s_{2k-1}^*(q)-, q; a_*(s_{2k-1}^*(q)-, q)) (s_{2k-1}^*(q))^{Y_i(s_{2k-1}^*(q), q)} \tag{3.32}$$

$$= V(s_{2k-1}^*(q), s_{2k-1}^*(q), q; a_*(s_{2k-1}^*(q), q), b_*(s_{2k-1}^*(q), q)),$$

$$\sum_{i=1}^2 C_i(s_{2k-1}^*(q)-, q; a_*(s_{2k-1}^*(q)-, q)) Y_i(s_{2k-1}^*(q)-, q) (s_{2k-1}^*(q))^{Y_i(s_{2k-1}^*(q), q)} \tag{3.33}$$

$$= s_{2k-1}^*(q) \partial_x V(s_{2k-1}^*(q), s_{2k-1}^*(q), q; a_*(s_{2k-1}^*(q), q), b_*(s_{2k-1}^*(q), q))$$

where the functions  $C_i(s, q; a_*(s, q))$ , for  $i = 1, 2$ , are given by (3.16) and the function  $V(x, s, q; a_*(s, q), b_*(s, q))$  has the form of (3.13) with (3.14), while the boundary  $a_*(s, q)$  provides a unique solution of the first-order nonlinear ordinary differential equation in (3.24) in the region  $\tilde{R}_{2k}(a_*)$  from (3.31) satisfying the (starting) conditions of (3.32)–(3.33) above, for each  $q > 0$  fixed, and every  $k \in \mathbb{N}$ , respectively.

On the other hand, we can set  $q_0^*(s) = 0$ , define the functions  $q_{2l-1}^*(s) = \inf\{q > q_{2l-2}^*(s) \mid a_*(s, q) < q\}$  and  $q_{2l}^*(s) = \inf\{q > q_{2l-1}^*(s) \mid a_*(s, q) > q\}$ , whenever they exist, and put  $q_{2l-1}^*(s) = q_{2l}^*(s) = \infty$  otherwise, so that the inequalities  $0 \leq q_{2l-2}^*(s) \leq q_{2l-1}^*(s) \leq \infty$  hold for all  $l \in \mathbb{N}$  and each  $s > 0$  fixed. In other words, the boundary  $a_*(s, q)$  exits the region  $E$  from the side of the diagonal  $d_2 = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < q = x \leq s\}$  passing through the points  $(q_{2l-1}^*(s), s, q_{2l-1}^*(q))$  and comes back to  $E$  from the side of the diagonal  $d_2$  passing through the points  $(q_{2l-1}^*(s), s, q_{2l-1}^*(s))$ , for  $k \in \mathbb{N}$ , for each  $s > 0$  fixed. Hence, the candidate value



function  $V(x, s, q; a_*(s, q), b_*(s, q))$  admits the representation of (3.13) with (3.14), and the candidate stopping boundaries  $a_*(s, q)$  and  $b_*(s, q)$  solve the system of arithmetic equations in (3.21) in the regions

$$\widehat{R}_{2l-1}(a_*, b_*) = \{(x, s, q) \in E' \mid q_{2l-2}^*(s) < q \leq q_{2l-1}^*(s)\}, \tag{3.34}$$

while the candidate value function  $V(x, s, q; b_*(s, q))$  admits the representation of (3.17) with (3.18), and the candidate stopping boundary  $b_*(s, q)$  solves the first-order nonlinear ordinary differential equation in the regions

$$\widehat{R}_{2l}(b_*) = \{(x, s, q) \in E' \mid q_{2l-1}^*(s) < q \leq q_{2l}^*(s)\}, \tag{3.35}$$

both representing subsets of the continuation region  $C_*$  in (2.24), for each  $s > 0$  fixed and every  $l \in \mathbb{N}$ . Furthermore, we observe that the process  $(X, S, Q)$  can enter the region  $\widehat{R}_{2l-1}(a_*, b_*)$  in (3.35) from the region  $\widehat{R}_{2l}(b_*)$  in (3.34) only through the point  $(q_{2l-1}^*(s), s, q_{2l-1}^*(s))$ , for any  $l \in \mathbb{N}$ , by hitting the plane  $d_2 = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < q = x \leq s\}$ , that is, by decreasing its third component  $Q$ . Therefore, the candidate value function should satisfy the instantaneous-stopping and smooth-fit conditions at the points  $(q_{2l-1}^*(s), s, q_{2l-1}^*(s))$ , which are expressed by the equalities

$$\begin{aligned} & \sum_{i=1}^2 C_i(s, q_{2l-1}^*(s) + ; b_*(s, q_{2l-1}^*(s) + )) (q_{2l-1}^*(s))^{\gamma_i(s, q_{2l-1}^*(s))} \\ & = V(q_{2l-1}^*(s), s, q_{2l-1}^*(s); a_*(s, q_{2l-1}^*(s)), b_*(s, q_{2l-1}^*(s))), \end{aligned} \tag{3.36}$$

$$\begin{aligned} & \sum_{i=1}^2 C_i(s, q_{2l-1}^*(s) + ; b_*(s, q_{2l-1}^*(s) + )) \gamma_i(s, q_{2l-1}^*(s)) (q_{2l-1}^*(s))^{\gamma_i(s, q_{2l-1}^*(s))} \\ & = q_{2l-1}^*(s) \partial_x V(q_{2l-1}^*(s), s, q_{2l-1}^*(s); a_*(s, q_{2l-1}^*(s)), b_*(s, q_{2l-1}^*(s))), \end{aligned} \tag{3.37}$$

where the functions  $C_i(s, q; b_*(s, q))$ , for  $i = 1, 2$ , are given by (3.18) and the function  $V(x, s, q; a_*(s, q), b_*(s, q))$  has the form of (3.13) with (3.14), while the boundary  $b_*(s, q)$  provides a unique solution of the first-order nonlinear ordinary differential equation in (3.27) in the region  $\widehat{R}_{2l}(b_*)$  from (3.35) satisfying the (starting) conditions of (3.36)–(3.37) above, for each  $s > 0$  fixed and every  $k \in \mathbb{N}$ . Note that the process  $(X, S, Q)$  cannot come from the region  $\widehat{R}_{2k}(a_*)$  in (3.31) directly to the region  $\widehat{R}_{2l}(b_*)$  in (3.35), or vice versa, without crossing the region  $\widehat{R}_{2k}(a_*, b_*)$  in (3.30) or  $\widehat{R}_{2l}(a_*, b_*)$  in (3.34), respectively, for every  $k, l \in \mathbb{N}$ .

Finally we observe that if we have  $\gamma_i(s, q) = \gamma_i(q)$ , for  $i = 1, 2$ , in (3.2), then the appropriate left-hand exercise boundary for the floating-strike lookback game option in (2.24)–(2.25) takes the form  $a_*(s, q) = a'(s, q) \vee g_*(q)s$  with some function  $0 < g_*(q) < 1$ , while, if we have  $\gamma_i(s, q) = \gamma_i(s)$ , for  $i = 1, 2$ , in (3.2), then the appropriate right-hand exercise boundary for that contract there takes the form  $b_*(s, q) = b'(s, q) \wedge h_*(s)q$  with some function  $h_*(s) > 1$ , for all  $0 < q < s$ . In these cases, we have solely the regions  $\widehat{R}_1(a_*, b_*)$ ,  $\widehat{R}_2(a_*)$  in (3.30)–(3.31) with  $k = 1$  and  $\widehat{R}_1(a_*, b_*)$ ,  $\widehat{R}_2(b_*)$  in (3.34)–(3.35) with  $l = 1$ , respectively. Moreover, if we have  $\gamma_i(s, q) = \gamma_i$ , for  $i = 1, 2$ , in (3.2), then we have  $a_*(s, q) = a'(s, q) \vee a_*(s) = a'(s, q) \vee g_*s$  and  $b_*(s, q) = b'(s, q) \wedge b_*(q) = b'(s, q) \wedge h_*q$  with some constants  $0 < g_* < 1$  and  $h_* > 1$ , for all  $0 < q < s$ . The latter property can be explained by the fact that the original problem of (1.4), and thus the three-dimensional problem of (2.3), can be reduced to an optimal stopping zero-sum game for the process  $(S/X, Q/X) = (S_t/X_t, Q_t/X_t)_{t \geq 0}$  representing a two-dimensional Markov diffusion process with reflection, by means of the change-of-measure arguments from [65] (see also [21]).

### 3.4. The system of arithmetic equations for the boundaries

Let us now extend the arguments from [26, Example 4.2] (see also [18, Section 3] and [60, Theorem 1]) to show that the system of arithmetic power equations in (3.21) admits a unique solution. For this purpose, by virtue of straightforward calculations, we first observe that the system of equations in (3.21) is equivalent to the following one:

$$\Xi_i(a(s, q); s, q) = \Upsilon_i(b(s, q); s, q) \tag{3.38}$$

for  $a'(s, q) \vee q \leq a(s, q) < b(s, q) \leq s \wedge b'(s, q)$ , where we set

$$\Xi_i(a; s, q) = \frac{\gamma_i(s, q)(s - Ka) + Ka}{a^{\gamma_3-i(s, q)}} \quad \text{and} \quad \Upsilon_i(b; s, q) = \frac{\gamma_i(s, q)(q - Lb) + Lb}{b^{\gamma_3-i(s, q)}} \tag{3.39}$$

for all  $0 < q < a < b < s$  and every  $i = 1, 2$ .

In order to show the existence and uniqueness of a solution of the system of arithmetic power equations in (3.38) with (3.39), we develop the ideas used to prove the existence and uniqueness of solutions to the systems of arithmetic power equations in [26, Example 4.2] (see also the systems (4.73)–(4.74) in [67, Chapter IV, Section 2], the system (3.16)–(3.17) in [18, Section 3], and [60, Theorem 1]). For this purpose, we observe that, for the derivatives of the functions  $\Xi_i(a)$  and  $\Upsilon_i(b)$ , for  $i = 1, 2$ , defined in (3.38), the equations

$$\Xi'_i(a; s, q) = \frac{(\gamma_1(s, q) - 1)(\gamma_2(s, q) - 1)K(a - \bar{a}(s, q))}{a^{\gamma_3-i(s, q)+1}} \tag{3.40}$$

and

$$\Upsilon'_i(b; s, q) = \frac{(\gamma_1(s, q) - 1)(\gamma_2(s, q) - 1)L(b - \underline{b}(s, q))}{b^{\gamma_3-i(s, q)+1}} \tag{3.41}$$

hold, so that the inequalities

$$\Xi'_i(a; s, q) > 0 \quad \text{for} \quad a < \bar{a}(s, q) \quad \text{and} \quad \Upsilon'_i(b; s, q) < 0 \quad \text{for} \quad b > \underline{b}(s, q) \tag{3.42}$$

are satisfied, for  $a'(s, q) \vee q < a < b < s \wedge b'(s, q)$  and every  $i = 1, 2$ , where the functions  $a'(s, q)$  and  $b'(s, q)$  are given by (2.9). Hence, we may conclude that the following properties hold, for each  $0 < q < s$  fixed. The function  $\Xi_1(a; s, q)$  is increasing on the interval  $(0, \bar{a}(s, q))$ , with  $\Xi_1(0+; s, q) = 0$  and  $\Xi_1(\bar{a}(s, q); s, q) > 0$ , so that the range of its values is given by the interval  $(0, \Xi_1(\bar{a}(s, q); s, q))$ . The function  $\Upsilon_1(b; s, q)$  is decreasing on the interval  $(\underline{b}(s, q), \infty)$ , with  $\Upsilon_1(\underline{b}(s, q); s, q) > 0$  and  $\Upsilon_1(\infty; s, q) = -\infty$ , so that the range of its values is given by the interval  $(-\infty, \Upsilon_1(\underline{b}(s, q); s, q))$ . The function  $\Xi_2(a; s, q)$  is increasing on the interval  $(0, \bar{a}(s, q))$  with  $\Xi_2(0+; s, q) = -\infty$  and  $\Xi_2(\bar{a}(s, q); s, q) > 0$ , so that the range of its values is given by the interval  $(-\infty, \Xi_2(\bar{a}(s, q); s, q))$ . The function  $\Upsilon_2(b; s, q)$  is decreasing on the interval  $(\underline{b}(s, q), \infty)$  with  $\Upsilon_2(\underline{b}(s, q); s, q) > 0$  and  $\Upsilon_2(\infty; s, q) = 0$ , so that the range of its values is given by the interval  $(0, \Upsilon_2(\underline{b}(s, q); s, q))$ .

We now observe that, when  $\Xi_i(\bar{a}(s, q); s, q) \leq \Upsilon_i(\underline{b}(s, q); s, q)$  holds, one can determine some  $\widehat{b}_i(s, q) \geq \underline{b}(s, q)$  from the equation  $\Xi_i(\bar{a}(s, q); s, q) = \Upsilon_i(\widehat{b}_i(s, q); s, q)$ , while, when  $\Xi_i(\bar{a}(s, q); s, q) \geq \Upsilon_i(\underline{b}(s, q); s, q)$  holds, one can determine some  $\widehat{a}_i(s, q) \leq \bar{a}(s, q)$  from the equation  $\Xi_i(\widehat{a}_i(s, q); s, q) = \Upsilon_i(\underline{b}(s, q); s, q)$ , for each  $0 < q < s$  fixed, and every  $i = 1, 2$ . Hence, it follows from the equations in (3.38) with (3.39) that, for each  $a \in (\underline{a}(s, q), \widehat{a}_1(s, q) \wedge \bar{a}(s, q))$ , there exists a unique number  $b \in (\underline{b}(s, q) \vee \widehat{b}_1(s, q), \bar{b}(s, q))$ , while, for each  $a \in$




FIGURE 3. A computer drawing of the boundary functions  $b_1(a)$  and  $b_2(a)$  in the case  $a'(s, q) \leq a_*(s, q) < b_*(s, q) \leq b'(s, q)$ , for each  $0 < q < s$  fixed.

( $\underline{a}(s, q), \widehat{a}_2(s, q) \wedge \bar{a}(s, q)$ ), there exists a unique number  $b \in (\underline{b}(s, q) \vee \widehat{b}_2(s, q), \bar{b}(s, q))$ , for each  $0 < q < s$  fixed. (Recall that the values  $\bar{a}(s, q)$  and  $\underline{b}(s, q)$  are specified in Part (ii) of Subsection 2.2 above and given by the expressions in (2.19), while the values  $\underline{a}(s, q)$  and  $\bar{b}(s, q)$  are determined in Theorem 2 below, for each  $0 < q < s$  fixed.)

In other words, we may conclude that the first and second equations in (3.38), respectively, uniquely determine the function  $b_1(a)$  on  $(\underline{a}(s, q), \widehat{a}_1(s, q) \wedge \bar{a}(s, q))$  with the range  $(\underline{b}(s, q) \vee \widehat{b}_1(s, q), \bar{b}(s, q))$  and the function  $b_2(a)$  on  $(\underline{a}(s, q), \widehat{a}_2(s, q) \wedge \bar{a}(s, q))$  with the range  $(\underline{b}(s, q) \vee \widehat{b}_2(s, q), \bar{b}(s, q))$ , for each  $0 < q < s$  fixed. These arguments also imply that the expression  $\underline{b}(s, q) \vee \widehat{b}_1(s, q) < b_1(\underline{a}(s, q)) < \bar{b}(s, q) < b_2(\underline{a}(s, q))$  holds. Moreover, the same arguments and assumptions directly yield that there exists exactly one intersection point, with the coordinates  $a_*(s, q)$  and  $b_*(s, q)$ , of the curves associated with the functions  $b_1(a)$  and  $b_2(a)$  on the interval  $a \in (\underline{a}(s, q), \widehat{a}_2(s, q) \wedge \bar{a}(s, q))$  such that  $\underline{b}(s, q) \vee \widehat{b}_1(s, q) < b_1(a_*(s, q)) \equiv b_*(s, q) \equiv b_2(a_*(s, q)) < \bar{b}(s, q)$  holds, for each  $0 < q < s$  fixed (see Figure 3 above).

More precisely, let us assume that there exist at least two intersection points,  $(a_*(s, q), b_*(s, q))$  and  $(\tilde{a}(s, q), \tilde{b}(s, q))$ , of the curves  $b_1(a)$  and  $b_2(a)$ , such that  $\underline{a}(s, q) < \tilde{a}(s, q) < a_*(s, q) \leq \bar{a}(s, q) \wedge \widehat{a}_2(s, q)$  and  $\underline{b}(s, q) \vee \widehat{b}_1(s, q) \leq \tilde{b}(s, q) < b_*(s, q) < \bar{b}(s, q)$  (or  $\underline{a}(s, q) < a_*(s, q) < \tilde{a}(s, q) \leq \bar{a}(s, q) \wedge \widehat{a}_2(s, q)$  and  $\underline{b}(s, q) \vee \widehat{b}_1(s, q) \leq b_*(s, q) < \tilde{b}(s, q) < \bar{b}(s, q)$ ), as well as  $b_2(a) > b_1(a)$ , for  $a \in (\tilde{a}(s, q), a_*(s, q))$  and any  $0 < q < s$  fixed. Observe that, by virtue of the assumptions made above and according to the implicit function theorem, it follows from the representations in (3.40)–(3.41) that the expressions

$$b'_i(a) = \frac{\Xi'_i(a)}{\Upsilon'_i(b)} = \frac{K(a - \bar{a}(s, q))}{L(b - \underline{b}(s, q))} \left(\frac{b}{a}\right)^{\gamma_{3-i}(s, q)+1} < 0 \tag{3.43}$$

hold, for every  $i = 1, 2$ , for all  $a \in (\tilde{a}(s, q), a_*(s, q))$  and  $b \in (\tilde{b}(s, q), b_*(s, q))$ , from which it directly follows that the inequality

$$\frac{b'_2(a)}{b'_1(a)} = \left(\frac{b}{a}\right)^{\gamma_1(s, q) - \gamma_2(s, q)} > 1 \tag{3.44}$$

is satisfied for all  $a \in (\tilde{a}(s, q), a_*(s, q))$ . Since the derivatives  $b'_i(a)$ , for  $i = 1, 2$ , from (3.43) are continuous functions on  $(\tilde{a}(s, q), a_*(s, q))$ , we may conclude that there exists an open interval  $(\tilde{a}(s, q) - \varepsilon, \tilde{a}(s, q) + \varepsilon)$ , for some relatively small  $\varepsilon > 0$ , such that the inequality  $b'_2(a) > b'_1(a)$  holds, so that the inequality  $b_2(a) > b_1(a)$  should hold for  $a \in (\tilde{a}(s, q) - \varepsilon, \tilde{a}(s, q) + \varepsilon)$  too. However, the latter fact contradicts the assumption that the equality  $b_1(\tilde{a}(s, q)) = b_2(\tilde{a}(s, q))$  holds, which means that the curves  $b_1(a)$  and  $b_2(a)$  may have only one intersection point; this completes the proof of the claim.

Furthermore, we recall that the functions  $a_*(s, q)$  and  $b_*(s, q)$  determined by means of the arguments above provide the candidate optimal stopping boundaries whenever the inequalities  $a'(s, q) \leq a_*(s, q) < b_*(s, q) \leq b'(s, q)$  hold for the solution of the system in (3.38) with (3.39), with  $a'(s, q)$  and  $b'(s, q)$  from (2.9), for each  $0 < q < s$  fixed. Otherwise, on the one hand, it can be shown by means of arguments similar to the ones used above that, if the inequalities  $a'(s, q) \leq a_*(s, q) < b'(s, q) < b_*(s, q)$  hold for the solution of the system in (3.38) with (3.39), then the right-hand candidate stopping boundary  $b_*(s, q)$  should coincide with  $b'(s, q)$ , while the left-hand candidate stopping boundary  $a_*(s, q) < b'(s, q)$  represents the smallest root (or the minimal solution) of the arithmetic equation in (3.22), which takes the form

$$\sum_{i=1}^2 (-1)^i \left( \frac{\gamma_{3-i}(s, q)s}{b'(s, q)} + (1 - \gamma_{3-i}(s, q))K \frac{a(s, q)}{b'(s, q)} \right) \left( \frac{a(s, q)}{b'(s, q)} \right)^{-\gamma_i(s, q)} \tag{3.45}$$

$$= (\gamma_1(s, q) - \gamma_2(s, q))(q/b'(s, q) - L)$$

for  $0 < q < s$ . On the other hand, it can be shown by means of arguments similar to the ones used above that, if the inequalities  $a_*(s, q) < a'(s, q) < b_*(s, q) \leq b'(s, q)$  hold for the solution of the system in (3.38) with (3.39), then the left-hand candidate stopping boundary  $a_*(s, q)$  should coincide with  $a'(s, q)$ , while the right-hand candidate stopping boundary  $b_*(s, q) > a'(s, q)$  represents the largest root (or the maximal solution) of the arithmetic equation in (3.23), which takes the form

$$\sum_{i=1}^2 (-1)^i \left( \frac{\gamma_{3-i}(s, q)q}{a'(s, q)} + (1 - \gamma_{3-i}(s, q))L \frac{b(s, q)}{a'(s, q)} \right) \left( \frac{b(s, q)}{a'(s, q)} \right)^{-\gamma_i(s, q)} \tag{3.46}$$

$$= (\gamma_1(s, q) - \gamma_2(s, q))(s/a'(s, q) - K)$$

for  $0 < q < s$ . We finally note that, in the case in which neither the system of arithmetic equations in (3.38) with (3.39) nor the equations in (3.45) and (3.46) admit solutions for which either the inequalities  $a_*(s, q) \geq a'(s, q)$  or  $b_*(s, q) \leq b'(s, q)$  hold, respectively, we may conclude that both the candidate stopping boundaries  $a_*(s, q)$  and  $b_*(s, q)$  should coincide with the boundaries  $a'(s, q)$  and  $b'(s, q)$ , for  $0 < q < s$ , respectively.

#### 4. Main results and proofs

In this section, building on the facts proved above, we formulate and prove the main result of the paper, which concerns the three-dimensional optimal stopping zero-sum game of (2.3) in the model from (1.1)–(1.2) and (1.3).

**Theorem 1.** *Let the process  $(X, S, Q)$  be defined in (1.1)–(1.2) and (1.3), where  $r > 0$  is a constant, and  $\delta(s, q) > 0$  and  $\sigma(s, q) > 0$  are continuously differentiable bounded functions on  $[0, \infty]^2$ . Assume that the function  $\delta(s, q)$  is increasing in both the variables  $s$  and  $q$  on  $[0, \infty]^2$ .*




FIGURE 4. A computer drawing of the value function  $V_*(x, s, q)$  and optimal exercise boundaries  $a'(s, q) < a_*(s, q) < b_*(s, q) < b'(s, q)$ , for each  $0 < q < s$  fixed.

Then the value function in (2.3) of the perpetual floating-strike lookback game option takes the form

$$V_*(x, s, q) = \begin{cases} V(x, s, q; a_*(s, q), b_*(s, q)), & \text{if } 0 < q \leq a_*(s, q) < x < b_*(s, q) \leq s, \\ V(x, s, q; a_*(s, q)), & \text{if } 0 < q \leq a_*(s, q) < x \leq s < b_*(s, q), \\ V(x, s, q; b_*(s, q)), & \text{if } 0 < a_*(s, q) < q \leq x < b_*(s, q) \leq s, \\ F(x, s), & \text{if } 0 < q \leq x \leq a_*(s, q) < s, \\ G(x, s, q), & \text{if } 0 < q < b_*(s, q) \leq x \leq s, \end{cases} \quad (4.1)$$

where the functions  $F(x, s)$  and  $G(x, s, q)$  are defined in (1.5), for some  $0 < L < K < L + 1$  given and fixed, and the optimal exercise times forming a Nash equilibrium in the game are given by

$$\tau_* = \inf \{t \geq 0 \mid X_t \leq a_*(S_t, Q_t)\} \quad \text{and} \quad \zeta_* = \inf \{t \geq 0 \mid X_t \geq b_*(S_t, Q_t)\}, \quad (4.2)$$

where the stopping boundaries satisfy the inequalities  $a'(s, q) \vee \underline{a}(s, q) \leq a_*(s, q) \leq \bar{a}(s, q) \wedge s \wedge b'(s, q)$  and  $a'(s, q) \vee q \vee \underline{b}(s, q) \leq b_*(s, q) \leq b(s, q) \wedge b'(s, q)$ , with  $a'(s, q)$  and  $b'(s, q)$  given by (2.9) and with  $\bar{a}(s, q)$  and  $\underline{b}(s, q)$  given by (2.19), while  $\underline{a}(s, q)$  and  $\bar{b}(s, q)$  are determined in Theorem 2 below. The candidate value functions are further specified as follows:

- (i) The function  $V(x, s, q; a_*(s, q), b_*(s, q))$  is given by (3.13)–(3.14), while either the boundaries  $a_*(s, q)$  and  $b_*(s, q)$  provide a unique solution to the system of arithmetic equations in (3.21) whenever  $a'(s, q) \leq a_*(s, q) < b_*(s, q) \leq b'(s, q)$  (see Figure 4 above); or  $b_*(s, q) = b'(s, q)$  and  $a_*(s, q)$  provides the smallest root of the arithmetic equation in (3.22) whenever  $a'(s, q) \leq a_*(s, q) < b'(s, q)$  (see Figure 5 above); or  $a_*(s, q) = a'(s, q)$  and  $b_*(s, q)$  provides the largest root of the arithmetic equation in (3.23) whenever  $b'(s, q) \geq b_*(s, q) > a'(s, q)$  (see Figure 6 above); or  $a_*(s, q) = a'(s, q)$




FIGURE 5. A computer drawing of the value function  $V_*(x, s, q)$  and optimal exercise boundaries  $a'(s, q) = a_*(s, q) < b_*(s, q) < b'(s, q)$ , for each  $0 < q < s$  fixed.




FIGURE 6. A computer drawing of the value function  $V_*(x, s, q)$  and optimal exercise boundaries  $a'(s, q) < a_*(s, q) < b_*(s, q) = b'(s, q)$ , for each  $0 < q < s$  fixed.

and  $b_*(s, q) = b'(s, q)$ , in the regions  $\tilde{R}_{2k-1}(a_*, b_*)$  and  $\hat{R}_{2l-1}(a_*, b_*)$  from (3.30) and (3.34), for  $k, l \in \mathbb{N}$ , for  $0 < q < s$ .

- (ii) The function  $V(x, s, q; a_*(s, q))$  is given by (3.15)–(3.16), while the boundary  $a_*(s, q)$  either provides a unique solution to the first-order nonlinear ordinary differential equation in (3.24) started at  $(s_{2k-1}^*(q), s_{2k-1}^*(q), q)$ , whenever  $a_*(s, q) \geq a'(s, q)$ , or coincides with  $a'(s, q)$ , in the regions  $\tilde{R}_{2k}(a_*)$  from (3.31), for  $k \in \mathbb{N}$ , for  $0 < q < s$ .

(iii) The function  $V(x, s, q; b_*(s, q))$  is given by (3.17)–(3.18), while the boundary  $b_*(s, q)$  either provides a unique solution to the first-order nonlinear ordinary differential equation in (3.27) started at  $(q_{2l-1}^*(s), s, q_{2l-1}^*(s))$ , whenever  $b_*(s, q) \leq b'(s, q)$ , or coincides with  $b'(s, q)$ , in the regions  $\tilde{R}_{2l}(b_l^*)$  from (3.35), for  $l \in \mathbb{N}$ , for  $0 < q < s$ .

Observe that we can put  $s = q = x$  to obtain the value of the original perpetual floating-strike lookback game option pricing problem of (1.4) from the value function of the optimal stopping zero-sum game of (2.3).

*Proof.* In order to verify the assertion stated above, it remains for us to show that the function defined in (4.1) coincides with the value function in (2.3) and that the stopping times  $\tau_*$  and  $\zeta_*$  in (4.2) form a Nash equilibrium with the boundaries  $a_*(s, q)$  and  $b_*(s, q)$  specified in the previous section. For this purpose, let us denote by  $V(x, s, q)$  the right-hand side of the expression in (4.1) associated with these boundaries  $a_*(s, q)$  and  $b_*(s, q)$ . Then it follows from the straightforward calculations presented in the previous section that the function  $V(x, s, q)$  solves the system of (2.28)–(2.30), while the smooth-fit and normal-reflection conditions of (2.33)–(2.34) are satisfied in the appropriate regions  $\tilde{R}_{2k-1}(a_*, b_*)$ ,  $\tilde{R}_{2k}(a_*)$  from (3.30)–(3.31), for  $k \in \mathbb{N}$ , and  $\tilde{R}_{2l-1}(a_*, b_*)$ ,  $\tilde{R}_{2l}(b_*)$  from (3.34)–(3.35), for  $l \in \mathbb{N}$ , respectively. We also observe that the function  $V(x, s, q)$  is  $C^{2,1,1}$  on the closure  $\bar{C}_*$  of  $C_*$  from (2.24) and  $D_*$  from (2.25), by construction. Hence, taking into account the fact that the boundaries  $a_*(s, q)$  and  $b_*(s, q)$  are assumed to be continuously differentiable, for  $0 < q < s$ , by applying the change-of-variable formula from [54, Theorem 3.1] (see also [59, Chapter II, Section 3.5] for a summary of the related results and further references) to the process  $e^{-r(t \wedge \theta)} V(X_{t \wedge \theta}, S_{t \wedge \theta}, Q_{t \wedge \theta})$ , we deduce that

$$\begin{aligned}
 e^{-r(t \wedge \theta)} V(X_{t \wedge \theta}, S_{t \wedge \theta}, Q_{t \wedge \theta}) &= V(x, s, q) + M_{t \wedge \theta} \tag{4.3} \\
 + \int_0^{t \wedge \theta} e^{-ru} (\mathbb{L}V - rV)(X_u, S_u, Q_u) I(X_u \neq S_u, X_u \neq Q_u, X_u \neq a_*(S_u, Q_u), X_u \neq b_*(S_u, Q_u)) du \\
 + \int_0^{t \wedge \theta} e^{-ru} \partial_s V(X_u, S_u, Q_u) I(X_u = S_u) dS_u + \int_0^{t \wedge \theta} e^{-ru} \partial_q V(X_u, S_u, Q_u) I(X_u = Q_u) dQ_u,
 \end{aligned}$$

for all  $t \geq 0$ , holds with  $\theta$  defined in (2.10). Here, the process  $(M_{t \wedge \theta})_{t \geq 0}$  defined by

$$M_{t \wedge \theta} = \int_0^{t \wedge \theta} e^{-ru} \partial_x V(X_u, S_u, Q_u) I(X_u \neq S_u, X_u \neq Q_u) \sigma(S_u, Q_u) X_u dB_u \tag{4.4}$$

is a continuous local martingale under the probability measure  $\mathbb{P}_{x,s,q}$ . Note that, since the time spent by the process  $(X, S, Q)$  at the boundary surfaces  $\{(x, s, q) \in E' \mid x = a_*(s, q)\}$  and  $\{(x, s, q) \in E' \mid x = b_*(s, q)\}$  and at the planes  $d_1 = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < q \leq x = s\}$  and  $d_2 = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < q = x \leq s\}$  is of Lebesgue measure zero (see e.g. [10, Chapter II, Section 1]), the indicators in the second line of the formula of (4.3) and in the formula of (4.4) can be ignored. Moreover, since the process  $S$  increases only when the process  $(X, S, Q)$  is located on the plane  $d_1$ , while the process  $Q$  decreases only when the process  $(X, S, Q)$  is located on the plane  $d_2$ , the indicators in the third line of (4.3) can be set equal to one. Finally, taking into account the fact that the candidate value function  $V(x, s, q)$  satisfies the normal-reflection conditions of (2.34) at the diagonals  $d_1$  and  $d_2$  in the regions  $\tilde{R}_{2k}(a_*)$ , for  $k \in \mathbb{N}$ , in (3.35) and  $\tilde{R}_{2l}(b_*)$ , for  $l \in \mathbb{N}$ , we may conclude that the integrals in the third line of (4.3) are actually equal to zero.

By using straightforward calculations and the arguments from the previous section, it can be verified that the inequality  $(\mathbb{L}V - rV)(x, s, q) \leq 0$  holds for all  $a'(s, q) < x < b_*(s, q)$  such that  $x \neq a_*(s, q)$ , and the inequality  $(\mathbb{L}V - rV)(x, s, q) \geq 0$  holds for all  $x > a_*(s, q)$  such that  $x \neq b_*(s, q)$ , as well as  $x \neq s$  and  $x \neq q$ . Moreover, we observe directly from the expressions in (3.13), (3.15), and (3.17) that the function  $V(x, s, q) - (s - Kx)$  is increasing in the variable  $x$  on the interval  $q \vee a_*(s, q) < x < b_*(s, q) \wedge s$ , from 0 (when  $a_*(s, q) \geq q$ ) to the value  $q - s + (K - L)b_*(s, q)$  (when  $b_*(s, q) \leq s$ ), because the expression  $\partial_x V(x, s, q) + K$  for the first-order partial derivative in (3.19) is positive there, for  $(x, s, q) \in E'$ . We also note from the expressions in (3.13), (3.15), and (3.17) that the function  $V(x, s, q)$  is convex in the variable  $x$  in a left-hand neighborhood of  $q \vee a_*(s, q)$  and concave in a right-hand neighborhood of  $b_*(s, q) \wedge s$ , because its second-order partial derivative  $\partial_{xx} V(x, s, q)$  is positive in a left-hand neighborhood of  $q \vee a_*(s, q)$  and negative in a right-hand neighborhood of  $b_*(s, q) \wedge s$ , for  $(x, s, q) \in E'$ . Thus, we may conclude that the inequalities in (2.31) hold, which together with the conditions of (2.29)–(2.30) and (2.33) imply that the inequalities  $s - Kx \leq V(x, s, q) \leq q - Lx$  are satisfied, for all  $(x, s, q) \in E'$ , under  $0 < L < K < L + 1$  given and fixed. It therefore follows from the expression in (4.3) that the inequalities

$$e^{-r(\tau \wedge \zeta_*)} F(X_{\tau \wedge \zeta_*}, S_{\tau \wedge \zeta_*}) \leq e^{-r(\tau \wedge \zeta_*)} V(X_{\tau \wedge \zeta_*}, S_{\tau \wedge \zeta_*}, Q_{\tau \wedge \zeta_*}) \leq V(x, s, q) + M_{\tau \wedge \zeta_*} \tag{4.5}$$

and

$$e^{-r(\tau_* \wedge \zeta)} G(X_{\tau_* \wedge \zeta}, S_{\tau_* \wedge \zeta}, Q_{\tau_* \wedge \zeta}) \geq e^{-r(\tau_* \wedge \zeta)} V(X_{\tau_* \wedge \zeta}, S_{\tau_* \wedge \zeta}, Q_{\tau_* \wedge \zeta}) \geq V(x, s, q) + M_{\tau_* \wedge \zeta} \tag{4.6}$$

hold for any stopping times  $\tau$  and  $\zeta$  of the process  $(X, S, Q)$ , because  $\tau_* \wedge \zeta_* \leq \theta$  ( $\mathbb{P}_{x,s,q}$ -a.s.) holds with  $\theta$  as defined in (2.10).

Now, consider the localising sequence  $(\varkappa_n)_{n \in \mathbb{N}}$  for the local martingale  $(M_{t \wedge \theta})_{t \geq 0}$  from (4.4) such that  $\varkappa_n = \inf\{t \geq 0 \mid |M_{t \wedge \theta}| \geq n\}$ , for each  $n \in \mathbb{N}$ . Then, inserting  $\tau \wedge \varkappa_n$  and  $\zeta \wedge \varkappa_n$  instead of  $\tau$  and  $\zeta$  in (4.5) and (4.6) and taking the expectations with respect to the probability measure  $\mathbb{P}_{x,s,q}$  in (4.5) and (4.6), by means of Doob’s optional sampling theorem, we get

$$\begin{aligned} & \mathbb{E}_{x,s,q} \left[ e^{-r(\tau \wedge \zeta_* \wedge \varkappa_n)} \left( F(X_{\tau \wedge \varkappa_n}, S_{\tau \wedge \varkappa_n}) I(\tau \wedge \varkappa_n < \zeta_*) + G(X_{\zeta_*}, S_{\zeta_*}, Q_{\zeta_*}) I(\zeta_* \leq \tau \wedge \varkappa_n) \right) \right] \\ & \leq \mathbb{E}_{x,s,q} \left[ e^{-r(\tau \wedge \zeta_* \wedge \varkappa_n)} V(X_{\tau \wedge \zeta_* \wedge \varkappa_n}, S_{\tau \wedge \zeta_* \wedge \varkappa_n}, Q_{\tau \wedge \zeta_* \wedge \varkappa_n}) \right] \\ & \leq V(x, s, q) + \mathbb{E}_{x,s,q} [M_{\tau \wedge \zeta_* \wedge \varkappa_n}] = V(x, s, q) \end{aligned} \tag{4.7}$$

and

$$\begin{aligned} & \mathbb{E}_{x,s,q} \left[ e^{-r(\tau_* \wedge \zeta \wedge \varkappa_n)} \left( F(X_{\tau_*}, S_{\tau_*}) I(\tau_* < \zeta \wedge \varkappa_n) + G(X_{\zeta \wedge \varkappa_n}, S_{\zeta \wedge \varkappa_n}, Q_{\zeta \wedge \varkappa_n}) I(\zeta \wedge \varkappa_n \leq \tau_*) \right) \right] \\ & \geq \mathbb{E}_{x,s,q} \left[ e^{-r(\tau_* \wedge \zeta \wedge \varkappa_n)} V(X_{\tau_* \wedge \zeta \wedge \varkappa_n}, S_{\tau_* \wedge \zeta \wedge \varkappa_n}, Q_{\tau_* \wedge \zeta \wedge \varkappa_n}) \right] \\ & \geq V(x, s, q) + \mathbb{E}_{x,s,q} [M_{\tau_* \wedge \zeta \wedge \varkappa_n}] = V(x, s, q) \end{aligned} \tag{4.8}$$

for all  $(x, s, q) \in E'$  and each  $n \in \mathbb{N}$ . Observe that, taking into account the arguments from [64, pp. 635–636], it follows from the structure of the stopping times  $\tau_*$  and  $\zeta_*$  in (4.2) that the property

$$\mathbb{E}_{x,s,q} \left[ \sup_{t \geq 0} e^{-r((\tau_* \wedge \zeta_*) \wedge t)} S_{(\tau_* \wedge \zeta_*) \wedge t} \right] = \mathbb{E}_{x,s,q} \left[ \sup_{t \geq 0} e^{-r((\tau_* \wedge \zeta_*) \wedge t)} (X_{(\tau_* \wedge \zeta_*) \wedge t} \vee s) \right] < \infty \tag{4.9}$$



holds, and the variables  $e^{-r(\tau_* \wedge \zeta_*)}(S_{\tau_* \wedge \zeta_*} - KX_{\tau_* \wedge \zeta_*})$  and  $e^{-r(\tau_* \wedge \zeta_*)}(Q_{\tau_* \wedge \zeta_*} - LX_{\tau_* \wedge \zeta_*})$  are finite on the set  $\{\tau_* \wedge \zeta_* = \infty\}$  ( $\mathbb{P}_{x,s,q}$ -a.s.); moreover,  $\mathbb{P}_{x,s,q}(\tau_* \wedge \zeta_* < \infty) = 1$  for all  $(x, s, q) \in E'$ . Hence, letting  $n$  go to infinity and using Fatou's lemma, we obtain that the inequalities

$$\mathbb{E}_{x,s,q}[e^{-r(\tau \wedge \zeta_*)} (F(X_\tau, S_\tau) I(\tau < \zeta_*) + G(X_{\zeta_*}, S_{\zeta_*}, Q_{\zeta_*}) I(\zeta_* \leq \tau))] \leq V(x, s, q) \tag{4.10}$$

and

$$\mathbb{E}_{x,s,q}[e^{-r(\tau_* \wedge \zeta)} (F(X_{\tau_*}, S_{\tau_*}) I(\tau_* < \zeta) + G(X_\zeta, S_\zeta, Q_\zeta) I(\zeta \leq \tau_*))] \geq V(x, s, q) \tag{4.11}$$

hold for any stopping times  $\tau$  and  $\zeta$  such that  $\tau \wedge \zeta \leq \theta$  ( $\mathbb{P}_{x,s,q}$ -a.s.) holds, and for all  $(x, s, q) \in E'$ . Therefore, using the fact that the function  $V(x, s, q)$  and the continuously differentiable boundaries  $a_*(s, q)$  and  $b_*(s, q)$  solve the second-order ordinary differential equation in (2.28) and satisfy the conditions of (2.29)–(2.30) and (2.33)–(2.34), inserting  $\tau_*$  in place of  $\tau$  and  $\zeta_*$  in place of  $\zeta$  into (4.10) and (4.11), we obtain that the equality

$$\mathbb{E}_{x,s,q}[e^{-r(\tau_* \wedge \zeta_*)} (F(X_{\tau_*}, S_{\tau_*}) I(\tau_* < \zeta_*) + G(X_{\zeta_*}, S_{\zeta_*}, Q_{\zeta_*}) I(\zeta_* \leq \tau_*))] = V(x, s, q) \tag{4.12}$$

holds, so that the candidate function  $V(x, s, q)$  coincides with the value function  $V_*(x, s, q)$  of the optimal stopping game in (2.3) for all  $(x, s, q) \in E'$ , and the optimal stopping times  $\tau_*$  and  $\zeta_*$  form a Nash equilibrium of the zero-sum game. Finally, we recall from the results of Part (ii) of Subsection 2.2 above, which are implied by standard comparison arguments applied to the value functions of the appropriate optimal stopping problems, that the inequalities  $a'(s, q) \vee \underline{a}(s, q) \leq a_*(s, q) < \bar{a}(s, q) \wedge s \wedge b'(s, q)$  and  $a'(s, q) \vee q \vee \underline{b}(s, q) < b_*(s, q) \leq \bar{b}(s, q) \wedge b'(s, q)$  should hold, for  $0 < q < s$ . This completes the verification.  $\square$

### 5. Appendix

In this section, we derive closed-form expressions for the value functions and optimal stopping boundaries of some auxiliary optimal stopping problems, which provide upper and lower bounds for the value function and optimal stopping boundaries of the original optimal stopping zero-sum game of (2.3).

#### 5.1. The optimal stopping and free-boundary problems

In order to provide the upper and lower bounds for the value functions and optimal stopping boundaries in the optimal stopping game of (2.3) above, let us introduce the value functions  $\bar{V}(x, s, q)$  and  $\underline{V}(x, s, q)$  of the optimal stopping problems

$$\bar{V}(x, s, q) = \sup_{\tau} \mathbb{E}_{x,s,q}[e^{-r\tau} (S_\tau - KX_\tau)] \quad \text{and} \quad \underline{V}(x, s, q) = \inf_{\zeta} \mathbb{E}_{x,s,q}[e^{-r\zeta} (Q_\zeta - LX_\zeta)] \tag{5.1}$$

for some given constants  $K, L > 0$ , where the supremum and infimum are taken with respect to all stopping times  $\tau$  and  $\zeta$  of the process  $X$ . It can be shown by means of arguments similar to the ones used in Part (ii) of Subsection 2.2, based on the representations (2.14) and (2.15) for the reward functional, under the assumptions  $\zeta = \infty$  and  $\tau = \infty$ , respectively, or by using the (easily proved) convexity and concavity of the value functions  $\bar{V}(x, s, q)$  and  $\underline{V}(x, s, q)$ , that the optimal stopping times in the problems of (5.1) have the form

$$\bar{\tau} = \inf \{t \geq 0 \mid X_t \leq \underline{a}(S_t, Q_t)\} \quad \text{and} \quad \underline{\zeta} = \inf \{t \geq 0 \mid X_t \geq \bar{b}(S_t, Q_t)\} \tag{5.2}$$

with some functions  $\underline{a}(s, q)$  and  $\bar{b}(s, q)$ , for  $0 < q < s$ , to be determined (see [36, 51] as well as [23, Section 3] and [21]). By means of arguments similar to the ones applied in [13, Subsection 3.2] and [52, Proposition 2.1], the existence of such boundaries  $\underline{a}(s, q)$  and  $\bar{b}(s, q)$  can be explained by the facts that the costs for the holder (maximiser of the expected discounted payoff) of waiting until the process  $X$  from (1.1) coming from a small  $x > 0$  increases to the current value of the running maximum process  $S$  and the costs for the writer (minimiser of the expected discounted payoff) of waiting until the process  $X$  coming from a large  $x > 0$  decreases to the current value of the running minimum process  $Q$  may be too large, owing to the structure of the integrands in the reward functionals of (2.16) and (2.17), respectively.

Extending the arguments from [36, 51] (see also [23, Section 3] and [21]) to the three-dimensional model under consideration here, we may conclude that the unknown value functions  $\bar{V}(x, s, q)$  and  $\underline{V}(x, s, q)$  from (5.1) and the unknown boundaries  $\underline{a}(s, q)$  and  $\bar{b}(s, q)$  from (5.2) should solve the equivalent free-boundary problems

$$(\mathbb{L}V - rV)(x, s, q) = 0 \quad \text{for } q \vee a(s, q) < x < s \quad \text{or} \tag{5.3}$$

$$(\mathbb{L}V - rV)(x, s, q) = 0 \quad \text{for } q < x < b(s, q) \wedge s, \tag{5.4}$$

$$V(x, s, q)|_{x=a(s,q)+} = s - K a(s, q), \quad V(x, s, q)|_{x=b(s,q)-} = q - L b(s, q), \tag{5.5}$$

$$\partial_x V(x, s, q)|_{x=a(s,q)+} = -K, \quad \partial_x V(x, s, q)|_{x=b(s,q)-} = -L, \tag{5.6}$$

$$\partial_s V(x, s, q)|_{x=s-} = 0, \quad \partial_q V(x, s, q)|_{x=q+} = 0, \tag{5.7}$$

$$V(x, s, q) = s - Kx \quad \text{for } x < a(s, q), \quad V(x, s, q) = q - Lx \quad \text{for } x > b(s, q), \tag{5.8}$$

$$V(x, s, q) > s - Kx \quad \text{for } a(s, q) < x \leq s, \quad V(x, s, q) < q - Lx \quad \text{for } q \leq x < b(s, q), \tag{5.9}$$

$$(\mathbb{L}V - rV)(x, s, q) < 0 \quad \text{for } x < a(s, q), \quad (\mathbb{L}V - rV)(x, s, q) > 0 \quad \text{for } x > b(s, q), \tag{5.10}$$

where the conditions in (5.5)–(5.7) are satisfied, for each  $0 < q < s$ .

### 5.2. Solutions to the free-boundary problems

It follows from the arguments of [36, 51] (see also [23, Section 3] and [21]) that the solution to the left-hand system in (5.3) and (5.5)–(5.10) has the form of (3.15) with (3.16), for  $0 < q \leq a(s, q) < x \leq s$ , while the boundary  $a(s, q)$  solves the first-order nonlinear ordinary differential equation in (3.24), for any  $q > 0$  fixed. We also see that the solution to the right-hand system in (5.4) and (5.5)–(5.10) has the form of (3.17) with (3.18), for  $0 < q \leq x < b(s, q) \leq s$ , while the boundary  $b(s, q)$  solves the first-order nonlinear ordinary differential equation in (3.27), for any  $s > 0$  fixed.

We further define the *maximal and minimal admissible* solutions of the first-order nonlinear ordinary differential equations as the largest and smallest possible solutions  $\underline{a}(s, q)$  and  $\bar{b}(s, q)$  of the equations in (3.24) and (3.27) which satisfy the inequalities  $\underline{a}(s, q) < s$  and  $\bar{b}(s, q) > q$  for all  $0 < q < s$ . By virtue of the classical results on the existence and uniqueness of solutions for first-order nonlinear ordinary differential equations, we may conclude that these equations admit (locally) unique solutions, in view of the fact that the right-hand sides in (3.24) and

(3.27) are (locally) continuous in  $(s, q, a(s, q))$  and  $(s, q, b(s, q))$  and (locally) Lipschitz in  $a(s, q)$  and  $b(s, q)$ , for each  $(s, q)$  fixed (see [52, Subsection 3.9] for similar arguments based on the analysis of other first-order nonlinear ordinary differential equations). It can then be shown by means of technical arguments based on Picard’s method of successive approximations that there exist unique solutions  $a(s, q)$  and  $b(s, q)$  to the equations in (3.24) and (3.27), started at some points  $(s_0, s_0, q)$  and  $(s, q_0, q_0)$  such that  $s_0 > 0$  and  $q_0 > 0$ , for each  $0 < q < s$  fixed (see also [34, Subsection 3.2] and [52, Example 4.4] for similar arguments based on the analysis of other first-order nonlinear ordinary differential equations).

Hence, in order to construct the appropriate functions  $\underline{a}(s, q)$  and  $\bar{b}(s, q)$  which satisfy the equations in (3.24) and (3.27) and stay strictly above or below the appropriate diagonal for  $0 < q < s$ , we can follow the arguments from [58, Subsection 3.5] (among others), which are based on the construction of sequences of so-called bad–good solutions which intersect the diagonals. For this purpose, for any positive sequences  $(s_k, q_k)_{k \in \mathbb{N}}$  and  $(s_l, q_l)_{l \in \mathbb{N}}$  such that  $s_k \uparrow \infty$  as  $k \rightarrow \infty$  and  $q_l \downarrow 0$  as  $l \rightarrow \infty$ , we can construct the sequence of solutions  $a_k(s, q)$ , for  $k \in \mathbb{N}$ , and  $b_l(s, q)$ , for  $l \in \mathbb{N}$ , to the equations (3.24) and (3.27) such that  $a_k(s_k, q_k) = s_k$  and  $b_l(s_l, q_l) = q_l$  holds, for each  $k, l \in \mathbb{N}$ . It follows from the structure of the equations in (3.24) and (3.27) that the properties  $\partial_s a_k(s_k, q_k) < 1$  and  $\partial_q b_l(s_l, q_l) > 1$  hold, for each  $k, l \in \mathbb{N}$  (see also [51, pp. 979–982] for the analysis of solutions of the non-parametrised version of the first-order nonlinear differential equation of (3.24)). Observe that, by virtue of the uniqueness of solutions mentioned above, we know that the two curves  $s \mapsto a_k(s, q)$  and  $s \mapsto a_m(s, q)$  cannot intersect, and similarly  $q \mapsto b_l(s, q)$  and  $q \mapsto b_n(s, q)$  cannot intersect, for  $l, k, m, n \in \mathbb{N}$  such that  $k \neq m$  and  $l \neq n$ ; thus, the sequence  $(a_k(s, q))_{k \in \mathbb{N}}$  is increasing and the sequence  $(b_l(s, q))_{l \in \mathbb{N}}$  is decreasing, so that the limits  $\underline{a}(s, q) = \lim_{k \rightarrow \infty} a_k(s, q)$  and  $\bar{b}(s, q) = \lim_{l \rightarrow \infty} b_l(s, q)$  exist, for each  $0 < q < s$ . We may therefore conclude that  $\underline{a}(s, q)$  and  $\bar{b}(s, q)$  provide the maximal and minimal solutions to the equations in (3.24) and (3.27) such that  $\underline{a}_k(s, q) < s$  holds for each  $k \in \mathbb{N}$  and  $\bar{b}_l(s, q) > q$  holds for each  $l \in \mathbb{N}$ , for all  $0 < q < s$ .

Moreover, since the right-hand sides of the first-order nonlinear ordinary differential equations in (3.24) and (3.27) are (locally) Lipschitz in  $s$  and  $q$ , for each  $0 < q < s$ , one can deduce by means of Gronwall’s inequality that the functions  $a_k(s, q)$  and  $b_l(s, q)$ , for each  $k, l \in \mathbb{N}$ , are continuous, so that the functions  $\underline{a}(s, q)$  and  $\bar{b}(s, q)$  are continuous too. The appropriate maximal admissible solutions of first-order nonlinear ordinary differential equations, and the associated maximality principle for solutions of optimal stopping problems, which is equivalent to the superharmonic characterisation of the payoff functions, were established in [52] and further developed in [5, 20, 23, 30, 32–37, 46, 50–51, 57–58, 63], as well as other subsequent papers (see [59, Chapter I; Chapter V, Section 17] for further references).

### 5.3. The results

Summarising the facts shown above, we state the following result, which can be proved by means of the same arguments as used for Theorem 1 above, in combination with the arguments from [21] (see [64]–[66] for the original derivation and [20]–[23] for the related comparison arguments).

**Theorem 2.** *Let the process  $(X, S, Q)$  be as defined in (1.1) and (1.3), where  $r > 0$  is a constant, and  $\delta(s, q) > 0$  and  $\sigma(s, q) > 0$  are continuously differentiable bounded functions on  $[0, \infty]^2$ . Assume that the function  $\delta(s, q)$  is increasing in both the variables  $s$  and  $q$  on  $[0, \infty]^2$ . Then the following assertions hold:*

- (i) The value function  $\bar{V}(x, s, q)$  of the left-hand optimal stopping problem in (5.1) with  $K > 0$  takes the form

$$\bar{V}(x, s, q) = \begin{cases} V(x, s, q; \underline{a}(s, q)), & \text{if } 0 < \underline{a}(s, q) < x \leq s, \\ s - Kx, & \text{if } 0 < x \leq \underline{a}(s, q), \end{cases} \quad (5.11)$$

and the optimal stopping time  $\bar{\tau}$  has the form of (5.2), where the function  $V(x, s, q; \underline{a}(s, q))$  is given by (3.15)–(3.16), while the optimal stopping boundary  $\underline{a}(s, q) [\leq rs/(K\delta(s, q))]$  provides the maximal solution of the first-order nonlinear ordinary differential equation in (3.24) staying below the diagonal  $d_1 = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < q \leq x = s\}$ , for  $0 < q < s$ .

- (ii) The value function  $\underline{V}(x, s, q)$  of the right-hand optimal stopping problem in (5.1) with  $L > 0$  takes the form

$$\underline{V}(x, s, q) = \begin{cases} V(x, s, q; \bar{b}(s, q)), & \text{if } 0 < q \leq x < \bar{b}(s, q), \\ q - Lx, & \text{if } x \geq \bar{b}(s, q), \end{cases} \quad (5.12)$$

and the optimal stopping time  $\underline{\zeta}$  has the form of (5.2), where the function  $V(x, s, q; \bar{b}(s, q))$  is given by (3.17)–(3.18), while the optimal stopping boundary  $\bar{b}(s, q) [\geq rq/(L\delta(s, q))]$  provides the minimal solution of the first-order nonlinear ordinary differential equation in (3.27) staying above the diagonal  $d_2 = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < q = x \leq s\}$ , for  $0 < q < s$ .

### Acknowledgements

The author is grateful to the editors and two anonymous referees for their patience and valuable suggestions, which provided essential help in improving the motivation and presentation of the paper.

### Funding information

There are no funding bodies to thank in relation to the creation of this article.

### Competing interests

There were no competing interests to declare which arose during the preparation or publication process of this article.

### References

- [1] ASMUSSEN, S., AVRAM, F. AND PISTORIUS, M. (2003). Russian and American put options under exponential phase-type Lévy models. *Stoch. Process. Appl.* **109**, 79–111.
- [2] AVRAM, F., KYPRIANOU, A. E. AND PISTORIUS, M. (2004). Exit problems for spectrally negative Lévy processes and applications to (Canadized) Russian options. *Ann. Appl. Prob.* **14**, 215–238.
- [3] BAURDOUX, E. J. AND KYPRIANOU, A. E. (2004). Further calculations for Israeli options. *Stochastics* **76**, 549–569.
- [4] BAURDOUX, E. J. AND KYPRIANOU, A. E. (2008). The McKean stochastic game driven by a spectrally negative Lévy process. *Electron. J. Prob.* **8**, 173–197.
- [5] BAURDOUX, E. J. AND KYPRIANOU, A. E. (2009). The Shepp–Shiryaev stochastic game driven by a spectrally negative Lévy process. *Theory Prob. Appl.* **53**, 481–499.
- [6] BAURDOUX, E. J., KYPRIANOU, A. E. AND PARDO, J. C. (2011). The Gapeev–Kühn stochastic game driven by a spectrally positive Lévy process. *Stoch. Process. Appl.* **121**, 1266–1289.

- [7] BENSOUSSAN, A. AND FRIEDMAN, A. (1974). Non-linear variational inequalities and differential games with stopping times. *J. Funct. Anal.* **16**, 305–352.
- [8] BENSOUSSAN, A. AND FRIEDMAN, A. (1977). Nonzero-sum stochastic differential games with stopping times and free-boundary problems. *Trans. Amer. Math. Soc.* **231**, 275–327.
- [9] BISMUT, J. M. (1977). Sur un problème de Dynkin. *Z. Wahrscheinlichkeitsthe.* **39**, 31–53.
- [10] BORODIN, A. N. AND SALMINEN, P. (2002). *Handbook of Brownian Motion*, 2nd edn. Birkhäuser, Basel.
- [11] CVITANIĆ, J. AND KARATZAS, I. (1996). Backward stochastic differential equations with reflection and Dynkin games. *Ann. Prob.* **24**, 2024–2056.
- [12] DETEMPLE, J. (2006). *American-Style Derivatives: Valuation and Computation*. Chapman and Hall/CRC, Boca Raton.
- [13] DUBINS, L., SHEPP, L. A. AND SHIRYAEV, A. N. (1993). Optimal stopping rules and maximal inequalities for Bessel processes. *Theory Prob. Appl.* **38**, 226–261.
- [14] EKSTRÖM, E. AND PESKIR, G. (2008). Optimal stopping games for Markov processes. *SIAM J. Control Optimization* **47**, 684–702.
- [15] EKSTRÖM, E. AND VILLENEUVE, S. (2006). On the value of optimal stopping games. *Ann. Appl. Prob.* **16**, 1576–1596.
- [16] FRIEDMAN, A. (1973). Stochastic games and variational inequalities. *Arch. Rational Mech. Anal.* **51**, 321–346.
- [17] FRIEDMAN, A. (1976). *Stochastic Differential Equations and Applications II*. Academic Press, New York.
- [18] GAPEEV, P. V. (2005). The spread option optimal stopping game. *Exotic Option Pricing and Advanced Levy Models*, eds A. Kyprianou, W. Schoutens and P. Wilmott, John Wiley, Chichester, pp. 293–305.
- [19] GAPEEV, P. V. (2006). Discounted optimal stopping for maxima in diffusion models with finite horizon. *Electron. J. Prob.* **11**, 1031–1048.
- [20] GAPEEV, P. V. (2007). Discounted optimal stopping for maxima of some jump-diffusion processes. *J. Appl. Prob.* **44**, 713–731.
- [21] GAPEEV, P. V. (2020). Optimal stopping problems for running minima with positive discounting rates. *Statist. Prob. Lett.* **167**, article no. 108899.
- [22] GAPEEV, P. V. (2022). Perpetual American double lookback options on drawdowns and drawups with floating strikes. *Methodology Comput. Appl. Prob.* **24**, 749–788.
- [23] GAPEEV, P. V., KORT, P. M. AND LAVRUTICH, M. N. (2021). Discounted optimal stopping problems for maxima of geometric Brownian motions with switching payoffs. *Adv. Appl. Prob.* **53**, 189–219.
- [24] GAPEEV, P. V., KORT, P. M., LAVRUTICH, M. N. AND THIJSSSEN, J. J. J. (2022). Optimal double stopping problems for maxima and minima of geometric Brownian motions. *Methodology Comput. Appl. Prob.* **24**, 789–813.
- [25] GAPEEV, P. V. AND KÜHN, C. (2005). Perpetual convertible bonds in jump-diffusion models. *Statist. Decisions* **23**, 15–31.
- [26] GAPEEV, P. V. AND LERCHE, H. R. (2011). On the structure of discounted optimal stopping problems for one-dimensional diffusions. *Stochastics* **83**, 537–554.
- [27] GAPEEV, P. V. AND LI, L. (2022). Optimal stopping problems for maxima and minima in models with asymmetric information. *Stochastics* **94**, 602–628.
- [28] GAPEEV, P. V. AND LI, L. (2022). Perpetual American defaultable standard and lookback options in models with incomplete information. *SIAM J. Financial Math.* **13**, 773–801.
- [29] GAPEEV, P. V. AND AL MOTAIRI, H. (2022). Discounted optimal stopping problems in first-passage time models with random thresholds. *J. Appl. Prob.* **59**, 714–733.
- [30] GAPEEV, P. V. AND RODOSTHENOUS, N. (2014). Optimal stopping problems in diffusion-type models with running maxima and drawdowns. *J. Appl. Prob.* **51**, 799–817.
- [31] GAPEEV, P. V. AND RODOSTHENOUS, N. (2016). On the drawdowns and drawups in diffusion-type models with running maxima and minima. *J. Math. Anal. Appl.* **434**, 413–431.
- [32] GAPEEV, P. V. AND RODOSTHENOUS, N. (2016). Perpetual American options in diffusion-type models with running maxima and drawdowns. *Stoch. Process. Appl.* **126**, 2038–2061.
- [33] GLOVER, K., HULLEY, H. AND PESKIR, G. (2013). Three-dimensional Brownian motion and the golden ratio rule. *Ann. Appl. Prob.* **23**, 895–922.
- [34] GRAVERSEN, S. E. AND PESKIR, G. (1998). Optimal stopping and maximal inequalities for geometric Brownian motion. *J. Appl. Prob.* **35**, 856–872.
- [35] GRAVERSEN, S. E. AND PESKIR, G. (1998). Optimal stopping and maximal inequalities for linear diffusions. *J. Theoret. Prob.* **11**, 259–277.
- [36] GUO, X. AND SHEPP, L. A. (2001). Some optimal stopping problems with nontrivial boundaries for pricing exotic options. *J. Appl. Prob.* **38**, 647–658.
- [37] GUO, X. AND ZERVOS, M. (2010).  $\pi$  options. *Stoch. Process. Appl.* **120**, 1033–1059.
- [38] HENRY-LABORDÈRE, P. (2009). Calibration of local stochastic volatility models to market smiles: a Monte-Carlo approach. *Risk Magazine* **22**, 10.

- [39] KALLSEN, J. AND KÜHN, C. (2004). Pricing derivatives of American and game type in incomplete markets. *Finance Stoch.* **8**, 261–284.
- [40] KARATZAS, I. AND SHREVE, S. E. (1991). *Brownian Motion and Stochastic Calculus*, 2nd edn. Springer, New York.
- [41] KIFER, Y. (2000). Game options. *Finance Stoch.* **4**, 443–463.
- [42] KITAPBAYEV, Y. (2014). On the lookback option with fixed strike. *Stochastics* **86**, 510–526.
- [43] KRYLOV, N. V. (1971). Control of Markov processes and  $W$ -spaces. *Math. USSR Izv.* **5**, 233–266.
- [44] KÜHN, C. AND KYPRIANOU, A. E. (2007). Callable puts as composite exotic options. *Math. Finance* **17**, 487–502.
- [45] KYPRIANOU, A. E. (2004). Some calculations for Israeli options. *Finance Stoch.* **8**, 73–86.
- [46] KYPRIANOU, A. E. AND OTT, C. (2014). A capped optimal stopping problem for the maximum process. *Acta Appl. Math.* **129**, 147–174.
- [47] LEPELTIER, J. P. AND MAINGUENAU, M. A. (1984). Le jeu de Dynkin en théorie générale sans l’hypothèse de Mokobodski. *Stochastics* **13**, 25–44.
- [48] LIPTSER, R. S. AND SHIRYAEV, A. N. (2001). *Statistics of Random Processes I*, 2nd edn. Springer, Berlin.
- [49] NEVEU, J. (1975). *Discrete-Parameter Martingales*. North-Holland, Amsterdam.
- [50] OTT, C. (2013). Optimal stopping problems for the maximum process with upper and lower caps. *Ann. Appl. Prob.* **23**, 2327–2356.
- [51] PEDERSEN, J. L. (2000). Discounted optimal stopping problems for the maximum process. *J. Appl. Prob.* **37**, 972–983.
- [52] PESKIR, G. (1998). Optimal stopping of the maximum process: the maximality principle. *Ann. Prob.* **26**, 1614–1640.
- [53] PESKIR, G. (2005). The Russian option: finite horizon. *Finance Stoch.* **9**, 251–267.
- [54] PESKIR, G. (2007). A change-of-variable formula with local time on surfaces. In *Séminaire de Probabilités XL*, eds C. Donati-Martin, M. Émery, A. Rouault and C. Stricker, Springer, Berlin, Heidelberg, pp. 70–96.
- [55] PESKIR, G. (2008). Optimal stopping games and Nash equilibrium. *Theory Prob. Appl.* **53**, 558–571.
- [56] PESKIR, G. (2012). A duality principle for the Legendre transform. *J. Convex Anal.* **19**, 609–630.
- [57] PESKIR, G. (2012). Optimal detection of a hidden target: the median rule. *Stoch. Process. Appl.* **122**, 2249–2263.
- [58] PESKIR, G. (2014). Quickest detection of a hidden target and extremal surfaces. *Ann. Appl. Prob.* **24**, 2340–2370.
- [59] PESKIR, G. AND SHIRYAEV, A. N. (2006). *Optimal Stopping and Free-Boundary Problems*. Birkhäuser, Basel.
- [60] QIU, S. (2020). American strangle options. *Appl. Math. Finance* **27**, 228–263.
- [61] REN, Y., MADAN, D. AND QIAN, M. Q. (2007). Calibrating and pricing with embedded local volatility models. *Risk Magazine* **20**, 138.
- [62] REVUZ, D. AND YOR, M. (1999). *Continuous Martingales and Brownian Motion*. Springer, Berlin.
- [63] RODOSTHENOUS, N. AND ZERVOS, M. (2017). Watermark options. *Finance Stoch.* **21**, 157–186.
- [64] SHEPP, L. A. AND SHIRYAEV, A. N. (1993). The Russian option: reduced regret. *Ann. Appl. Prob.* **3**, 631–640.
- [65] SHEPP, L. A. AND SHIRYAEV, A. N. (1994). A new look at the pricing of Russian options. *Theory Prob. Appl.* **39**, 103–119.
- [66] SHEPP, L. A. AND SHIRYAEV, A. N. (1996). A dual Russian option for selling short. In *Probability Theory and Mathematical Statistics: Lectures Presented at the Semester Held in St. Petersburg, Russia, March 2–April 23, 1993*, eds Ibragimov, I. A. et al. Gordon and Breach, Amsterdam, 209–218.
- [67] SHIRYAEV, A. N. (1978). *Optimal Stopping Rules*. Springer, Berlin.
- [68] SHIRYAEV, A. N. (1999). *Essentials of Stochastic Finance*. World Scientific, Singapore.
- [69] STETTNER, L. (1982). On a general zero-sum stochastic game with optimal stopping. *Prob. Math. Statist.* **3**, 103–112.
- [70] STETTNER, L. (1982). Zero-sum Markov games with stopping and impulsive strategies. *Appl. Math. Optimization* **9**, 1–24.