

**A. Bauer, S. Pawlby, D. T. Plant, D. King, C. M. Pariante,
[M. Knapp](#)**

Perinatal depression and child development: exploring the economic consequences from a South London cohort

**Article (Accepted version)
(Refereed)**

Original citation:

Bauer, A., Pawlby, S., Plant, D. T., King, D., Pariante, C. M. and Knapp, M. (2015) *Perinatal depression and child development: exploring the economic consequences from a South London cohort*. *Psychological Medicine*, 45 (1). pp. 51-61. ISSN 0033-2917

DOI: [10.1017/S0033291714001044](http://dx.doi.org/10.1017/S0033291714001044)

© 2014 [Cambridge University Press](#)

This version available at: <http://eprints.lse.ac.uk/57718/>

Available in LSE Research Online: December 2014

LSE has developed LSE Research Online so that users may access research output of the School. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LSE Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain. You may freely distribute the URL (<http://eprints.lse.ac.uk>) of the LSE Research Online website.

This document is the author's final accepted version of the journal article. There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

Perinatal depression and child development: exploring the economic consequences from a South London cohort

Annette Bauer* ¹, Susan J Pawlby³, Dominic T Plant³, Derek King¹, Carmine M Pariante³, Martin Knapp^{1 2}

¹ Personal Social Services Research Unit, London School of Economics and Political Science

² Centre for the Economics of Mental and Physical Health, Institute of Psychiatry, King's College London

³ Department of Psychological Medicine, Institute of Psychiatry, King's College London

Background. Depression in mothers during pregnancy and in the postnatal period has been recognised to have wide-ranging adverse impacts on offspring. Our study examines some of the outcomes and long-term economic implications experienced by offspring who have been exposed to perinatal depression.

Method. We analysed the effects of perinatal depression on child development outcomes of children at ages 11 and 16 years from a community-based South London Child Development Study. Economic consequences were attached to those outcomes through simple decision-analytic techniques, building on evidence from studies of epidemiology, health-related quality of life, public sector costs and employment. The economic analysis takes a life-course perspective from the viewpoints of public sector, individual and society.

Results. Additional risks that children exposed to perinatal depression develop emotional, behavioural or cognitive problems ranged from 5% to 21%. In addition, there was a high risk (24%) that children would have special educational needs. We present results in the form of cost consequences attached to adverse child outcomes. For each child exposed to perinatal depression, public sector costs exceeded £3,030, costs due to reduced earnings were £1,400 and health-related quality of life loss was valued at £3,760.

Conclusion. Action to prevent or treat mothers' depression during pregnancy and after birth is likely to reduce public sector costs, increase earnings and improve quality of life for children who were exposed to the condition.

Key words: child development, perinatal depression, long-term, cost consequences, economic modelling

*Address for correspondence: Annette Bauer, Personal Social Services Research Unit, London School of Economics and Political Science, Houghton Street, London WC2A 2AE, UK; email: a.bauer@lse.ac.uk

Introduction

Depression in mothers during pregnancy and the postnatal period can not only substantially affect the quality of life for mothers, but also have wide-ranging adverse impacts on their offspring. Studies evidence the negative impact of perinatal depression on children and their higher risk of developing emotional, behavioural and intellectual problems (Murray & Cooper 1997; Hay *et al.* 2001; Essex *et al.* 2001; Marchand *et al.* 2002; Grace *et al.* 2003; McMahon *et al.* 2008; Pawlby *et al.* 2009; Fihrer *et al.* 2009; Velders *et al.* 2011). Maternal depression, particularly during pregnancy and in the postnatal period, can contribute to intergenerational transmission of socio-economic disadvantage, impacting on the child's quality of life and future life prospects, including in the labour market (Johnston *et al.* 2011).

As demonstrated by a number of studies, there is a substantial economic impact associated with mental health problems starting in childhood and reaching into adulthood and possibly later life, with wide-ranging implications for society (Scott *et al.* 2001; Healey *et al.* 2004; McCrone *et al.* 2005; Romeo *et al.* 2006; Knapp *et al.* 2011). The purpose of this research was to explore some of the early adulthood economic consequences of adverse child development effects linked to maternal depression during pregnancy and after birth.

Method

In overview, our approach was first to apply logistic regression to data from a South London cohort in order to estimate the additional risk that children develop adverse outcomes linked to their earlier exposure to perinatal depression. We then looked for evidence streams that are concerned with the costs attached to those outcomes, including those that reach into adulthood. With simple decision-analytic modelling we then combined the risk data with the various sources of evidence. The aim was to model cost consequences in adulthood until the age of 50 where evidence was strong enough to allow this.

Primary data source

We analysed data from the still on-going community-based South London Child Development Study (Hay *et al.* 2001; 2003). In this study 252 women were initially recruited from two antenatal clinics between January and December 1986; a 75% random selection was assessed by psychiatric interview during pregnancy and 85% of those entered the follow-up study after the first postnatal year. Our study focused on the 120 mothers and children for whom full information was available and who have been found to be representative of the group that entered the follow-up study (Hay *et al.* 2010). Data were available to us for a range of outcomes for the child at ages 11 and 16 years. Maternal depression was assessed and diagnosed during pregnancy and after birth using the Clinical Interview Schedule (CIS; Goldberg *et al.* 1970) for an ICD-9 diagnosis. Because of the very long follow-up period, this topic-specific study offered a good platform for examining some of the long-term implications of maternal perinatal depression on children.

Statistical analysis

First, we generated the additional probabilities that children who had been exposed to perinatal depression would experience adverse child development outcomes compared to children whose mothers were not depressed during this period. For this purpose we undertook multivariate logistic regression analysis in Stata (StataCorp 2009) to test the significance of the relationship between antenatal and/or postnatal depression and child development outcomes. We considered covariates in the analysis which had previously been found by the researchers of the South London based study to influence the relationship between either antenatal or postnatal depression and child development outcomes (Hay *et al.* 2001; 2003; 2010). These included child's gender, mother's age at birth, mother's age at birth of first child, breastfeeding, maternal depression before pregnancy, relationship breakdown in the year after birth and mother's IQ. We modelled the effect of perinatal depression for child outcomes using backward elimination logistic regression. We generated predicted probabilities for the child variables with perinatal depression as the exposure factor and adjusted for the covariates. We then calculated the additional risk for each child outcome from the difference in predicted probabilities between those children exposed and not exposed to perinatal depression.

Variables

For the regression we prepared the data as follows: we created an additional binary variable for whether the mother had experienced depression at 36 weeks into pregnancy and/or at 3 months postpartum. Then, we selected variables for adverse child experiences; our choice was guided by previous findings of significant relationships in the literature and appropriateness for use in economic evaluation. Emotional and conduct problems at age 11 years were measured on the respective subscales of the teacher-rated Strengths and Difficulties Questionnaire (SDQ; Goodman 1997); we took a threshold of five to define cases for emotional problems and a threshold of three for conduct problems (Goodman; www.sdqinfo.com). Cognitive impairment, which had been measured at age 11 years, was defined as IQ score below 81 on the Wechsler Intelligence Scale for Children (WISC-III; Wechsler 1992; Petrou *et al.* 2010). We combined three variables to create one binary variable concerning the special educational needs (SEN) status of the child. The status had been reported by parents or teachers at 11 years for the child being 'currently' on the register, or having been on the register 'ever', and at age 16 years for 'on the register since 11 years'. The status 'on the register' indicated that the child had been formally assessed for their special needs. Furthermore, a classification into five stages provided information of the level of support the child was likely to have received (under national guidance): stage one described initial action and stage five comprised a formal entitlement for additional school resources (so called 'statement'). The binary variable we created summarized whether or not the child was deemed stage 1 or above on any of those variables. Because the sample size was too small to sensibly carry out statistical analysis within those subgroups, we approximated probabilities for a child being on the register for primary school, secondary school, or throughout both, and being with a statement (stage five) or

without one (stages one to four) from the whole sample distribution. Finally, we created a binary variable for whether the adolescent at age 16 left school with or without qualifications based on teacher-reported results.

Evidence assessment

We reviewed studies concerned directly or indirectly with the economic implications of adverse development outcomes during childhood and (where possible) adulthood. Our search strategies were pragmatic and we used expert advice and snowballing to identify the relevant literature. We prioritised and selected sources which reported peer-reviewed evidence from the UK of a more recent date, based on larger samples and with the appropriate study design (following the hierarchy of evidence). Data sets looked at included: surveys or studies that measured public service use by children or adults with mental health problems; employment and earnings surveys for people with mental health problems or those leaving school without qualifications; we also looked at epidemiological studies which investigated the prevalence and incidence of mental health conditions at different ages and average durations; and we searched for studies which evaluated (incremental) health utility scores associated with different development conditions in paediatric and adult populations.

Decision analytic modelling approach

In MS EXCEL 2010 we used decision-analytic modelling to link the additional risk data from the analyses of the South London study with data we identified in the evidence search.

We estimated costs to the public sector by assigning unit costs to service use data where this had not already been undertaken in the original studies. Where possible we took unit cost estimates from the 2011 version of PSSRU annual compendium, a national source commonly used for economic evaluations in this area (Curtis 2011). All public service costs were reported at 2010-11 price levels. Next, still taking a government perspective, we valued payments of social security benefits at £80 per week, which was a weighted average of incapacity benefits, income support and Job Seeker Allowance calculated from statistics on expenditure and number of claimants in those categories (DWP 2011). We did not consider loss of tax revenues.

Then, reflecting costs to the individual and society, we estimated the additional risk of reduced earnings for, and the quality of life impacts on the individual and society. We valued productivity losses at average daily wage rates for the cost of a lost work day or an average per annum wage rate for people moving from incapacity benefit into employment (Perkins *et al.* 2009). We valued quality of life impairments for children who experienced adverse outcomes in childhood and adulthood by assigning the willingness-to-pay threshold of £20,000 for an additional year spent in best health (measured on a standardised scale). Values between £20,000 to £30,000 are used by the National Institute for Health and Care Excellence to inform decisions about the cost-effectiveness of interventions (NICE 2008). We estimated the health-related quality of life (HRQOL) loss experienced by children with adverse conditions during childhood from study findings by Petrou *et al.* (2010) who evaluated

incremental utility scores between children with and without each condition. Where a distinction was made in this source between health utility scores measured on two different scales (HUI2 and HUI3), and for different severities of the condition, we calculated a weighted (incremental) health utility score. We incorporated health-related quality of life (HRQOL) loss in adulthood where evidence was available. All costs (public service use including payments for benefit claims, productivity losses and HRQOL impairments) were discounted back to the time of birth at a rate of 3.5% (NICE 2004). The sum of the discounted costs (i.e. present value) was then multiplied by the additional risk that a child exposed to perinatal depression would experience adverse outcomes during childhood and adulthood.

Whilst this described our general approach, we carried out the following additional steps specific to each area of child development; these were determined by the ways outcomes had been measured and the available evidence in this field.

Modelling emotional problems

Based on findings from an earlier study in this field by Halligan *et al.* (2007) that child emotional disorder was rooted in childhood anxiety rather than depression, we prioritised information specific to anxiety disorder where this was possible. We estimated costs per year from data from a British cohort on the use of mental health specialist services, their frequency and duration for children with emotional problems (Meltzer *et al.* 2003; Snell *et al.* 2013).

We calculated treatment costs from the age of 11 to 16 years assuming the additional risk to be constant over this period and applying those cost per year estimates. We approximated treatment costs and the value of HRQOL impairments in adulthood based on assumptions drawn from the literature: we assumed an average onset of anxiety in adolescence around the age of 16 and an average persistence of 16 years (Grant *et al.* 2005). In addition, we assumed that the average probability that a child at age 16 years would continue having these mental health problems in adulthood would be 86%, a figure used in previous economic studies and which had been derived from data from a UK birth cohort study which measured depression and anxiety of individuals from age 13 to 56 (Colman *et al.* 2007, Friedli & Parsonage 2009). Because of the considerable overlap between the two conditions in adulthood and a treatment response that in practice was expected to be similar, we estimated treatment costs for adults based on annual cost for depression (Thomas & Morris 2003).

We derived estimates for HRQOL loss in adulthood from a European study which evaluated incremental health utilities for mild, moderate and severe forms of generalised anxiety disorder in adults compared to those without the condition via the Short-Form-12 (Toghanian *et al.* 2010).

Productivity losses and costs to the public sector as a result of additional benefit claims were estimated from the additional probabilities of unemployment for children with psychological problems until the age of 50 years as provided by a study of the British National Child Development cohort (Goodman *et al.* 2011).

Modelling behavioural problems

Studies of the economic consequences of conduct problems have found that costs to the public sector and society can be very high (Bonin *et al.* 2011). In our analysis we used existing cost estimates from longitudinal studies (Scott *et al.* 2001; Friedli & Parsonage 2007; 2009). Data were available to predict public service costs between ages 10 and 28. This included resource use for treating mental illness and substance misuse, and costs to the criminal justice and education systems. Cost estimates were provided for those with conduct problems; it was assumed that, of those, 10% were severe enough to be classed a disorder (Friedli & Parsonage 2007; 2009).

Adolescents with behavioural problems have been found to be at a higher risk of leaving school without qualifications (Colman *et al.* 2009). Productivity losses were taken directly from Friedli and Parsonage. HRQOL losses during childhood were estimated as described earlier. How conduct problems or disorder affect the HRQOL in adults is still uncertain, and so no cost could be calculated (see for example, McCabe *et al.* 2005).

Modelling intellectual problems

We estimated public sector costs for additional school expenditure associated with an increased risk that a child would be deemed stage 1, captured through our newly created variable. Because the South London study sample was too small to identify the additional risk distinguished by school periods or stages, we derived parameters for those from the distribution in the sample as a whole. If it had been reported at 11 years that the child had been on the register 'ever' but not 'currently', then we assumed that the child had been on the register for the primary school period only; children who were reported at age 16 years to have been on the register since age 11 were assumed to have been there for the whole secondary school period; where responses were positive to all three questions it was assumed that the child had been on the register for their whole time during primary and secondary school. We excluded observations for which responses were contradictory, for example when it was reported that the child at 11 years was 'currently' on the register but at age 16 years not reported as having been on register since 11 years. In addition, from the sample, we derived probabilities that a child would be on the register without a statement (stages 1 to 4) or with a statement (stage 5) during primary and secondary school.

Costs for school support provided to the child were applied to these probabilities. Because the only cost estimates which provided a split into costs for children with and without a statement for SEN was an older source (Boyle & Burton, 2004) we derived an alternative estimate from recent school budget data provided by the Department for Education. This second approach, however, did not allow a distinction to be made between costs with or without statement, and instead we took an average figure per pupil with SEN.

We used the additional risk of a child leaving school without qualifications to predict productivity losses and additional government expenditure for paying benefit claims. First, we calculated per annum values linked to the risk of lower earnings for someone who left school without qualifications compared to someone who left with

qualifications; we then estimated annual costs linked to the additional risk for someone leaving school without qualifications to be unemployed – one, in the form of productivity losses and, two, in the form of social security payments. Probability values were derived from employment rates published in labour market reports (Pigott 2011); we used earning statistics for average salary information for people with different qualifications. Employment-related costs were valued for the period when a person leaves school at 16 until retirement age (conservatively assumed to start early when people reach their fifties). HRQOL losses due to cognitive impairment in childhood were modelled for the period 11 to 16 years, for which we calculated a weighted mean of moderate and severe conditions based on an assumption that 10% of problems were classed a disorder.

Results

The average age of mothers at the birth of their index child was 26.7 (SD=5.0; range 17-44) and for almost half of the women (48%) this was their first child. At the time of birth 65% were married and another 28% were living with their partners; this proportion changed substantially during the study period with only 51% of the offspring living with both biological parents by age 11 years. The vast majority (88%) of women were from a working class background and 72% had at least basic school qualifications. Seventy-two per cent of the sample was White British. Thirty-one per cent of the women were diagnosed with depression during pregnancy and 22% with depression at 3 months postpartum. This compares with a very wide range of prevalence rates reported in the literature (from as low as 5% to more than 25%), a range explained at least in part by variations in assessment tools and methods, timing and population (Gavin *et al.* 2005). In this study the comprehensive diagnosis process that was carried out is likely to have contributed to a more accurate identification of cases; at the same time some of the population characteristics such as a higher proportion of working-class families and families from ethnic minorities could explain a relatively high prevalence. The sample, although unlikely to differ greatly from many other urban areas in England, was not representative of the country as a whole.

Additional risk for developing adverse child outcomes

Results from the statistical analysis, including probabilities for the development of adverse child outcomes depending on whether or not the mother was depressed during and/or after pregnancy, are presented in *Table 1*. Perinatal depression, measured through the combined variable, was significantly associated with child emotional problems and whether a child was registered at school because of special educational needs. The significant effect we found on behavioural problems originated in depression during pregnancy whilst the child's cognitive impairment (measured as IQ below 81) was affected by postnatal depression. Perinatal depression in combination with breastfeeding predicted school performance. Because of the important link between mother's depression and breastfeeding we decided to reconstruct the relationship by linking the effect of perinatal depression on breastfeeding from the literature to the additional risk (from our study) of 10%

that a child who had not been breastfed would leave school without qualifications. From a study by Henderson *et al.* (2003) we derived the additional probability of 2.5%. This was based on a reported overall risk for mothers to stop breastfeeding between the second and sixth months of 22% and a reported higher risk of 1.25 for depressed mothers compared to non-depressed mothers. Values for additional risks derived from differences in probabilities are shown in *Table 2*.

Costs linked to emotional problems

Parameters used for the analysis are shown in *Table 2*. The costs attributable to perinatal depression for treating emotional problems during adolescence and adulthood were £80 at the time of birth. The direct costs to the government for paying out-of-work benefits linked to a child's greater risk of unemployment were estimated at £180. Productivity losses linked to the additional risk of a person being unemployed were valued at £490. The value of HRQOL during childhood was estimated at £620 at the time of birth. HRQOL loss during adulthood was estimated at £630 so that the combined loss was valued at £1,250.

Costs linked to behavioural problems

Parameters considered in the analysis are presented in *Table 2*. The public sector costs associated with the additional risk of a child developing conduct problems (including the risk of developing a disorder) were estimated at £1,040. Costs from reduced earnings linked to perinatal depression were £890 at the time of birth and those from reduced HRQOL were £1,040.

Costs linked to intellectual problems

For a child reported by their parents or teachers to be/have been on the SEN register (as shown by the binary variable we created) we found: the probability of being on the register during both primary and secondary school without a statement (stages 1 to 4) was 35% and the probability for the same length with a statement was 17%; the probability for being registered during primary school only and without a statement was 26% whilst it was 22% for being on the register (stages 1 to 4) during secondary school only. Children who had a statement appeared to have it for their whole school career so that there were zero probabilities for having a statement during primary or secondary school only. Costs for additional school support were estimated at £3,030 if the data from Boyle & Burton (2004) were applied and at £3,320 when we applied school budget data. We included the lower value in the results. The productivity loss for someone leaving school without qualifications attributable to perinatal depression was £1,400 and respective out-of-work payments amounted to £20 per child. HRQOL loss related to cognitive impairment during childhood was estimated at a value of £3,760. The data informing these calculations are presented in *Table 2*.

Cost findings combined

Findings on cost consequences per child exposed to perinatal depression linked to the three areas of child development are shown in *Table 3*. The costs for each area of child development could not be simply aggregated because some of the consequences overlapped; for example, the calculated reduced earnings for

someone leaving school without qualifications overlapped with the productivity loss calculated from adverse emotional and behavioural outcomes. For each cost category (public sector, productivity, HRQOL) the highest value across the three child outcome areas can be interpreted as the cost minimum; that is, the costs that would occur if there was a complete overlap between the consequences. Following on from this, costs for public service spending were then at a minimum £3,030; loss of earnings exceeded £1,400 per child and quality of life impairments were valued at a minimum of £3,760.

Discussion

Summary

The aim of this research was to get a better understanding of the economic implications of perinatal depression through its effect on children. We used the South London Development study to extract outcomes data for children (exposed; not exposed) at ages 11 and 16 years, and linked those to UK literature on economic consequences. We established the additional probabilities that children exposed to antenatal and/or postnatal depression develop negative outcomes in three areas of child development (emotional, behavioural and cognitive). Only considering a subset of the economic costs of adverse child development linked to mother's depression – because of data limitations – we find that the cumulative costs (discounted back to time of birth) are high, with public sector costs per case of perinatal depression of at least £3,030 and a minimum economic value of £8,190 if non-cashable productivity and quality of life losses are included.

Strengths and limitations

Our study explored some but not all of the economic consequences that are likely to be linked to perinatal depression and its effects on children. We calculated costs as they arise for the government in the form of expenditure for public services and welfare payments, and costs to the individual in the form of quality of life losses and productivity losses. With the exception of the welfare payments, these together are costs to society (welfare payments decrease government expenditure but increase individuals' income by the same amount, so that from a societal perspective they represent a zero-sum exchange).

These findings complement other evidence on early childhood factors and longer-term economic effects that can be potentially reduced if risk factors of perinatal depression are identified and addressed through prevention or treatment. It contributes to knowledge on the costs of a mental health condition that, although defined by a relatively short time-frame, can have particularly wide-ranging economic impacts. Decision-analytic modelling allowed us to draw on a broad range of epidemiological and economic data sets and include a wide range of economic consequences over a long time horizon.

Our study only explored a subset of economic consequences. Our approach of searching the literature for information about those was thorough, but we were not able to conduct a systematic review. We relied on data which provided information on reported emotional and behavioural problems at age 11 years, but this meant we could not include treatment costs for earlier years, and these have shown to be substantial for a small number of children with symptoms at the severe end of, for example, the antisocial behaviour spectrum (Romeo *et al.* 2006). We did not include internalising and externalising outcome measures in our analysis because they are broader constructs that are often measured at a time when symptoms are not distinctively manifested.

A range of public sector costs and costs to individuals could not be included in the analysis because of the absence of evidence. For example, evidence from the US indicates that by far the largest proportion (85%) of the cost of anxiety disorders is attributable to an excess use of health services (Kessler & Greenberg, 2002). Furthermore, a current lack of knowledge about disorder persistence, progression and regression of some mental illnesses into old age prevented us from projecting the economic consequences into old age. The analysis also did not include the knock-on effects and costs that accrue to other individuals (such as family members or the victims of crime).

A further limitation is that our estimates of additional probabilities were derived from a relatively small sample, with consequences for statistical significance.

Implications

Numerous studies have emphasised the need to focus on supporting mothers during the sensitive period around childbirth. Our work indicates that there could be not only health and wellbeing arguments but also economic arguments for action to prevent or treat perinatal depression. Earlier work which estimated the burden of postnatal depression on mothers themselves also demonstrated the benefits for investing in this area (Morrell *et al.* 2009; Brugha *et al.* 2011; Dagher *et al.* 2011; Bauer *et al.* 2011). Although in this paper we do not evaluate interventions for mothers, our work is, as far as we are aware, the first to explore some of the economic consequences from a child's perspective. It provides further argument for policy makers and service providers to focus on planning an effective support infrastructure for new mothers and mothers-to-be.

Our findings only relate to higher-risk populations and further evidence is needed to understand a fuller set of economic consequences which relate to all groups of childbearing women. We know from previous research that the influence of other risk factors needs to be considered, such as whether or not children are breastfed and the nature of the mother-infant relationship, including child maltreatment or parent hostility (Plant *et al.* 2013; Pawlby *et al.* 2011; Barker *et al.* 2011; Velders *et al.* 2011). There is an important role for studies which evaluate the (cost-)effectiveness of interventions that aim to reduce those predictors of adverse child outcomes such

as perinatal depression, and for measures of long-term effects (for example meta-analysis by Cuijpers *et al.* 2008).

Acknowledgements. We are grateful to a number of people for comments and help over a long period, including Cheryll Adams, Jane Morrell, Madeleine Stevens, Eva Bonin, Nika Fuchkan.

Declaration of interest. None

References

- Barker ED, Jaffee SR, Uher R, Maughan B** (2011). The contribution of prenatal and postnatal maternal anxiety and depression to child maladjustment. *Depression and Anxiety* **28**, 696-702.
- Bauer A, Knapp M, McDaid D** (2011). Health visiting and reducing post-natal depression. In *Mental health promotion and mental illness prevention: The economic case* (ed. Knapp M, McDaid D, Parsonage M), pp. 4-5. Department of Health, London.
- Bonin EM, Stevens M, Beecham J, Byford S, Parsonage M** (2011). Costs and longer-term savings of parenting programmes for the prevention of persistent conduct disorder: a modelling study. *BMC Public Health* **11**, 803.
- Boyle D, Burton E** (2004). *Making sense of SEN – Special educational needs, A guide for donors and grant-makers*. New Philanthropy Capital, London.
- Brugha T, Morrell CJ, Slade P, Walters SJ** (2011). Universal prevention of depression postnatally: cluster randomized trial evidence in primary care. *Psychological Medicine* **41**, 739-748
- Colman I, Ploubidis G, Wadsworth M, Jones P and Croudace T (2007), A longitudinal typology of symptoms of depression and anxiety over the life course. *Biological Psychiatry* **62**: 1265-1271.
- Cuijpers P, Brännmärk JG, Straten van A (2008), Psychological treatment of postpartum depression: A meta-analysis. *Journal of Clinical Psychology* **64**: 103-118.
- Curtis L** (2011). *Unit Costs of Health and Social Care 2011*. Personal Social Services Research Unit, University of Kent.
- Dagher RK, McGovern PM, Dowd BE, Lundberg U** (2011). Postpartum depressive symptoms and the combined load of paid and unpaid work: a longitudinal analysis. *International Archives of Occupational and Environmental Health* **84**, 735-43.
- Danese A, Moffitt TE, Harrington H, Milne BJ, Polanczyk G, Pariante CM, Poulton R, Caspi A** (2009). Adverse Childhood Experiences and Adult Risk Factors for Age-Related Disease. *Archives of Paediatrics* **163**, 1135-1143.
- DWP** (2011). *Statistical and Accounting Data, Benefit Expenditure by Country, Region and Local Authority, 2009/10*, Department of Work and Pension, (http://research.dwp.gov.uk/asd/asd4/la_expenditure.xls). Last accessed 15 April 2011.
- Essex M, Klein M, Mich R, Smider N** (2001). Timing of initial exposure to maternal major depression and children's mental health symptoms in kindergarten. *British Journal of Psychiatry* **179**, 151-56.

Fergusson DM, Horwood LJ, Ridder EM (2005). Show me the child at seven: the consequences of conduct problems in childhood for psychosocial functioning in adulthood. *Journal of Child Psychology and Psychiatry and Allied Disciplines* **46**, 837-49.

Fihrer I, McMahon CA, Taylor AJ (2009). The impact of postnatal and concurrent maternal depression on child behaviour during the early school years. *Journal of Affective Disorders* **119**, 116-23.

Friedli L, Parsonage M (2007). *Mental Health Promotion: Building an Economic Case*. Northern Ireland Association for Mental Health, Belfast.

Friedli L, Parsonage M (2009). *Promoting mental health and preventing mental illness: the economic case for investment in Wales*. Mental Health Promotion Network, Wales.

Gavin NI, Gaynes BI, Lohr KN, Meltzer-Brody S, Gartlehner G, Swinson T (2005). Perinatal depression: a systematic review of prevalence and incidence, *Journal of Obstetrics and Gynecology*, **106**, 1071-83.

Goldberg DP, Cooper B, Eastwood MR, Kedward HB, Shepherd M (1970). A standardized psychiatric interview for use in community surveys. *British Journal of Preventive Social Medicine* **24**, 18-23.

Goodman R *Interpreting Symptom Scores and Defining "Caseness" from Symptom Scores*. (www.sdqinfo.com).

Goodman R (1997). The Strengths and Difficulties Questionnaire: a research note. *Journal of Child Psychology and Psychiatry*, **38**, 581 -586.

Goodman A, Joyce R, Smith, JP (2011). The long shadow cast by childhood physical and mental problems on adult life. *Proceedings of the National Academy of Sciences of the United States of America* **108**, 6032-37.

Grace SL, Evindar A, Stewart DE (2003). The effect of postpartum depression on child cognitive development and behaviour: A review and critical analysis of the literature. *Archives of Women's Mental Health* **6**, 263–74.

Grant BF, Hasin DS, Blanco C, Stinson FS, Chou SP, Goldstein RB, Dawson DA, Smith S, Saha TD, Huang B (2005). The Epidemiology of Social Anxiety Disorder in the United States: results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Journal of Clinical Psychiatry* **66**, 1351-61.

Green H, McGinnity A, Meltzer H, Ford T, Goodman R (2005). Mental health of children and young people in Great Britain 2004. *Applied Physics Letters* 1-383.

- Hay DF, Pawlby S, Sharp D, Asten P, Mills A, Kumar R** (2001). Intellectual problems shown by 11-year-old children whose mothers had postnatal depression. *Journal of Child Psychology and Psychiatry and Allied Disciplines* **42**, 871-89.
- Hay DF, Pawlby S, Angold A, Harold GT, Sharp D** (2003). Pathways to violence in the children of mothers who were depressed postpartum. *Developmental Psychology* **39**, 1083-94.
- Hay DF, Pawlby S, Waters CS, Perra O, Sharp D** (2010). Mothers' antenatal depression and their children's antisocial outcomes. *Child Development* **81**, 149-165
- Halligan SL, Murray L, Martins C, Cooper PJ** (2007). Maternal depression and psychiatric outcomes in adolescent offspring: A 13-year longitudinal study. *Journal of Affective Disorders* **97**, 145-54.
- Healey A, Knapp M, Farrington DP** (2004). Adult labour market implications of antisocial behaviour in childhood and adolescence: findings from a UK longitudinal study, *Applied Economics* **36**, 93-105.
- Henderson JJ, Evans SF, Straton JAY, Priest SR, Hagan R** (2003). Impact of Postnatal Depression on Breastfeeding Duration, *Birth* **30**, 175-80.
- Johnston DW, Schurer S, Shields MA** (2011). *Evidence on the Long Shadow of Poor Mental Health across Three Generations*. Discussion Paper Series IZA DP No. 6014, Institute for the Study of Labor.
- Kessler RC, Greenberg PE** (2002). The economic burden of anxiety and stress disorders. In *Neuropsychopharmacology: The Fifth Generation of Progress* (ed. K. L. Davis, D. Charney, J. T. Coyle, C. Nemeroff), American College of Neuropsychopharmacology.
- Knapp M, King D, Healey A, Thomas C** (2011). Economic outcomes in adulthood and their associations with antisocial conduct, attention deficit and anxiety problems in childhood. *Journal of Mental Health Policy and Economics* **14**, 137-47.
- Manning WG, Wells KB** (1992). Effects of psychological distress and psychological well-being on use of medical services. *Medical Care* **30**, 541-53.
- Marchand JF, Hock E, Widaman K** (2002). Mutual relations between mothers' depressive symptoms and hostile-controlling behaviour and young children's externalizing and internalizing behavior problems. *Parenting: Science and Practice*, **2**: 335-53.
- McCabe C, Sutcliffe P, Kaltenthaler E** (2005). *Parent-Training Programmes in the Management of Conduct Disorder: A report from the NICE Decision Support Unit and the SchARR Technology Assessment Group*, July 2005.

- McCrone P, Knapp M, Fombonne E** (2005). The Maudsley long-term follow-up of child and adolescent depression. Predicting costs in adulthood. *European Child Adolescent Psychiatry* **14**, 407-13.
- McMahon C, Trapolini T, Barnett B** (2008). Maternal state of mind regarding attachment predicts persistence of postnatal depression in the preschool years. *Journal of Affective Disorders* **107**, 199–203.
- Meltzer H, Gatward R, Corbin T, Goodman R, Ford T** (2003). *Persistence, onset, risk factors and outcomes of childhood mental disorders*. London: TSO.
- Morrell CJ, Warner R, Slade P, Dixon S, Walters S, Paley G, Brugha T** (2009). Psychological interventions for postnatal depression: cluster randomised trial and economic evaluation: the PoNDER trial. *Health Technology Assessment* **13**, 1-153.
- Murray L, Cooper P** (1997). Effects of postnatal depression on infant development. *Archives of Disease in Childhood* **77**, 99-101
- NICE** (2004). Guide to the methods of technology appraisal. London: National Institute for Health and Clinical Excellence, 2004.
- NICE** (2008). *Guide to the Methods of Technology Appraisal*. National Institute for Health and Clinical Excellence, London.
- ONS** (2011). *Earnings by qualification, 2011*; Office for National Statistics, last accessed 12th April 2013; http://www.ons.gov.uk/ons/dcp171776_229888.pdf
- ONS** (2012). *Live Births, Stillbirths and Infant deaths: Babies born in 2009 in England and Wales*. Statistical Bulletin; Office for National Statistics, London.
- Pawlby S, Hay DF, Sharp D, Waters CS, O’Keane V** (2009). Antenatal depression predicts depression in adolescent offspring: prospective longitudinal community-based study. *Journal of Affective Disorders* 2009 **113**, 236-43.
- Pawlby S, Hay DF, Sharpe D, Waters CS, Pariante, CM** (2011). Antenatal depression and offspring psychopathology: the influence of childhood maltreatment. *British Journal of Psychiatry* **199**, 106-12.
- Perkins R, Farmer P, Litchfield P** (2009). *Realising ambitions: Better employment support for people with a mental health condition, A Review to Government*. Department of Work and Pensions, London.
- Petrou S, Johnson S, Wolke D, Hollis C, Kochhar P, Marlow N** (2010). Economic costs and preference-based health-related quality of life outcomes associated with childhood psychiatric disorders. *The British Journal of Psychiatry* **197**, 395-404.

Pigott G (2011). *Focus on London 2011: Labour Market: Beyond Headlines*. Greater London Authority, London.

Plant DT, Barker ED, Waters CS, Pawlby S, Pariante CM (2013). Intergenerational transmission of maltreatment and psychopathology: the role of antenatal depression. *Psychological Medicine* **43**, 519-528.

Romeo R, Knapp M, Scott S (2006). Economic cost of severe antisocial behaviour in children - and who pays it. *British Journal of Psychiatry* **188**, 547.

Scott S, Knapp M, Henderson J, Maughan B (2001). Financial cost of social exclusion: follow up study of antisocial children into adulthood. *British Medical Journal* **323**, 1-5.

Snell T, Knapp M, Healey A, Guglani S, Evans-Lacko S, Fernandez JL, Meltzer H, Ford T (2013). Economic impact of childhood psychiatric disorder on public sector services in Britain: estimates from national survey data. *Journal of Child Psychology and Psychiatry*. doi: 10.1111/jcpp.12055.

StataCorp (2009). *Stata Statistical Software Release 10.1*. StataCorp, College Station, Texas.

Thomas CM, Morris S (2003). Cost of depression among adults in England in 2000. *British Journal of Psychiatry* **183**, 514-19.

Toghanian S, DiBonaventura M, Jarbrink K, Locklear J (2010). *The economic and humanistic burden of generalised anxiety disorder: a retrospective database analysis in Europe*. Presented at the 13th Annual European Congress of the International Society for Pharmacoeconomics and Outcomes Research, Prague.

Velders FP, Dieleman G, Henrichs J, Jaddoe VW, Hofman A, Verhulst FC, Hudziak JJ, Tiemeier H (2011). Prenatal and postnatal psychological symptoms of parents and family functioning: the impact on child emotional and behavioural problems. *European Child & Adolescent Psychiatry* **20**, 341-350.

Wechsler D (1992). *The Wechsler Intelligence Scale for Children (WISC-III UK)*. London: The Psychological Corporation.

Table 1. Results from statistical analysis using backward elimination logistic regression: p-value for retaining effect=0.20, p-value for addition to the model=0.10; perinatal depression effect always retained. Values in table are odds-ratio (95% confidence interval)

	SDQ emotional problems score above 5 (n=129)	SDQ conduct problems score above 3 (n=129)	IQ below 81 (n=139)	Special education needs (n=133)
Pre- and/or post-natal depression: yes (relative to no)	6.67 * (1.13, 39.17)	3.53 * (1.27, 9.83)	2.93 * (1.03, 8.34)	2.82 ** (1.29, 6.18)
Child's sex: Male (relative to female)	8.58 (0.94, 78.22)			
Child breastfed: Yes (relative to no)	0.24 (0.04, 1.40)	2.48 (0.86, 7.11)	0.49 (0.18, 1.34)	0.33 ** (0.15, 0.71)
Mother's age at birth of index child		0.46 (0.23, 0.93)		
Mother's depression before pregnancy: yes (relative to no)				
Relationship changes 1 partner change bet. birth and 1yr 2 partner changes bet. birth and 1yrrelative to no partner changes				
Mother's IQ			0.93 (0.010)**	
Constant	0.01 ** (0.001, 0.11)	0.21 (0.05, 0.80)	29.02 (0.144)	0.65 (0.35, 1.19)
Model chi-squared test p-value	0.0057	0.0060	0.0004	0.0005
Pseudo R squared	0.2334	0.1108	0.1674	0.0888
Probability of positive outcome if pre- and/or post-natal depression - No	0.012	0.09	0.1	0.25
Probability of positive outcome if pre- and/or post-natal depression - Yes	0.07	0.21	0.31	0.49

* p<0.05; ** p<0.01

Table 2. Parameters for estimating costs of emotional problems in child- and adulthood; MHS = mental health specialist; HUI = Health Utility Index

Parameter	Mean	Source and description
Additional risk of adverse outcomes for children exposed to perinatal depression		
Δ probability child (11yrs) develops emotional problems	5 %	Primary data
Δ probability child (11yrs) develops conduct problems	12 %	
Δ probability child (11yrs) has cognitive impairment (IQ below 81)	21 %	
Δ probability child reported by parent or teacher to be on special educational needs register	24 %	
Δ probability child to leave secondary school without qualification	0.25 %	Primary data, Henderson <i>et al.</i> 2003
Incremental health utility scores in childhood from Petrou <i>et al.</i> 2010		
Emotional problems	0.164	HUI Mark 3: 0.199; HUI Mark 2 0.128
Conduct problems	0.115	HUI Mark 3 (0.143) and HUI Mark 2 (0.086)
Cognitive impairment	0.237	Incremental utility score for moderate symptoms (assumed 90% prevalence) HUI Mark 3 (0.273) and HUI Mark 2 (0.158), midpoint 0.216 Incremental utility score for severe symptoms (assumed 10% prevalence) HUI Mark 3 (0.571) and HUI Mark 2 (0.286), midpoint 0.429
Parameters for estimating costs of emotional problems in child- and adulthood		
Probability that child sees MHS	20 %	Meltzer <i>et al</i> 2003
Mean number of appointments		
- Clinical psychologist	4.7	
- Psychiatrist	4.8	
- Counsellor	4.5	
- Mental health nurse	9.1	
Mean duration per appointment (in hours)		
- Clinical psychologist	0.9	
- Psychiatrist	0.75	
- Counsellor	0.83	
- Mental health nurse	0.78	
Unit costs per hour of face-to-face contact (in £)		PSSRU Unit costs for health and social care 2011
- Clinical psychologist	£135	
- Psychiatrist	£353	
- Counsellor	£66	
- Mental health nurse	£65	
Treatment cost for depression per year during adulthood	£200	Thomas & Morris 2003, in 2010/11 prices
Additional annual probability of unemployment (from 23yrs to 50yrs)	11 %	Goodman <i>et al</i> 2011; assumed constant based on additional probabilities of being unemployed at different ages: 0.116(23years), 0.0103 (33years),0.107 (42years)0.107 (50yrs)
Incremental health utility of anxiety in adulthood	0.0989	Derived from Toghianian <i>et al</i> 2010; health utilities for mild (0.64), moderate (0.6)

		and severe (0.54) anxiety multiplied with probabilities of mild (0.442) moderate (0.448) and severe (0.11) anxiety; average health utility in people without anxiety was 0.71
Parameters for estimating costs of conduct problems in child- and adulthood		
Public service costs linked to conduct <i>problems</i> from age 10 to 28yrs	£17,310	Derived from Scott <i>et al</i> 2001, in 2010/11 prices and discounted to time at birth
Public service costs linked to conduct <i>disorder</i> from age 10 to 28yrs	£64,920	Derived from Scott <i>et al</i> 2001, in 2010/11 prices and discounted to time at birth
Value for reduced life-time earnings linked to conduct <i>problems</i>	£6,930	Derived from Friedli & Parsonage 2009: £75,000 out of which 9% are reduced earnings; in 2010/11 prices
Value for reduced life-time earnings linked to conduct <i>disorder</i>	£10,770	Derived from Friedli & Parsonage 2009: £150,000 out of which 7% are reduced earnings; in 2010/11 prices
Parameters for estimating costs of intellectual problems in child- and adulthood		
Additional annual spend per pupil with SEN without statement in primary (secondary) school	£1,130 (£1,270)	Boyle & Burton (2004), based on data from the Audit Commission and DfES; in 2010/11 prices
Additional annual spend per pupil with SEN with statement in primary (secondary) school	£6,770 (£4,510)	Boyle & Burton (2004), based on data from the Audit Commission and DfES; in 2010/11 prices
Additional annual spend per pupil with SEN in mainstream school (SEN budget for mainstream schools of £3.8bn and number of pupils with SEN in mainstream schools of 1.5m for England 2010/11)	£2,400	Department for Education, Statistical First Release 19/2010; School budget data reports 2010/11
Δ earnings per year for someone with compared to someone without qualifications	£3,420	ONS 2011; assumed are 1,800 working hours per year
Probability for someone without school qualifications to be employed	44 %	Data Management and Analysis Group, Social Exclusion Update 07/08
Δ probability for someone to be unemployed with/ without school qualifications	30 %	Pigott 2011

Table 3. Costs per child exposed to perinatal depression, at 2010/11 prices, discounted to time of birth

	Public sector costs (in £)*	Productivity loss (in £)	Quality of life impairments (in £)
Costs attached to additional risk that child develops emotional problems	260	490	1,250
Costs attached to additional risk that child develops behaviour problems	2,660	890	1,040
Costs attached to additional risk that child develops cognitive problems <ul style="list-style-type: none"> - Special educational needs - No school qualifications - IQ <81 	3,030	1,400	3,760

* Includes government payments for benefit claims

