

Martin Orrell, Elisa Aguirre, Aimee Spector, Zoë Hoare, Robert T. Woods, Amy Streater, Helen Donovan, Juanita Hoe, [Martin Knapp](#), Christopher Whitaker and Ian Russell
Maintenance cognitive stimulation therapy for dementia: single-blind, multicentre, pragmatic randomised controlled trial

**Article (Accepted version)
(Refereed)**

Original citation:

Orrell, Martin, Aguirre, Elisa, Spector, Aimee, Hoare, Zoë, Woods, Robert T., Streater, Amy, Donovan, Helen, Hoe, Juanita, Knapp, Martin, Whitaker, Christopher and Russell, Ian (2014) *Maintenance cognitive stimulation therapy for dementia: single-blind, multicentre, pragmatic randomised controlled trial*. [The British Journal of Psychiatry](#), 204 (6). pp. 454-461. ISSN 0007-1250 DOI: [10.1192/bjp.bp.113.137414](https://doi.org/10.1192/bjp.bp.113.137414)

© 2014 [The Royal College of Psychiatrists](#)

This version available at: <http://eprints.lse.ac.uk/60469/>
Available in LSE Research Online: December 2014

LSE has developed LSE Research Online so that users may access research output of the School. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LSE Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain. You may freely distribute the URL (<http://eprints.lse.ac.uk>) of the LSE Research Online website.

This document is the author's final accepted version of the journal article. There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

Maintenance Cognitive Stimulation Therapy for dementia: a single-blind, multi-centre, pragmatic randomised controlled trial.

Martin Orrell^{1,2}, Elisa Aguirre^{1,2}, Aimee Spector^{2,3}, Zoe Hoare⁴, Robert T Woods⁵, Amy Streater^{1,2}, Helen Donovan⁶, Juanita Hoe^{1,2}, Martin Knapp⁷, Christopher Whitaker⁴, Ian Russell⁸

1 Unit of Mental Health Sciences, University College London, Charles Bell House, 67-73 Riding House Street, London, W1W 7EJ, England
Martin Orrell, Professor of Ageing and Mental Health
Elisa Aguirre, researcher assistant
Amy Streater, research assistant
Juanita Hoe, senior clinical research associate, .

2 Research and Development Department, North East London Foundation Trust, Goodmayes Hospital, Barley Lane, Ilford, Essex

3 Research Department of Clinical, Educational and Health Psychology, University College London, 1-19 Torrington Place, London, WC1E 7HB, England
Aimee Spector, Senior Lecturer in Clinical Psychology

4 North Wales Organisation for Randomised Trials in Health (NWORTH)
Institute of Medical & Social Care Research, Bangor, Wales
Zoe Hoare, senior statistician,
Christopher Whitaker, principal statistician

5 DSDC Wales, Bangor University, 45 College Road, Bangor, Gwynedd, LL57 2AS, Wales
Robert T Woods, Professor of Clinical Psychology of Older People

6 Clinical Psychology Service, South Essex Partnership NHS Foundation Trust, Gilbert Hitchcock House, 21 Kimbolton Road, Bedford, MK40 3QT, England

Helen Donovan, Clinical Psychologist

7 Personal Social Services Research Unit, London School of Economics and Political Science, Houghton Street, London WC2A 2AE

Martin Knapp, Professor of Social Policy and Director of the PSSRU

8 Swansea University, College of Medicine, Singleton Park, Swansea, SA2 8PP, UK

Ian Russell, Professor of Clinical Trials

Correspondence to: m.orrell@ucl.ac.uk

Key words dementia, cognitive stimulation, quality of life, acetylcholinesterase inhibitors (ACHEIs), cognition

ABSTRACT

Background

There is good evidence for the benefits of short-term Cognitive Stimulation Therapy for dementia but little is known about possible long-term effects.

Aims

To evaluate the effectiveness of maintenance Cognitive Stimulation Therapy for people with dementia in a single-blind, pragmatic randomised controlled trial including a sub-study with participants taking acetylcholinesterase inhibitors (ACHEIs).

Method

The participants were 236 people with dementia from nine care homes and nine community services. Prior to randomisation all participants received the seven-week 14-session CST programme. The intervention group received the weekly Maintenance CST group programme for 24 weeks. The control group received usual care. Primary outcomes were cognition and quality of life.

Results

For the intervention group at the six-month primary end point there were significant benefits for self-rated quality of life (QoL-AD $p=0.03$). At three months there were improvements for proxy-rated quality of life (QoL-AD $p=0.01$: DEMQOL $p=0.03$), and activities of daily living ($p=0.04$). The intervention sub-group taking ACHEIs showed cognitive benefits (on the MMSE) at three ($p=0.03$) and six months($p=0.03$) .

Conclusions Continuing CST improves quality of life; and improves cognition for those taking ACHEIs.

WORD COUNT: main text 3571

INTRODUCTION

There is good evidence for the benefits of cognitive stimulation for people with dementia.¹ A recent Cochrane review showed that cognitive stimulation improved both cognition and quality of life.² The review concluded the benefits of cognitive stimulation enhanced those of medication, and that it was effective whether or not acetylcholinesterase inhibitors (ACHEIs) were prescribed.² The 2011 World Alzheimer report concluded “there is strong evidence to support cognitive stimulation programmes and these interventions should therefore be routinely offered”.³ Cognitive stimulation is a psychological intervention for dementia that targets cognitive and social functioning and is designed to enhance general cognitive abilities. Cognitive Stimulation Therapy (CST) is a well-defined evidence-based version of cognitive stimulation which is standardised, including 2 training manuals and a DVD.⁴ CST was developed following review of a related approach known as Reality Orientation (RO)⁵ and evaluated CST in a pilot trial,⁶ followed by a full trial,⁴ and developed a manual⁷ and a training DVD. CST is now used widely across the UK and in several other countries. A pilot study of maintenance CST which continued for an extra 16 weekly sessions beyond the standard seven-week (14-session) CST programme⁸ found a significant improvement in cognitive function compared with CST alone. The Cochrane Review found no link between duration or frequency of the programme and degree of improvement.² Some studies have continued cognitive stimulation for six months or more,^{9,10} but there is little evidence about how far potential benefits may continue after sessions end. The Cochrane Review suggested that after the sessions finished the effects on cognition were evident for at

most three months;² and another study found no continuing effects at ten months.¹¹ This trial aimed to evaluate the effectiveness of Maintenance CST in improving cognition and quality of life in people with dementia who have completed the standard CST programme. Hence the intervention group would continue with maintenance CST whereas the control group would have standard CST only followed by treatment as usual.¹² In addition, a sub-study focused on the effects of maintenance CST on people with dementia taking ACHEIs.

METHODS

Study Design

This was a single-blind, multi-centre, pragmatic randomised controlled trial comparing (1) Maintenance CST groups after completing standard CST vs (2) standard CST only followed by treatment as usual.¹³ There was no modification in design or eligibility criteria from the study protocol¹² available at <http://www.trialsjournal.com/content/pdf/1745-6215-11-46.pdf>. The clinical trial registration number is ISRCTN26286067.

Participants

Potential centres were screened for eligibility to determine whether there were sufficient numbers of potential participants with dementia, using the inclusion criteria flow chart. Participants all met the DSM-IV criteria for dementia¹⁴ using the diagnostic algorithm and most had either Alzheimers or vascular dementia. All had mild (45%) to moderate dementia (55%) on the Clinical

Dementia Rating scale;¹⁵ could communicate, hear and see well enough to participate in the group; had no major physical illness or disability, or diagnosed learning disability. All trial participants completed seven weeks of CST₄ comprising fourteen twice-weekly 45-minute sessions according to the CST manual.⁷

Approximately half of the participants from nine care homes, and half from nine community services within London, Essex and Bedfordshire. The community centres included four voluntary sector specialist dementia day centres, and five centres based in local Community Mental Health Teams for older people. The nine care homes included; five provided by Social Services, one by the private sector, and three by a voluntary organisation. Of 21 centres approached, one refused and two had too few eligible participants. The study was approved by the Barking & Havering Local Research Ethics Committee in October 2008, ethical approval reference number: 08/H0702/68.

Intervention

After completion of the CST programme participants were randomised within each centre to either the (1) intervention group 24-week Maintenance CST (MCST) programme;¹⁶ or (2) the treatment as usual control group. Treatment as usual (TAU) varied across the 18 centres but other activities were generally available to both groups.

The Maintenance CST programme was based on the theory of cognitive stimulation as applied to the original CST programme.⁴ guided by the Medical Research Council framework for complex interventions.^{17,18} Each

Maintenance CST session has a specific theme or activity (e.g. current affairs; my life; word games) within a consistent structure including orientation-based activity, refreshments and a group song. Each group had two facilitators, one from the research team and one staff member from the participating centre (i.e. care home or community service). All facilitators had at least one year of experience in dementia care, and had attended the one-day CST training course.

Outcome measures

Participants were interviewed at baseline, before randomisation, at three months (intermediate end point) and after six months (primary end point). Researchers collected the proxy ratings of the quality of life measures, the NPI and the ADCS-ADL in structured interviews: with staff for participants in care homes; and with family carers for those in the community.

Primary outcomes

(1) Alzheimer's Disease Assessment Scale – Cognition Subscale (ADAS-Cog). This is the standard cognitive test used in clinical trials for dementia.¹⁹ This comprises 11 tasks measuring memory, language, praxis, attention and other cognitive abilities. Lower scores reflect better cognition.

(1) Quality of Life in Alzheimer's Disease scale (QoL-AD)²⁰. This is recommended by the European consensus on outcome measures for psychosocial interventions in dementia.²¹ The QoL-AD includes both self rating and proxy rating (by family care or staff) versions and covers 13

domains of quality of life. It has good internal consistency, validity and reliability. Higher scores reflect better quality of life.

Secondary outcomes

(1) Mini-Mental State Examination (MMSE), a brief but widely used generic test of cognitive function with higher scores indicating better cognition.²² This is easier to complete than the ADAS-Cog, but still has good reliability and validity.

(2) Dementia Quality of Life scale (DemQoL).²³ The DemQoL covers five domains of quality of life and uses both self reporting and rating by family carer or staff member as proxy with higher scores indicating better quality of life. It has good internal consistency, inter-rater reliability and concurrent validity and can generate a measure of utility.

(3) Neuropsychiatric Inventory (NPI).²⁴ This assesses 10 behaviours that commonly occur in dementia and has good validity and reliability. Lower scores on this specific measure reflect better behaviour. Total score is the sum of frequency x severity of each behaviour.

(4) Alzheimer's Disease Co-operative Study-Activities of Daily Living (ADCS-ADL). This validated questionnaire assesses functional capacity over the range of dementia severity.²⁵ By summing competencies this measure gives high scores to more able respondents.

Sample size

Based on the Cochrane Review we estimated effect size for Maintenance CST of 0.39 on the ADAS-Cog with power of 80% when using 5% significance level and estimating attrition at 15% between baseline and six months. This required a sample size of 230 participants randomised at baseline and an estimated 195 at follow up. With an estimated 60 participants with Alzheimer's disease and taking ACHEIs, this provided sufficient numbers for the maintenance CST/ACHEIs sub-study to estimate effect size and the feasibility of a full scale trial.

Randomisation

All participants completed the initial CST programme¹³ and were then allocated at random between (1) the intervention group receiving weekly Maintenance CST for 24 weeks or (2) the control group receiving treatment as usual (TAU). The N.WORTH Clinical Trials Unit remotely randomised participants in equal proportions between groups after stratifying for: centre (community service or care home), whether ACHEI was prescribed, and previous CST group. The random allocation sequence was computer-generated and in the ratio of 1:1. N.WORTH emailed the individual allocation to the site researcher delivering the intervention and stored the allocation list under a secure password, which was not available to any study site staff. The scheduled treatment sessions, session records and participant records were saved at the site, strictly separated, and distant from the coordinating study centre. Once the trial was completed in each centre, records were transferred to the coordinating study centre and stored by the study centre administrator who was not involved in the assessment process or data analysis. This was in

order to avoid contamination. The nature of the intervention prevented us from blinding participants to their allocated group. However blind researchers conducted initial and subsequent interviews, generally in care homes or participants' own homes. The statistician conducting the data analysis was also blind to group assignment.

Statistical analysis

We used the MACRO system to manage the data. Data was entered manually and audited internally for typing errors by hand, in order to ensure a low error rate. Data was transferred to SPSS and audited externally by NWORDTH with hard copies of assessments. These audits entailed cross checking a random 10% sample of the electronic data with the paper records to ensure accurate entry. Both random and systematic data entry errors were identified and corrected. As the audits were carried out in parallel with data entry systematic errors could be corrected at an early stage. The dataset is available from the corresponding author at m.orrell@ucl.ac.uk. Participants' consent was obtained, but the data presented are anonymised and risk of identification is low. For participants with some follow-up data, we imputed individual data missing within a scale according to the validated rules for that scale; and missing total scores by multiple regression on variables including allocated group, age, gender, ethnicity, marital status, whether prescribed an ACHEI, staff or family caregiver, centre type and individual centre (using random effects). We adopted a forward stepwise model, and used baseline scores to help predict scores at three months, then both of these to predict

scores at six months, since no participant missing at three months returned at six months.

Primary analyses by treatment allocated used analysis of covariance to adjust all imputed data for baseline differences in age, gender, ethnicity, marital status, prescription of ACHEIs, proportion of family caregivers, individual centre (using random effects) and baseline score on the variable under analysis. We then estimated the effect of treatment from the resulting model. The maintenance CST/ ACHEIs trial platform followed the same methodology as for the primary analysis and used the interaction term between ACHEIs and the treatment group to identify any effect between the two factors for the outcome measures.

Results

The recruitment period took place between January 2009 and September 2010. The final 24-week follow up was completed in May 2011. Of 272 people with dementia that started the CST groups and were considered for the trial, 36 were withdrawn (Table 1). We followed up 218 participants (92% of 236; 96% of those still alive) at 3 months and 199 (84% of 236; 89% of those still alive) at 6 months. The CONSORT flowchart (Figure 1) records the reasons for subsequent withdrawals. The withdrawal rate was similar in both arms of the trial. Of the 236 participants, 123 were allocated to the Maintenance CST group and 113 to usual care (TAU). The groups were well matched at baseline and randomisation avoided imbalances (Table 2). The mean age was 83 years and most participants were white females. On average

participants allocated to the Maintenance CST groups attended 18 of the 24 available sessions.

Outcomes

At the six-month primary end-point (Table 3), the Maintenance CST group had higher scores than the TAU group on self-rated QoL-AD (first primary outcome) which reached statistical significance with mean difference 1.78 (95% CI 0.00 to 3.60; $p=0.03$). There were no significant differences on ADAS-Cog (second primary outcome). There were no significant differences in secondary outcomes at six months. There were two types of centre studied, care homes and community services. There were significant differences among the centres over and above that explained by centre type.

At three months there were no significant differences on either primary outcome. For secondary outcomes, participants randomised to the intervention group had significantly better scores than controls on proxy ratings of quality of life (QoL-AD and DEMQOL) and daily activities. The mean difference on the proxy QoL-AD was 1.53 (95% CI 0.35 to 2.71; $p=0.01$); and for the proxy DEMQOL it was 3.24 (95% CI 0.24 to 6.24; $p=0.03$). The difference on the ADCS-ADL was 2.64 (95% CI 0.04 to 5.24; $p=0.04$).

Quality of maintenance CST programme provision

To estimate the quality of the maintenance CST provision after each session the researchers made ratings on a range of factors related to the successful running of the groups: manager's attitude (0-2), centre atmosphere (0-2), co facilitators input (0-2), group atmosphere (0-2), and average number of

sessions attended by participants (0 = less than 12, 1 = 13–20, 2 = 21-24) with higher scores indicating better quality. Centres were divided into low quality (score 0-5) and high quality (score 6-10). Eight out of 9 community centres scored as high quality compared to only 6 out of 9 care homes. The quality indicator was incorporated into the model of analysis with primary outcome results, with baseline score, centre type, age and allocation as a fixed effect and within a random effect of centre nested within the interaction of quality and type. The analysis showed that both centre type and quality of CST provision were not significant in the model using either QoL-AD or ADAS-Cog. There were differences among the centres that could not be explained by amount of sessions attended or quality of CST provision.

Maintenance CST/ ACHEI trial platform results

There were no significant results in relation to primary outcomes. Table 4 shows the observed means and SD at baseline. The means and SDs presented at follow-ups 1 and 2 are adjusted for the factors and covariates in the fitted model including the treatment group by ACHEIs interaction term. The follow-up means are standardised to a common baseline mean value. The significance levels quoted below are for the interaction term. Only for MMSE at both three and six months follow-up were significant interactions found. The results show that starting from a mean baseline MMSE of 17.8 there was the smallest decrease to 17.25 (95% CI 14.63 to 19.87, $p = 0.03$) at follow-up 2 in those taking ACHEIs and receiving maintenance CST. The largest decrease occurred in those taking ACHEIs but with no maintenance CST where the mean was 14.62 (95% CI 11.81 to 17.43, $p = 0.03$). There

were no other significant differences between groups in any other outcome measures.

Between baseline and second follow up 92% had no changes to their to ACHEI status with three participants stopping (1 TAU/2 MCST) and 11 starting (4 TAU/7MCST) medication.

There were no differences between the groups (intervention and control) in the number of reported adverse events or severity. In the intervention group there were five deaths and four withdrawals due to health issues. In the usual care group there were six deaths and five withdrawals due to health issues. All events were judged as unrelated to trial treatment or assessment contacts by the study trial coordinator and Principal Investigator.

DISCUSSION

Main findings

Cognitive stimulation for people with dementia is recognised as being effective^{2,13} and cost effective²⁶, and CST in particular improves both cognition and quality of life.^{4,13} This trial finds that after the initial CST programme, a further 24-week course of weekly Maintenance CST improves quality of life at six-month follow up but confers no additional benefit to cognition. At six months it was only participants who reported improved quality of life (a small standardised difference of 0.35), whereas at three months only the proxy respondents (carers/care staff) noted the improvement (a small standardised difference of 0.30). Participants in the intervention group also

improved in their activities of daily living at three months (a very small standardised difference of 0.15). There were no significant differences in other outcomes at either three or six months. The sub-study results suggest that people on ACHEI medication may benefit cognitively from maintenance CST suggesting a synergistic effect. This is in line with other studies combining ACHEIs and cognitive stimulation^{4,8,13}, and the Cochrane review² which found that the effect of cognitive stimulation on cognition is over and above the effects of medication alone. The relevance in terms of clinically significant change is less clear. A mean decrease of 1 point versus 4 points on the MMSE scale may make a big difference for some people with dementia. The difference might translate into economic benefits since a difference of 1 point in the MMSE score may be associated with substantial reductions in the costs of caring for people with dementia²⁷. The CST programme prior to baseline resulted in mean improvements of 4.4 points on the ADAS-Cog and 2.7 points on the MMSE¹³. Since dementia is associated with progressive cognitive decline there may have been limited potential for further cognitive improvement with the maintenance programme. This means that at six-month follow-up both groups were likely to have declined from the baseline taken after the CST groups finished, and so significant differences in cognition were only likely to be found if the usual care (CST only) group had declined more than the maintenance group.

Strengths and limitations

As participants came from nine care homes and nine community services across London, Essex and Bedfordshire, this pragmatic trial is likely to be

generalisable in many respects. However, since participants were almost all white it is hard to say how far CST is useful for other ethnic or cultural groups. Nevertheless, we have recently adapted the CST programme for a south Asian population and successfully run a local group in Hindi and Gujarati. Although we took great care to blind our researchers to allocated treatment, we could not blind those care staff and family carers who provided proxy ratings for four measures (ADCS-ADL, NPI, QoL-AD and DEMQOL) and this means there is a risk of detection bias. Notably these measures provided three of the four significant findings. Compared to the original CST study this trial had more diversity in dementia severity due to a much higher proportion recruited from the community (50% vs 15%). This resulted in the standard deviations of the cognitive measures being much higher than in the original trial of CST₄. A larger trial might find significant differences in cognition after weekly Maintenance CST. However, it may be that more frequent groups would be more efficacious. This was the first rigorous trial of Maintenance CST. The results are encouraging but not conclusive and suggest that further trials are needed and it will be important for other groups to evaluate Maintenance CST.²⁸

Future research could look in more depth at the optimum frequency and duration of CST groups, for example to continue to provide CST twice a week (rather than once weekly) for a six month period. Another option would be to repeat the standard 7 week CST programme for example after a 4 month break. However, this option could be disruptive to the groups, and would not

mirror the standard approach used in drug interventions which are given without interruption rather than as a short course.

Clinical implications

In the previous stage of this study before and after CST (prior to randomisation) we found that both cognition and quality of life significantly improved, including for those people on ACHEIs.¹³ However following Maintenance CST at six-month follow-up we found no significant differences in cognition. There were no differences on the ADAS-Cog although the MMSE showed a 0.85 points advantage for the Maintenance CST group. This does not suggest that Maintenance CST has substantial effects on cognition over and above the original benefits of the initial CST programme.² MMSE scores in mild to moderate dementia generally decrease by 2 to 4 points per year,²⁹. Before the initial CST programme¹³ (2 months before the start of this RCT) the mean ADAS-Cog was 35.0 and the mean MMSE score was 15.8. Eight months later at six-month follow-up there was no overall cognitive decline with the mean ADAS-Cog scores being 35.9 and 35.3 and the mean MMSE scores being 16.3 and 15.5 in the treatment and control groups respectively. From a standardised baseline score of 17.8 on the MMSE; in the ACHEI only group, MMSE scores fell to 14.6 points in the 6 months of the maintenance CST trial; compared with a decrease to 17.3 in the maintenance CST/ACHEI (combined) treatment; and a decrease to 16.3 in the maintenance CST only group. This suggests that CST may continue to have some degree of protective effect on cognition over and above the effects of medication. Other studies using usual care control groups have also found that a programme of

cognitive stimulation sessions over a longer time period can be effective in reducing cognitive decline in dementia.^{30,31}

In chronic conditions quality of life may be more important for older adults than disease-specific outcomes and it is a key outcome that interventions for dementia should target. Benefits to cognition alone may not be sufficient to justify an extensive programme of intervention unless they are accompanied by other benefits such as quality of life³². Two recent systematic reviews highlighted that there are few well-designed studies on the effectiveness of either pharmacological³³ or psychosocial³⁴ interventions on quality of life. Like other follow up studies we found that individual changes in quality of life were apparent for nearly three-quarters of our sample^{35,36,37}. In contrast to the Cochrane review of cognitive stimulation our study found that activities of daily living improved at three-month follow-up. However, previous research³⁵ suggests that there may be a correlation between proxy rated quality of life and activities of daily living. It might be that the effects of the intervention on proxy rated quality of life was linked with the effects on activities of daily living. At six-month follow-up these proxy rated domains showed no difference. However, for the person with dementia a temporary improvement in quality of life, cognition, or activities of daily living may all be considered worthwhile

Future research

As this was the first rigorous trial of Maintenance CST, we encourage others to implement and evaluate this novel extension in other populations in other contexts with other staff. In our research programme we have three further cognitive stimulation therapy studies²⁸ Firstly, we are undertaking a pragmatic

cluster randomised implementation trial to compare staff trained in CST receiving either (1) additional support (web support, regular phone support) or (2) no support. This will evaluate whether additional staff support results in more CST group attendances. Secondly, we are conducting an implementation in practice study measuring minimal outcomes (cognition and quality of life) for centres running CST/Maintenance CST groups. Lastly, we have developed a version of CST for use by the family carer (individual CST) and this is currently being evaluated in a large multicentre trial funded by the NIHR/HTA³⁸.

In conclusion, standard CST can improve cognition and quality of life³⁹. This trial indicates that the weekly Maintenance CST over 24 weeks provides some potential benefit beyond the basic CST programme. Further research should evaluate whether long-term CST should be provided more frequently than once a week. Over the 8 months (from the original baseline before 2 months of CST and the 6-month follow-up), the average cognitive decline in both the maintenance CST and usual care groups was considerably less than would normally be expected in practice suggesting the original CST programme had some residual beneficial effect. Maintenance CST may offer short and long term benefits to quality of life. The sub study of maintenance CST with ACHEIs provides initial evidence that maintenance Cognitive Stimulation Therapy in combination with ACHEI medication may have longer term benefits to cognition. Pharmacological and psychosocial interventions may potentially work better together than either alone.

Acknowledgements

Maintenance Cognitive Stimulation Programme (ISRCTN26286067) is part of the Support at Home - Interventions to Enhance Life in Dementia (SHIELD) project (Application No. RP-PG-0606-1083) awarded to Prof Orrell (UCL/NELFT), based in North East London Foundation Trust (study sponsor), and funded by the NIHR Programme Grants for Applied Research funding scheme. Other grant holders include Woods (Bangor), Challis (Manchester), Moniz-Cook (Hull), Russell (Swansea), Knapp (LSE) and Dr Charlesworth (UCL). This report/article presents independent research commissioned by the National Institute for Health Research (NIHR) under its Programme Grants for Applied Research scheme (RP-PG-060-1083). The views expressed in this publication are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health. The NIHR Programme Grant for Applied Research scheme monitors progress of the programme RP-PG-060-1083 through regular reports. The NIHR encourages grant-holders to seek external peer-reviewed publication.

We would also like to thank all the trial participants and their carers, especially for completing a high proportion of responses; centre managers and staff for all their collaboration and support; a wide range of healthcare practitioners for generously referring participants to the trial; and the following researchers involved in data collection, entering the data and completing trial documentation – Lauren Yates, Kier Yong, Linda Smith, Stacie McCarry, Deepak Shankar, and Caroline O’Haire.

References

- 1 Orrell M, Woods B, Spector A. Should we use cognitive stimulation therapy to improve cognitive function in people with dementia? *BMJ* 2012; 344:e633 doi: 10.1136/bmj.e633.
- 2 Woods B, Aguirre E, Spector A, Orrell M. Cognitive stimulation to improve cognitive functioning in people with dementia. *Cochrane Database of Systematic Reviews* 2012; Issue 2. Art. No.: CD005562. DOI: 10.1002/14651858.CD005562.pub2.
- 3 Prince M, Bryce R, Ferri C. *World Alzheimer Report: the benefits of early diagnosis and intervention*. London: Alzheimer's Disease International; p49. 2011. www.alz.co.uk/worldreport2011.
- 4 Spector A, Thorgrimsen L, Woods B, Royan L, Davies S, Orrell M et al. Efficacy of an evidence-based cognitive stimulation therapy programme for people with dementia: randomised controlled trial. *Br J Psychiatry* 2003; 183: 248-54.
- 5 Spector A, Thorgrimsen L, Woods B, Orrell M. Reality Orientation for Dementia: a systematic review of the evidence of effectiveness from randomised controlled trials. *The Gerontologist* 2000; 40: 206,12.
- 6 Spector A, Davies S, Woods B, Orrell M. Can reality orientation be rehabilitated? Development and piloting of an evidence-based programme of cognition-based therapies for people with dementia. *Neuropsychological Rehabilitation* 2001; 11: 377-97.
- 7 Spector A, Thorgrimsen L, Woods B, Orrell M. Making a difference: an evidence based group programme to offer cognitive stimulation therapy

- (CST) to people with dementia: manual for group leaders. London: Hawker Publications; 2006.
- 8 Orrell M, Spector A, Thorgrimsen L, Woods B. A pilot study examining the effectiveness of Maintenance Cognitive Stimulation Therapy (Maintenance CST) for people with dementia. *International Journal of Geriatric Psychiatry* 2005; **20**: 446-51.
 - 9 Buschert VC, Friese U, Teipel SJ, Schneider P, Merensky W, Rujescu D et al. Effects of a newly developed cognitive intervention in amnesic mild cognitive impairment and mild Alzheimer's disease: a pilot study. *Journal of Alzheimer's Disease* 2011; **25**: 679–94.
 - 10 Requena C, Maestu F, Campo P, Fernandez A, Ortiz T. Effects of cholinergic drugs and cognitive training on dementia: 2-year follow up. *Dementia and Geriatric Cognitive Disorders* 2006; **22**: 339-45.
 - 11 Chapman SB, Weiner MF, Rackley A, Hynan LS, Zientz J. Effects of cognitive-communication stimulation for Alzheimer's disease patients treated with donepezil. *Journal of Speech, Language, and Hearing Research* 2004; **47**: 1149-63.
 - 12 Aguirre E, Spector A, Hoe J, Knapp M, Woods B, Orrell M et al. Maintenance Cognitive Stimulation Therapy (CST) for dementia: a single-blind, multi-centre, randomised controlled trial of Maintenance CST vs. CST for dementia. *Trials* 2010; **11** (46).
 - 13 Aguirre E, Hoare Z, Streater A, Spector A, Woods B, Orrell M et al. Cognitive Stimulation Therapy (CST) for people with dementia - who benefits most? *International Journal of Geriatric Psychiatry* 2012, doi: 10.1002/gps.3823

- 14 American Psychiatric Association. Diagnostic and statistical manual of mental health disorders, 4th ed. Washington DC: APA; 1994.
- 15 Hughes C, Berg L, Danziger W, Coben LA, Martin RL. A new clinical scale for the staging of dementia. *British Journal of Psychiatry* 1992; **140**: 566-72.
- 16 Aguirre E, Spector A, Hoe J, Streater A, Woods RT, Orrell M et al. Development of an evidence- based extended programme of maintenance cognitive stimulation therapy (CST) for people with dementia. *Non-pharmacological Therapies in Dementia Journal* 2011; **1**: 198-215.
- 17 Medical Research Council. A framework for development and evaluation of RCTs for complex interventions to improve health. London: MRC; 2000.
- 18 Craig P, Dieppe P, MacIntyre S, Mitchie S, Nazareth I, Petticrew M. Developing and evaluating complex interventions: the new Medical Research Council guidance. *BMJ* 2008; **337**: a1655.
- 19 Rosen WG, Mohs RC, Davis KL. A new rating scale for Alzheimer's disease. *Am J Psychiatry* 1984; **141**: 1356-64.
- 20 Logsdon RG, Gibbons LE, McCurry SM, Teri L. Assessing quality of life in older adults with cognitive impairment. *Psychosom Med* 2002; **64**: 510-9.
- 21 Moniz-Cook E, Vernooij-Dassen M, Woods R, Verhey F, Chattat R, Orrell M et al. A European consensus on outcome measures for psychosocial intervention research in dementia care. *Aging & Mental Health* 2008; **12**: 14-25.

- 22 Folstein MF, Folstein SE, McHugh PR. Mini Mental State. A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 1975; **12**: 189–98.
- 23 Smith SC, Lamping DL, Banerjee S, Harwood R, Foley B, Knapp M et al. Measurement of health-related quality of life for people with dementia: development of a new instrument (DEMQOL) and an evaluation of current methodology. *Health Technol Assess* 2005; **9** (10).
- 24 Cummings L, Mega M, Gray K, Rosenberg-Thompson S, Carusi DA, Gornbein J. The Neuropsychiatric Inventory: comprehensive assessment of psychopathology in dementia. *Neurology* 1994; **44**: 2308-14.
- 25 Galasko D, Bennet D, Sano M, Ernesto C, Thomas R, Ferris S et al. An inventory to assess activities of daily living for clinical trials in Alzheimer's disease: the Alzheimer Disease Cooperative Study. *Alzheimer Dis Assoc Disord* 1997; **11**: S33-39.
- 26 Knapp M, Thorgrimsen L, Patel A, Spector A, Hallam, A, Woods B, Orrell M (2006) Cognitive Stimulation Therapy for dementia: is it cost effective? *British Journal of Psychiatry*, **188**, 574-580.
- 27 Jonsson L, Lindgren P, Wimo A, et al. Costs of Mini Mental Examination-related cognitive impairment. *Pharmaeconomics* 1999, **16**: 409- 416.
- 28 Streater A, Spector A, Aguirre E, Hoe J, Hoare Z, Orrell M et al. Maintenance Cognitive Stimulation Therapy (CST) in practice: study protocol for a randomized controlled trial. *Trials*, **13** (91).

- 29 Clark CM, Sheppard L, Fillenbaum GG, Galasko D, Morris JC, Heyman A. Variability in annual Mini-Mental State Examination score in patients with probable Alzheimer Disease. *Arch Neurol* 1999; **56**: 857-62.
- 30 Zanetti O, Frisoni GB, De Leo D, Buono MD, Bianchetti A, Trabucchi M. Reality Orientation Therapy in Alzheimer's disease: useful or not? A controlled study. *Alzheimer's Disease and Associated Disorders* 1995; **9**: 132-8.
- 31 Metitieri T, Zanetti O, Geroldi C, Frisoni GB, De Leo D, Trabucchi M. Reality Orientation Therapy to delay outcomes of progression in patients with dementia: a retrospective study. *Clinical Rehabilitation* 2001, **15**: 471-8.
- 32 Woods B, Thorgrimsen L, Spector A, Royan L, Orrell M. Improved quality of life and cognitive stimulation therapy in dementia. *Aging and Mental Health* 2006; **10**: 219-26.
- 33 Cooper C, Mukadam N, Katona C, Lyketsos CG, Blazer D, Livingston G et al. Systematic review of the effectiveness of pharmacologic interventions to improve quality of life and well-being in people with dementia. *Am J Geriatr Psychiatry* 2012; PMID: 22245881.
- 34 Cooper C, Mukadam M, Katona C, Lyketsos C, Ames D, Livingston G et al. Systematic review of the effectiveness of non-pharmacological interventions to improve quality of life of people with dementia. *International Psychogeriatrics* 2012, doi:10.1017/S1041610211002614
- 35 Selwood A, Thorgrimsen L, Orrell M. Quality of life in dementia—a one-year follow-up study. *Int J Geriatr Psychiatry* 2005; **20**: 232–7.

- 36 Lyketsos CG, Gonzales-Salvador T, Chin J, Baker A, Black B, Rabins P. A follow-up study of change in quality of life among persons with dementia residing in a long-term care facility. *Int J Geriat Psychiatry* 2003; **18**: 275–81.
- 37 Missotten P, Ylief M, Di Notte D, Paquay, De Lepeleire J, Fontaine O. Quality of life in dementia: a 2-year follow-up study. *Int J Geriat Psychiatry* 2007; **22**: 1201–7.
- 38 Orrell M, Yates L, Burns A, Russell I, Woods RT, Hoare Z et al. Individual Cognitive Stimulation Therapy for Dementia (iCST Trial), *Trials*, 2012, **13**:172 doi:10.1186/1745-6215-13-172.
- 39 National Institute for Health and Clinical Excellence and the Social Care Institute for Excellence (NICE-SCIE). Dementia: supporting people with dementia and their carers in health and social care. Clinical Guideline 42. London: NICE and SCIE; 2007.

Figure 1. Consort flowchart of participants' progress




Table 1. Baseline characteristics of 236 participants in maintenance CST trial

Participants at Baseline 0: Before start of CST groups	272
Total lost from the beginning of CST groups	36
Reason for withdrawal	
Did not like CST groups and wanted to withdraw	17 (49%)
Health issues	15 (40%)
Difficulties with group time or other participants	2 (6%)
Moved to a different care home	2 (6%)
Participants at Baseline 1 - After completion of CST groups	236

Table 2. Baseline characteristics of 236 participants by allocated group

	Intervention (n=123)	Control (n=113)
Characteristics	Number (%)	Number (%)
Female	80 (65%)	70 (62%)
Ethnicity: white	111 (90%)	104 (92%)
Marital status (widow)	54 (44%)	57 (50%)
Dementia diagnosis (AD)	38 (31%)	35 (31%)
On AChEIs	42 (34%)	34 (30%)
In Care Home	51 (41%)	50 (44%)
	Mean (SD)	Mean (SD)
Age (years):	82.7 (7.9)	83.5 (7.2)
ADAS-Cog	31.1 (14.6)	33.2 (13.0)
QoL-AD	36.1 (4.8)	36.5 (5.7)
MMSE	17.8 (5.6)	17.8 (5.4)
DEMQOL	94.8 (10.9)	95.1 (11.7)
NPI	13.8 (12.9)	11.3 (9.1)
ADCS-ADL	42.7 (17.2)	41.5 (18.1)
Proxy QoL-AD	33.7 (5.9)	33.3 (4.9)
Proxy DEMQOL	102.2 (13.5)	102.2 (11.2)

AChEIs - Acetylcholinesterase Inhibitors

ADAS-Cog - Alzheimer's Disease Assessment Scale – Cognition Subscale

QoL-AD - Quality of Life in Alzheimer's Disease scale

MMSE - Mini-Mental State Examination

DEMQOL - Dementia Quality of Life scale

NPI - Neuropsychiatric Inventory

ADCS-ADL - Alzheimer's Disease Co-operative Study-Activities of Daily Living scale

Table 3. Effects of Maintenance CST on adjusted imputed outcomes at primary and secondary end points

Adjusted outcomes	Primary end point - 6 month follow up				Secondary end point 3-month follow up			
	Treatment	Control	Difference ^a	Significance level	Treatment	Control	Difference ^a	Significance level
	Mean (SE)	Mean (SE)	Mean (95% CI)	Median of 5 imputations	Mean (SE)	Mean (SE)	Mean (95% CI)	Median of 5 imputations
ADAS-Cog ^p	35.94 (2.79)	35.29 (2.85)	-0.65 (-3.71 to 2.42)	.67	35.32 (2.56)	34.47 (2.59)	-0.85 (-3.40 to 1.70)	.27
QoL-AD ^c	35.62 (1.43)	33.84 (1.53)	1.78 (-0.01 to 3.57)	.03	34.29 (1.03)	33.97 (1.04)	0.32 (-0.88 to 1.52)	.54
MMSE ^c	16.34 (1.21)	15.49 (1.25)	0.85 (-0.29 to 1.99)	.15	16.09 (0.88)	15.79 (0.91)	0.30 (-0.72 to 1.31)	.56
DEMQOL ^c	89.13 (3.55)	88.83 (3.56)	0.30 (-2.70 to 3.31)	.87	89.85 (2.34)	90.71 (2.38)	-0.86 (-3.45 to 1.73)	.54
NPI ^b	18.76 (3.78)	20.35 (3.94)	1.58 (-2.67 to 5.84)	.53	14.71 (2.84)	16.18 (2.76)	1.47 (-1.59 to 4.53)	.34
ADCS-ADL ^c	43.29 (2.88)	42.35 (2.87)	0.94 (-2.04 to 3.92)	.54	43.58 (2.32)	40.94 (2.32)	2.64 (0.08 to 5.20)	.04
Proxy QoL-AD ^c	34.12 (1.41)	34.05 (1.41)	0.07 (-1.39 to 1.53)	.95	33.93 (1.05)	32.40 (1.07)	1.53 (0.37 to 2.69)	.01
Proxy DEMQOL ^c	97.75 (3.23)	96.61 (3.21)	1.13 (-2.24 to 4.51)	.50	101.36 (2.67)	98.12 (2.67)	3.24 (0.29 to 6.19)	.03

a Positive differences favour maintenance CST.

b Lower scores show better outcome

c Higher scores show better outcome

Table 4. Effects of Maintenance CST on adjusted imputed outcomes at primary and secondary end point according to ACHEI treatment. The significance levels quoted are for the interaction term of treatment group and receipt of ACHEIs.

Group	n	Baseline		3 months		Interaction p	6 months		Interaction p
		Mean	SE	Model adjusted mean	SE		Model adjusted mean	SE	
ADAS-Cog						.13			.71
ACHEI	34	31.29 (27.17 to 35.41)	2.09	37.05 (31.23 to 42.87)	2.84		36.52 (29.47 to 43.58)	3.53	
TAU	79	34.03 (31.08 to 36.99)	1.5	32.35 (26.77 to 37.92)	2.68		34.67 (28.81 to 40.52)	2.97	
ACHEI/MCST	42	28.65 (25.14 to 32.16)	1.78	36.55 (29.94 to 43.16)	3.16		35.77 (29.33 to 42.22)	3.28	
MCST	81	32.4 (28.91 to 35.89)	1.77	33.85 (28.47 to 39.22)	2.57		35.99 (30.07 to 41.91)	2.98	
QOL-AD						.97			.48
ACHEI	34	37.73 (36.24 to 39.22)	.76	32.81 (30.4 to 35.22)	1.23		33.94 (30.23 to 37.65)	1.86	
TAU	79	35.99 (34.64 to 37.34)	.69	35.13 (32.98 to 37.27)	1.09		33.81 (30.83 to 36.79)	1.52	
ACHEI/MCST	42	37.08 (35.56 to 38.6)	.77	33.14 (30.66 to 35.63)	1.27		34.72 (31.39 to 38.06)	1.7	
MCST	81	35.62 (34.58 to 36.67)	.53	35.45 (33.39 to 37.5)	1.05		36.07 (33.25 to 38.89)	1.44	
NPI						.99			.26
ACHEI	34	12.13 (8.94 to 15.32)	1.62	17.23 (10.77 to 23.69)	3.28		23.78 (14.85 to 32.71)	4.54	
TAU	79	11 (9.01 to 12.99)	1.01	15.12 (9.51 to 20.74)	2.86		17.15 (9.08 to 25.23)	4.11	
MCST/ACHEI	42	16.15 (10.77 to 21.53)	2.73	15.85 (9.17 to 22.53)	3.39		18.21 (9.44 to 26.98)	4.47	

MCST	81	12.65 (10.58 to 14.72)	1.05	13.61 (8.03 to 19.19)	2.82	17.49 (10.05 to 24.93)	3.79
ADL					.80	.80	
ACHEI	34	44.03 (38.35 to 49.71)	2.88	41.51 (36.04 to 46.98)	2.79	42.45 (35.9 to 49)	3.34
TAU	79	40.42 (36.28 to 44.56)	2.10	40.37 (35.53 to 45.22)	2.47	42.22 (36.26 to 48.17)	3.04
MCST/ACHEI	42	48.24 (42.59 to 53.89)	2.87	43.83 (38.21 to 49.45)	2.87	43.91 (37.12 to 50.69)	3.46
MCST only	81	39.78 (36.32 to 43.23)	1.75	43.17 (38.59 to 47.75)	2.34	42.91 (37.18 to 48.64)	2.92
MMSE					.03	.03	
ACHEI	34	18.85 (17.29 to 20.41)	.79	15.26 (13.13 to 17.39)	1.08	14.62 (11.81 to 17.43)	1.40
TAU	79	17.33 (16.08 to 18.58)	.63	16.25 (14.44 to 18.05)	.92	16.26 (13.68 to 18.84)	1.28
MCST/ACHEI	42	18.27 (16.62 to 19.91)	.84	17.17 (15.09 to 19.25)	1.06	17.25 (14.63 to 19.87)	1.33
MCST	81	17.55 (16.29 to 18.8)	.64	15.77 (14.03 to 17.5)	.88	16.26 (13.71 to 18.8)	1.26
DEMQOL					.92	.97	
ACHEI	34	97.90 (94.93 to 100.87)	1.51	89.13 (83.62 to 94.65)	2.81	87.93 (80 to 95.87)	3.90
TAU	79	93.86 (91.07 to 96.66)	1.42	92.25 (87.33 to 97.17)	2.51	89.75 (82.03 to 97.46)	3.74
ACHEI/MCST	42	97.36 (94.35 to 100.36)	1.53	88.99 (83.33 to 94.66)	2.89	87.88 (80.13 to 95.62)	3.86
MCST	81	93.49 (91.02 to 95.97)	1.26	91.04 (86.4 to 95.68)	2.37	90.22 (82.54 to 97.9)	3.69