

Maximilian Salcher, [Huseyin Naci](#), Tyler J. Law, Titus Kuehne, Stephan Schubert, Marcus Kelm

Balloon dilatation and stenting for aortic coarctation: a systematic review and meta-analysis

Article (Accepted version)
(Refereed)

Original citation: Salcher, Maximilian, Naci, Huseyin, Law, Tyler J., Kuehne, Titus, Schubert, Stephan and Kelm, Marcus (2016) *Balloon dilatation and stenting for aortic coarctation: a systematic review and meta-analysis*. [Circulation: Cardiovascular Interventions](#) . ISSN 1941-7632
DOI: [10.1161/CIRCINTERVENTIONS.115.003153](https://doi.org/10.1161/CIRCINTERVENTIONS.115.003153)

© 2016 by [American Heart Association](#)

This version available at: <http://eprints.lse.ac.uk/66303/>

Available in LSE Research Online: June 2016

LSE has developed LSE Research Online so that users may access research output of the School. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LSE Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain. You may freely distribute the URL (<http://eprints.lse.ac.uk>) of the LSE Research Online website.

This document is the author's final accepted version of the journal article. There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

BALLOON DILATATION AND STENTING FOR AORTIC COARCTATION: A SYSTEMATIC REVIEW AND META-ANALYSIS

Maximilian Salcher MSc¹; Huseyin Naci PhD¹; Tyler J. Law MD^{1,2}; Titus Kuehne MD³;
Stephan Schubert MD³; Marcus Kelm MD³ on behalf of CARDIOPROOF Consortium

¹ LSE Health, London School of Economics and Political Science, London, United Kingdom

² Department of Anesthesia, University of Toronto, Ontario, Canada

³ Department of Paediatric Cardiology and Congenital Heart Diseases, Deutsches
Herzzentrum Berlin, Berlin, Germany

* Corresponding Author.

Address for correspondence: LSE Health and Social Care, Cowdray House; London School
of Economics and Political Science; Houghton Street, London WC2A 2AE, United Kingdom.

Tel: +44 2079556959

Fax: +44 2079556803

Email: m.salcher@lse.ac.uk

Subject codes: 27

Total word count: 7,345

ABSTRACT

Background: There is no systematic assessment of available evidence on effectiveness and comparative effectiveness of balloon dilatation and stenting for aortic coarctation.

Methods and results: We systematically searched four online databases to identify and select relevant studies of balloon dilatation and stenting for aortic coarctation based on a *priori* criteria (PROSPERO 2014:CRD42014014418). We quantitatively synthesized results for each intervention from single-arm studies, and obtained pooled estimates for relative effectiveness from pair-wise and network meta-analysis of comparative studies.

Our primary analysis included 15 stenting (423 participants) and 12 balloon dilatation studies (361 participants) including patients ≥ 10 years of age. Post-treatment blood pressure gradient reduction to ≤ 20 mm Hg and ≤ 10 mm Hg was achieved in 89.5% [95%CI 83.7-95.3] and 66.5% [44.1-88.9%] of patients undergoing balloon dilatation and in 99.5% [97.5-100.0%] and 93.8% [88.5-99.1%] of patients undergoing stenting, respectively. Odds of achieving ≤ 20 mm Hg were lower with balloon dilatation as compared to stenting (odds ratio [OR] 0.105 [0.010-0.886]). 30-day survival rates were comparable.

Numerically more patients undergoing balloon dilatation experienced severe complications during admission (6.4% [2.6-10.2%]) compared to stenting (2.6% [0.5-4.7%]). This was supported by meta-analysis of head-to-head studies (OR 9.617 [2.654-34.845]) and network meta-analysis (OR 16.23, 95% credible interval 4.27-62.77) in a secondary analysis in patients ≥ 1 month of age including 57 stenting (3,397 participants) and 62 balloon dilatation studies (4,331 participants).

Conclusions: Despite the limitations of the evidence base consisting predominantly of single-arm studies, our review indicates that stenting achieves superior immediate relief of a relevant pressure gradient compared to balloon dilatation.

Key words: coarctation; balloon; stents; heart defects, congenital; meta-analysis

Introduction

Coarctation of the aorta is a congenital heart disease (CHD) that significantly reduces life expectancy^{1,2} and is associated with increased morbidity even years after successful repair.^{3,4} With an incidence of 3 to 4 cases per 10,000 live births,^{5,6} aortic coarctation accounts for 5-8% of all congenital heart defects and is frequently associated with other CHD such as bicuspid aortic valve disease.⁷

After the initial treatment, ongoing monitoring of patients is recommended by the American Heart Association (AHA) and European Society of Cardiology (ESC) to detect relapse of the disease, disease progression and late complications.^{8,9} Key clinical challenges that may persist post-repair include re-coarctation, persisting arterial hypertension, exercise-induced hypertension and subsequent sequelae with atherosclerosis and coronary artery disease as major cause of death.¹⁰⁻¹²

Two transcatheter interventions exist for treatment of aortic coarctation, balloon dilatation and stenting. Balloon dilatation involves positioning the deflated balloon across the stenotic site and subsequent inflation, stretching the intimal and medial layers of the aorta.^{13,14} This mechanism bears the risk of damaging the aortic wall and can lead to aneurysm formation. Implantation of a stent across the coarcted segment possesses theoretical advantages over balloon dilatation, including lower risk for aortic wall injuries and more sustained relief of the obstruction.^{7,15} It is not clear whether these theoretical advantages hold true, particularly in the long term. Aortic wall injuries and restenosis were also seen in patients undergoing stent implantation,¹⁶ highlighting the need to assess the comparative effectiveness of balloon dilatation and stenting. Guidelines on the management of patients with aortic coarctation from both AHA¹⁷ and ESC⁹ do not provide recommendations on the choice of transcatheter interventions. Considerations regarding the effectiveness and safety of the treatment options are largely based on C-level (expert consensus) and only rarely on B-level (non-randomized) evidence.

The evidence available on the effectiveness and comparative effectiveness of balloon dilatation and stenting for aortic coarctation has not yet been collected systematically. Furthermore, no systematic comparison of the effectiveness of these two transcatheter interventions exists.

Methods

Study identification and selection

A review protocol was developed and subsequently made publicly available on the PROSPERO website of the University of York Centre for Reviews and Dissemination at the onset of our work (<http://goo.gl/ZhXomV>).¹⁸ The review was set up to assess the effectiveness of balloon dilatation and stenting as well as their comparative effectiveness in the treatment of patients with aortic coarctation. The main parameters of our systematic review are summarized in Table 1.

We searched online databases MEDLINE (via PubMed), CINAHL Plus, EMBASE, and the Cochrane Library with search strings containing word and phrase match terms as well as database specific subject headings. The search strategy was constructed to find relevant articles on balloon angioplasty and stenting in aortic coarctation and is available in the online supplement. We additionally searched reference lists of four review articles^{7,19–21} and three clinical practice guidelines.^{8,9,17} We did not distinguish between different types of stents (e.g. bare metal; covered; balloon- and self-expandable).

Pre-defined inclusion criteria included head-to-head comparisons of balloon angioplasty and stenting. Given the paucity of comparative studies, we also included single-arm studies which did not use any comparator ('case series'), as well as studies comparing one of the two relevant procedures to a third comparator, surgical repair.

Studies were eligible for inclusion if they had a minimum of 15 patients with native or re-coarctation per relevant study arm. We thus excluded very small studies but were still able to systematically capture the vast majority of evidence on this topic. We included studies published from 1990 through 2014 to limit the potential bias in treatment results originating from outdated knowledge and other contextual differences before that date. Stenting and balloon dilatation are considered treatment options mainly in adolescents and adults. For our primary analysis, we restricted the sample to studies and study arms including patients aged ≥ 10 years to capture the patient population for which both interventions are appropriate. For our secondary analysis, we relaxed the age restriction and included all studies including patients ≥ 1 month of age. We subjected the results of all secondary analyses to extensive sensitivity testing with respect to patient age.

We excluded studies that were not in English; published before 1990; did not report any of our pre-specified outcomes; or that were conducted in animals. We further excluded comments; editorials; letters; and conference abstracts.

Search results from the database searches were merged and obvious duplicates were removed. One researcher scanned articles at the title and abstract level for eligibility according to pre-specified inclusion and exclusion criteria. Full texts of articles deemed eligible were then independently assessed for inclusion by two researchers. Deviating decisions on eligibility were resolved by discussion and consensus between the two researchers. Each eligible study was then re-examined and pre-specified data were extracted independently by two researchers.

For the few studies with overlapping populations, we included the report with the most detailed or relevant data, or the most recent publication in order to maximize follow-up time. For larger studies, we contacted corresponding authors to ascertain whether study participants were indeed from the same cohort and used the information provided to exclude duplicate patients.

Outcomes

We report pre-specified outcomes as proportions of the total number of patients in each study arm. Our primary measure of immediate treatment success was the proportion of patients achieving a post-treatment blood pressure gradient ≤ 20 mm Hg. We adopted this cut-off value because of its widespread use in the literature and because it is the threshold below which patients are generally not considered candidates for intervention according to AHA and ESC guidelines.^{9,17} In addition, we used a second cut-off value, ≤ 10 mm Hg, as a stricter criterion for treatment success, reflecting some skepticism towards establishment of treatment success through a gradient threshold that is just below treatment indication. Adapting the categorization proposed by Vitiello et al.,²² we report the proportion of patients with (1) severe complications during intervention or before discharge, and (2) minor complications. Severe complications were defined as life-threatening events requiring immediate therapy; permanent functional or anatomic lesion; any aortic wall injury (dissection/ acute aneurysm); and unexpected major drug side effects. Minor complications included balloon rupture; stent migration; bleeding at access site; loss of femoral pulse; and other complications that were not deemed severe by study investigators. We further report 30-day survival rates, and the proportion of patients with reinterventions for restenosis or vascular complications related to the initial intervention at follow-up. We standardized reintervention rates in our analysis of single-arm studies to represent annual reinterventions per 100 patient-years of follow-up. Linearized reintervention rates were calculated as $[\text{events}/(\text{sample size} \times \text{mean follow-up time})] \times 100$. We then obtained separate pooled estimates of linearized reintervention rates for studies with mean follow-up time ≤ 1 year; between 1 and 3 years; and >3 years.

Although our data extraction included additional outcomes, lack of data precluded meaningful analysis.

Statistical analysis

Our primary analysis focused on patients ≥ 10 years of age. Studies in this age group included very small numbers of patients. We therefore relaxed the age restriction and broadened the eligible evidence base in secondary analyses, allowing for more precise estimates of treatment effectiveness and safety.

For both primary and secondary analyses, we conducted three types of statistical analyses. First, using information from all case series and study arms within one intervention type, we computed overall estimates for proportions of participants with any given outcome. We assessed statistical heterogeneity between study results using the I^2 statistic.^{23,24} A fixed-effect model was used for outcomes with low between-study heterogeneity ($I^2 < 25\%$) and a random-effects model for outcomes with moderate to high heterogeneity ($I^2 \geq 25\%$), as recommended by the Cochrane Collaboration. For studies with a proportion of 0 or 1 for any given outcome, we imputed the average of the variances of the other studies to obtain an estimate of the variance.²⁵

Second, we synthesised the results from comparative studies using pair-wise meta-analysis. In a similar fashion to our analysis of single-arm studies, we first visually inspected heterogeneity of results and then assessed between-study heterogeneity using the I^2 statistic. Informed by this, a fixed-effect or random-effects model was used to obtain odds ratios (OR) for each outcome, comparing the odds for each outcome (number of patients with event compared to patients without event) in patients undergoing balloon dilatation with the odds in patients undergoing stenting. Meta-analyses of single-arm and comparative studies were carried out in STATA, version 13.

Finally, we conducted network meta-analyses for studies including one or both interventions of interest and a third intervention, surgery for aortic coarctation. Unlike traditional pair-wise meta-analysis, which pools the results of direct head-to-head studies between two treatment options, network meta-analysis allows for the combination of both direct and indirect sources

of evidence to compare multiple treatments that may not have been directly compared to each other in head-to-head studies.²⁶ It combines the results of studies that compare treatments A vs. C and the results of studies that compare treatments B vs. C to indirectly estimate the comparative effectiveness of treatments A vs. B.

In this analysis, we were able to widen the evidence base for comparisons of balloon dilatation and stenting by including studies comparing either one of the two transcatheter interventions with surgery. The primary assumption of network meta-analysis is that the pooled studies are comparable in terms of relative treatment effect modifiers (i.e., in terms of variables that have a known influence on the outcomes).²⁷⁻²⁹ We qualitatively evaluated the comparability of the studies included in the network meta-analyses in terms of key baseline characteristics and visually inspected the influence of these baseline variables on outcomes.

Our network meta-analysis model combined study-level treatment effects using Bayesian methods in WinBUGS.²⁹ This was based on modelling the outcomes in every treatment group of every study, and specifying the relations among the relative effects across studies making different comparisons. The model adopted random-effects, which took into account potential heterogeneity by assuming that each treatment was drawn from the same distribution whose mean and variance were estimated from the data. We present the results from network meta-analysis as OR and 95% credible intervals (CrI). Credible intervals indicate a 95% probability that the true OR falls within the observed range of estimates. If a 95% CrI does not include the null value 1.00, this can be interpreted as indicating <5% probability that there is no difference between the two intervention groups.

We only report the findings of network meta-analyses for which history plots suggested successful convergence in WinBUGS. In all network meta-analyses, we qualitatively evaluated the consistency of relative treatment effects obtained from both the single-arm and comparative studies. The consistency of the relative treatment effects were visually inspected for potential differences between estimates obtained from three sets of analyses.

We checked for discrepancy in terms of the direction of effect, as well as its magnitude, and confirmed that all 95% intervals greatly overlapped, which suggested adequate consistency.

Results

The literature search process is presented in the PRISMA flow chart with the number of articles screened and excluded at every stage (Figure 1). None of the studies including both balloon dilatation and stenting patients was a randomized controlled trial. The evidence base therefore primarily consists of single-arm studies for both stenting and balloon dilatation, including all the participants of case series and respective study arms from multiple-arm studies. There was a limited number of multiple-arm studies directly comparing the interventions of interest.

For the primary analysis, we identified 15 stenting studies or study arms with 423 participants, 12 balloon dilatation studies or study arms with 361 participants, 2 studies comparing the two interventions,^{30,31} and 1 study comparing stenting with surgery³² (Figure 2, Panel A. Full list of included studies provided in the online supplement). Mean follow-up time ranged from 1 to 12 years in balloon dilatation studies and from 10 months to 4.7 years in stenting studies. Single-arm studies for balloon dilatation were published between 1992 and 2009, and single-arm studies for stenting between 2001 and 2013. Comparative studies including both balloon dilatation and stenting patients were published in 2003 and 2005.

For the secondary analysis, we identified 57 stenting studies (3,397 participants), 62 balloon dilatation studies (4,331 participants). 7 studies compared the two interventions (5 studies with patients undergoing stenting or balloon dilatation and 2 studies including surgery as common comparator. Figure 2, Panel B). We obtained additional unpublished data for one of the comparative studies included in the secondary analysis directly from the authors.³³

Exploration of differences between patients undergoing balloon dilatation and stenting

Baseline characteristics for patients undergoing balloon dilatation and stenting are shown in Table 2. Statistically significant differences were found in pre-treatment blood pressure gradient and the proportion of patients with native/recurrent coarctation between groups.

For variables that were statistically significantly different between groups, and other variables that were considered to have a systematic effect on outcomes, we constructed forest plots to examine any potential effect on four key outcomes: proportion of patients achieving a gradient reduction ≤ 20 mm Hg; proportion achieving a gradient reduction ≤ 10 mm Hg; 30-day mortality; proportion of patients with severe complications before discharge. An

example is Figure 3^{30,30,30,30,31,31,31,31,34,34,35,35,36,36,37,37,38,38,39,39,40,40,41,41–}

43,43,44,44,45,45,46,46,47,47,48,48,49,49,50,50[,] which demonstrates that for both interventions, average pre-treatment blood pressure gradient (Panel A) and proportion of patients with native coarctation (Panel B) do not systematically influence the proportion of patients with successful treatment. Further exploration of the effect of patient baseline characteristics on key outcomes is provided in the online supplement (Figures S1-S12). We also assessed the impact of mean age and found no discernible systematic effect on the four outcomes. We therefore pooled the results of individual studies.

Results from single-arm studies

Figure 4 shows pooled results and 95% CIs for each outcome in single-arm studies.

Treatment success before discharge was more often achieved in patients undergoing stenting compared to balloon dilatation. The proportion of patients achieving post-treatment gradient ≤ 20 mm Hg was 0.895 (95%CI 0.837-0.953; I^2 56.7%) in balloon dilatation studies vs 0.995 (95%CI 0.975-1.000; I^2 0.0%) in stenting studies, and proportion of patients achieving post-treatment gradient ≤ 10 mm Hg was 0.665 (95%CI 0.441-0.889; I^2 93.2%) in balloon dilatation studies vs 0.938 (95%CI 0.885-0.991; I^2 0.0%) in stenting studies.

For the proportion of patients with severe complications before discharge we obtained a pooled estimate of 0.064 (95%CI 0.026-0.102; I^2 31.3%) for patients undergoing balloon dilatation and 0.026 (95%CI 0.005-0.047; I^2 0.0%) for patients undergoing stenting. Pooled estimates for the proportion with minor complications before discharge were 0.128 (95%CI 0.012-0.244; I^2 91.0%) for patients undergoing balloon dilatation and 0.073 (95%CI 0.041-0.106; I^2 7.8%) for patients receiving stents.

All patients undergoing balloon dilatation in the identified studies survived at 30 days and the pooled estimate for patients undergoing stenting was 0.999 (95%CI 0.988-1.000; I^2 0.0%).

At follow-up, the pooled estimates for the proportion of patients with reinterventions for re-coarctation or vascular complications related to the initial intervention were 0.061 (95%CI 0.026-0.096; I^2 0.0%) for patients undergoing balloon dilatation and 0.085 (95%CI 0.039-0.131; I^2 60.5%) for patients undergoing stenting. The pooled linearized reintervention rate was 0.9 (95% CI 0.3-1.4; I^2 0.0%) per 100 patient-years of follow-up for patients undergoing balloon dilatation and 3.3 (95% CI 1.6-5.0; I^2 37.5%) per 100 patient-years for patients undergoing stenting.

Results from comparative studies

Figure 5 shows pooled OR and 95% CIs for comparative studies. Patients undergoing balloon dilatation were significantly less likely to achieve treatment success compared to patients undergoing stenting as measured by the proportion of patients achieving a blood pressure gradient ≤ 20 mm Hg (OR 0.105, 95%CI 0.010-0.886; I^2 0.0%).

No statistically significant difference was found for minor complications before discharge (OR 0.669, 95%CI 0.035-12.742; I^2 58.9%).

Focusing on the two main indications for reinterventions at follow-up the pooled OR included 1.00, but suggested a tendency towards increased risk after balloon dilatation. The OR for re-coarctation at follow-up in patients undergoing balloon dilatation vs. stenting was 7.010

(95%CI 0.794-61.92; I^2 0.0%), and the OR for aortic wall injuries at follow-up was 3.340 (95%CI 0.477-23.367; I^2 0.0%).

Secondary analysis

In the extended sample, including studies with patients <10 years of age, we found statistically significant differences in patient baseline age, weight, and pre-treatment gradient (Table S1 in online supplement). However, forest plots did not indicate a systematic impact of patient baseline characteristics on key outcomes (Figures S13-S24).

Compared to the primary analysis, pooled estimates of single-arm studies in the secondary analysis were generally more precise and showed less favorable results, but confirmed the overall direction of effect in stenting vs. balloon dilatation studies in all but two outcomes (Figure 4). The pooled linearized reintervention rate did not show a significant difference between balloon dilatation (3.8 events per 100 patient-years of follow-up, 95% CI 2.9-4.7; I^2 81.6%) and stenting (5.4 events, 95% CI 4.1-6.7; I^2 78.8%). A higher proportion of patients had re-coarctation at follow-up after balloon dilatation (0.165, 95% CI 0.136-0.194; I^2 77.1%) compared to stenting (0.048, 95% CI 0.029-0.067; I^2 54.2%).

For comparative studies, the secondary analysis confirmed the results of the primary analysis with more precise estimates (Figure 5). Due to data availability, we were also able to analyze additional outcomes. While the OR for the primary criterion of treatment success (blood pressure gradient \leq 20mm Hg) included 1.00 (OR 0.663, 95%CI 0.358-1.229; I^2 40.3%), the stricter threshold of \leq 10mm Hg was statistically significantly less likely to be achieved by patients undergoing balloon dilatation compared to patients undergoing stenting (OR 0.435, 95%CI 0.320-0.591; I^2 20.3%). The pooled OR of patients undergoing balloon dilatation with severe complications before discharge compared to patients undergoing stenting was 9.617 (95%CI 2.654-34.845; I^2 53.9%), indicating considerably higher rate of complications in balloon studies. Comparing the odds of reinterventions at follow-up in

patients after balloon dilatation to the odds after stenting, the OR was 0.65 (95%CI 0.38-1.10; I^2 0.0%).

Results from network meta-analysis

Network meta-analysis could only be conducted for the extended sample of studies because of limited data availability. We did not obtain precise estimates for the comparative effectiveness of balloon dilatation and stenting for all outcomes. We therefore only report the results for three outcomes for which our analyses achieved convergence.

Using surgery as a common comparator, we observed higher odds for experiencing severe complications before discharge in patients undergoing balloon dilatation compared to patients undergoing stenting (OR 16.23, 95%CrI 4.27-62.77). The majority of severe complications in balloon dilatation and stenting patients consisted of damages to the aortic wall.

In terms of minor complications, we found no statistically significant difference between the two interventions of interest: OR for patients undergoing balloon dilatation vs stenting 0.95 (95%CrI 0.23-4.16). Similarly, we found no statistically significant difference between the two transcatheter interventions for reinterventions at follow-up (OR patients undergoing balloon dilatation vs stenting 0.70, 95%CrI 0.35-1.28).

Sensitivity analysis

Results of sensitivity analyses are available in the online supplement.

To test the sensitivity of our comparative effectiveness results to potential overreporting of desirable and underreporting of undesirable events in case series, we obtained pooled estimates for each outcome excluding (1) the study reporting the most favorable results, and (2) the two studies reporting the most favorable results. We did not find materially different results for the comparative effectiveness of stenting and balloon dilatation (Figure S25 in the online supplement).

We plotted study publication year against the proportion of patients with post-treatment gradient ≤ 20 mm Hg and linearized reintervention rate to study the potential impact of advanced technology and experience. We did not detect systematically better results in more recent studies (Figures S27 and S28 in the online supplement).

Discussion

Immediate and follow-up outcomes

The ultimate aim of coarctation treatment has traditionally been the complete relief of a pressure gradient.⁵¹ While both treatments were capable of reducing the pressure gradient in patients aged ≥ 10 years, stenting was more frequently associated with a gradient reduction to ≤ 20 mm Hg and ≤ 10 mm Hg in our analyses and thus showed better immediate relief of the stenosis compared to balloon dilatation.

We observed a tendency towards higher risk of severe complications during intervention or before discharge after stenting compared to balloon dilatation in our primary analysis. This finding was amplified when we included patients below 10 years of age in our secondary analysis. Our results, which highlight the advantage of stenting with respect to patient safety, confirm and extend the findings reported by the studies of the Congenital Cardiovascular Interventional Study Consortium (CCISC).^{51–53} Our results were consistent across the three types of statistical analysis conducted (meta-analysis of case series; pairwise meta-analysis; and network meta-analysis). Severe complications consisted of damages to the aortic wall in most cases. Other severe complications were rare.

Sustained relief of the obstruction and therefore the prevention of recurrent coarctation is an often-cited advantage of stenting.³¹ Accordingly, a lower number of reinterventions for recurrent coarctation could be expected. Contradicting this theoretical advantage, we found no statistically significant difference in the proportion of patients with reinterventions at follow-up. The reason to re-intervene is not consistently reported across the studies, and

different arguments could lead to reintervention in the balloon dilatation group (e.g. restenosis due to growth in younger patients) than in the stenting group (e.g. neo-intimal proliferation). In comparison to balloon dilatation, less need for re-coarctation repair could be offset by the need for stent redilatation, which would only reflect a planned staged repair approach in very few patients with sub-atretic coarctation.

Even after successful stenting gradients frequently remain. The shape of the entire anatomical region as well as flow features can show an impact. While current guidelines recommend reintervention once gradient thresholds are reached, Computational Fluid Dynamic simulations carry the potential for more personalized decision making in the future.^{54,55}

Overall, our results focusing on short and mid-term outcome may be seen as clear arguments towards stent placement. However, the issue of long term outcome has only incompletely been studied and remains more difficult to assess. Considering that most children undergo their first intervention at infancy or early childhood, long term consequences will be affected by placement of material that was originally planned for smaller vascular anatomy. Criteria for the decision of which treatment to use include patient age, history, and anatomy of the coarcted segment.^{19,30} This suggests that a 'one treatment fits all' approach is not appropriate. Stent repair seems to be a preferred method in adults and older children, while its use in infants and younger children will be to bridge the time to surgical repair.

State of the evidence

There is widespread consensus that randomized controlled trials (RCTs) are the gold standard for establishing the effectiveness and safety of clinical interventions.⁵⁶ However, in our systematic review we found no RCT comparing balloon dilatation and stenting. We found that over 7,700 patients have been treated in the major clinical centers by either balloon dilatation or stenting for aortic coarctation over the last 25 years, and yet the evidence base

for interventional treatment for this condition is confined to mostly small case series and few large collaborative observational studies. Previous collaborative efforts such as the CCISC⁵¹ and the Valvuloplasty and Angioplasty of Congenital Anomalies Registry (VACA)⁵⁷ have shown that pooling patient data from a considerable number of centers is feasible. Such future collaborations across centers, regions, and countries would significantly improve the current state of evidence on the effectiveness of treatment alternatives for aortic coarctation and generate much needed information regarding the comparative effectiveness of balloon dilatation and stenting in this patient population, ideally using more rigorous study designs such as RCTs.

Our systematic review and meta-analysis shares the limitations of the individual studies. Due to a clear lack of controlled studies in the literature, we included case series which rank low in the hierarchy of evidence.⁵⁸ Indeed, even when using control groups, observational study designs are susceptible to bias in several ways.⁵⁹ One particular area of concern in our review was potential selection bias, as allocation of patients to a given treatment was at the cardiologist's discretion. However, our extensive sensitivity studies suggested that, while there are some differences in patient characteristics between studies evaluating balloon dilatation and stenting, these do not seem to systematically affect the outcomes. Our exploration of the potential impact of patient characteristics on outcomes was limited to the study level and it is therefore possible that confounding effects at the individual patient level were concealed. In our secondary analysis, the lowest mean age in stenting studies was 8 years, while it was around 1 month in balloon dilatation studies. Although we did not find a detectable effect of mean age on key outcomes, we cannot fully rule out a confounding effect in stenting patients of very young age in these secondary analyses. The findings therefore cannot necessarily be applied to young patients.

In our network meta-analysis we made use of surgery as a common comparator between our two interventions of interest. Surgery for aortic coarctation may not be used in the same patients as balloon dilatation and stenting.¹⁹ However, this is not necessarily a limitation of

our network meta-analysis. Similar characteristics of patients eligible for surgical intervention ensured that the common comparator was consistent across different studies included in the network meta-analysis. Nevertheless, we do not report the results of our network meta-analysis as the base-case. Rather, these findings support and extend the findings of our analyses based on single-arm and comparative studies.

Despite significant between-study heterogeneity in the evidence base, we decided to pool study results to gain insight about their comparative effectiveness. Applying narrative rather than quantitative synthesis can be misleading as it does not provide a clear approach towards heterogeneity.⁶⁰ Exploring heterogeneity in study results, we found that patient characteristics that could potentially impact on outcomes did not explain the observed variability.

Due to data availability, reporting limitations and inconsistencies, the list of outcomes reported in this paper does not include all outcomes that were prespecified in our review protocol.

In conclusion, our review suggests that stenting achieves better immediate relief of aortic coarctation. In addition, we found some evidence that patients undergoing stenting may experience fewer severe complications during their hospital admission compared to those undergoing balloon dilatation.

Funding Sources

This systematic review was carried out as part of CARDIOPROOF, a proof-of-concept study funded by the European Commission under FP7.

Appendix (author group)

CARDIOPROOF partners are the following:

Edwin Morley-Fletcher (Lynkeus, Rome, Italy)

Titus Kuehne (Deutsches Herzzentrum Berlin, Berlin, Germany)

Anja Hennemuth (Fraunhofer MEVIS, Bremen, Germany)

David Manset (gnúbila, Argonay, France)

Alistair Mcguire (London School of Economics and Political Science, London, United Kingdom)

Gernot Plank (Medizinische Universität Graz, Graz, Austria)

Olivier Ecabert (Siemens AG, Munich, Germany)

Giacomo Pongiglione (Ospedale Pediatrico Bambino Gesù, Rome, Italy)

Vivek Muthurangu (University College London, London, United Kingdom)

Disclosures

None.

References

1. Campbell M. Natural history of coarctation of the aorta. *Br Heart J*. 1970;32:633–640.
2. Verheugt CL, Uiterwaal CS, Grobbee DE, Mulder BJ. Long-term prognosis of congenital heart defects: a systematic review. *Int J Cardiol*. 2008;131:25–32.
3. Celermajer DS, Greaves K. Survivors of coarctation repair: fixed but not cured. *Heart*. 2002;88:113–114.
4. Toro-Salazar OH, Steinberger J, Thomas W, Rocchini AP, Carpenter B, Moller JH. Long-term follow-up of patients after coarctation of the aorta repair. *Am J Cardiol*. 2002;89:541–547.
5. Grech V. Diagnostic and surgical trends, and epidemiology of coarctation of the aorta in a population-based study. *Int J Cardiol*. 1999;68:197–202.

6. Hoffman JIE, Kaplan S. The incidence of congenital heart disease. *J Am Coll Cardiol.* 2002;39:1890–1900.
7. Tanous D, Benson LN, Horlick EM. Coarctation of the aorta: evaluation and management. *Curr Opin Cardiol.* 2009;24:509–515.
8. Warnes CA, Williams RG, Bashore TM, Child JS, Connolly HM, Dearani JA, del Nido P, Fasules JW, Graham JTP, Hijazi ZM, Hunt SA, King ME, Landzberg MJ, Miner PD, Radford MJ, Walsh EP, Webb GD. ACC/AHA 2008 Guidelines for the Management of Adults With Congenital Heart Disease. A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Develop Guidelines on the Management of Adults With Congenital Heart Disease) Developed in Collaboration With the American Society of Echocardiography, Heart Rhythm Society, International Society for Adult Congenital Heart Disease, Society for Cardiovascular Angiography and Interventions, and Society of Thoracic Surgeons. *J Am Coll Cardiol.* 2008;52:e143–e263.
9. Baumgartner H, Bonhoeffer P, De Groot NMS, de Haan F, Deanfield JE, Galie N, Gatzoulis MA, Gohlke-Baerwolf C, Kaemmerer H, Kilner P, Meijboom F, Mulder BJM, Oechslin E, Oliver JM, Serraf A, Szatmari A, Thaulow E, Vouhe PR, Walma E, Task Force on the Management of Grown-up Congenital Heart Disease of the European Society of Cardiology (ESC), Association for European Paediatric Cardiology (AEPC), ESC Committee for Practice Guidelines (CPG). ESC Guidelines for the management of grown-up congenital heart disease (new version 2010). *Eur Heart J.* 2010;31:2915–2957.
10. Correia AS, Gonçalves A, Paiva M, Sousa A, Oliveira SM, Lebreiro A, Cruz C, Maciel MJ, Correia AS, Gonçalves A, Paiva M, Sousa A, Oliveira SM, Lebreiro A, Cruz C, Maciel MJ. Long-term follow-up after aortic coarctation repair: The unsolved issue of exercise-induced hypertension. *Rev Port Cardiol.* 2013;32:879–883.

11. Cohen M, Fuster V, Steele PM, Driscoll D, McGoon DC. Coarctation of the aorta. Long-term follow-up and prediction of outcome after surgical correction. *Circulation*. 1989;80:840–845.
12. O’Sullivan J. Late hypertension in patients with repaired aortic coarctation. *Curr Hypertens Rep*. 2014;16:421.
13. Vergales JE, Gangemi JJ, Rhueban KS, Lim DS. Coarctation of the Aorta - The Current State of Surgical and Transcatheter Therapies. *Curr Cardiol Rev*. 2013;9:211–219.
14. Rosenthal E. Coarctation of the aorta from fetus to adult: curable condition or life long disease process? *Heart*. 2005;91:1495–1502.
15. Cardoso G, Abecasis M, Anjos R, Marques M, Koukoulis G, Aguiar C, Neves JP. Aortic Coarctation Repair in the Adult. *J Card Surg*. 2014;29:512–518.
16. Godart F. Intravascular stenting for the treatment of coarctation of the aorta in adolescent and adult patients. *Arch Cardiovasc Dis*. 2011;104:627–635.
17. Feltes TF, Bacha E, Beekman RH, Cheatham JP, Feinstein JA, Gomes AS, Hijazi ZM, Ing FF, Moor M de, Morrow WR, Mullins CE, Taubert KA, Zahn EM. Indications for Cardiac Catheterization and Intervention in Pediatric Cardiac Disease A Scientific Statement From the American Heart Association. *Circulation*. 2011;123:2607–2652.
18. Salcher M, Naci H. Balloon angioplasty vs stenting: systematic review of the effectiveness of two interventions for native and recurrent aortic coarctation PROSPERO: 2014CRD42014014418. 2014;Available from: http://www.crd.york.ac.uk/PROSPERO/display_record.asp?ID=CRD42014014418
19. Luijendijk P, Bouma BJ, Groenink M, Boekholdt M, Hazekamp MG, Blom NA, Koolbergen DR, de Winter RJ, Mulder BJM. Surgical versus percutaneous treatment of aortic coarctation: new standards in an era of transcatheter repair. *Expert Rev Cardiovasc Ther*. 2012;10:1517–1531.
20. Hu ZP, Wang ZW, Dai XF, Zhan BT, Ren W, Li LC, Zhang H, Ren ZL. Outcomes of surgical versus balloon angioplasty treatment for native coarctation of the aorta: a meta-analysis. *Ann Vasc Surg*. 2014;28:394–403.

21. Pádua LMS, Garcia LC, Rubira CJ, de Oliveira Carvalho PE. Stent placement versus surgery for coarctation of the thoracic aorta. *Cochrane Database Syst Rev* [Internet]. 2012. Available from:
<http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD008204.pub2/abstract>
22. Vitiello R, McCrindle BW, Nykanen D, Freedom RM, Benson LN. Complications associated with pediatric cardiac catheterization. *J Am Coll Cardiol*. 1998;32:1433–1440.
23. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med*. 2002;21:1539–1558.
24. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ*. 2003;327:557–560.
25. Tabaee A, Anand VK, Barrón Y, Hiltzik DH, Brown SM, Kacker A, Mazumdar M, Schwartz TH. Endoscopic pituitary surgery: a systematic review and meta-analysis. *J Neurosurg*. 2009;111:545–554.
26. Lu G, Ades AE. Combination of direct and indirect evidence in mixed treatment comparisons. *Stat Med*. 2004;23:3105–3124.
27. Lu G, Ades AE. Assessing Evidence Inconsistency in Mixed Treatment Comparisons. *J Am Stat Assoc*. 2006;101:447–459.
28. Jansen JP, Naci H. Is network meta-analysis as valid as standard pairwise meta-analysis? It all depends on the distribution of effect modifiers. *BMC Med*. 2013;11:159.
29. Dias S, Welton NJ, Caldwell DM, Ades AE. Checking consistency in mixed treatment comparison meta-analysis. *Stat Med*. 2010;29:932–944.
30. Zabal C, Attie F, Rosas M, Buendía-Hernández A, García-Montes JA. The adult patient with native coarctation of the aorta: balloon angioplasty or primary stenting? *Heart*. 2003;89:77–83.
31. Pedra CAC, Fontes VF, Esteves CA, Pilla CB, Braga SLN, Pedra SRF, Santana MVT, Silva MAP, Almeida T, Sousa JEMR. Stenting vs. balloon angioplasty for discrete

- unoperated coarctation of the aorta in adolescents and adults. *Catheter Cardiovasc Interv.* 2005;64:495–506.
32. Roselli EE, Qureshi A, Idrees J, Lima B, Greenberg RK, Svensson LG, Pettersson G. Open, hybrid, and endovascular treatment for aortic coarctation and postrepair aneurysm in adolescents and adults. *Ann Thorac Surg.* 2012;94:751-756.
 33. Forbes TJ, Garekar S, Amin Z, Zahn EM, Nykanen D, Moore P, Qureshi SA, Cheatham JP, Ebeid MR, Hijazi ZM, Sandhu S, Hagler DJ, Sievert H, Fagan TE, Ringewald J, Du W, Tang L, Wax DF, Rhodes J, Johnston TA, Jones TK, Turner DR, Pedra CAC, Hellenbrand WE. Procedural results and acute complications in stenting native and recurrent coarctation of the aorta in patients over 4 years of age: A multi-institutional study. *Catheter Cardiovasc Interv.* 2007;70:276–285.
 34. Bentham JR, English K, Ballard G, Thomson JDR. Effect of interventional stent treatment of native and recurrent coarctation of aorta on blood pressure. *Am J Cardiol.* 2013;111:731–736.
 35. Tanous D, Collins N, Dehghani P, Benson LN, Horlick EM. Covered stents in the management of coarctation of the aorta in the adult: initial results and 1-year angiographic and hemodynamic follow-up. *Int J Cardiol.* 2010;140:287–295.
 36. Macdonald S, Thomas SM, Cleveland TJ, Gaines PA. Angioplasty or stenting in adult coarctation of the aorta? A retrospective single center analysis over a decade. *Cardiovasc Interv Radiol.* 2003;26:357–364.
 37. Harrison DA, McLaughlin PR, Lazzam C, Connelly M, Benson LN. Endovascular stents in the management of coarctation of the aorta in the adolescent and adult: one year follow up. *Heart.* 2001;85:561–566.
 38. Wheatley GH 3rd, Koullias GJ, Rodriguez-Lopez JA, Ramaiah VG, Diethrich EB. Is endovascular repair the new gold standard for primary adult coarctation? *Eur J Cardiothorac Surg.* 2010;38:305–310.

39. Sadiq M, Rehman AU, Qureshi AU, Qureshi SA. Covered stents in the management of native coarctation of the Aorta - Intermediate and long-term follow-up. *Catheter Cardiovasc Interv.* 2013;82:511–518.
40. Haji-Zeinali AM, Ghazi P, Alidoosti M. Self-expanding nitinol stent implantation for treatment of aortic coarctation. *J Endovasc Ther.* 2009;16:224–232.
41. Tyagi S, Singh S, Mukhopadhyay S, Kaul UA. Self- and balloon-expandable stent implantation for severe native coarctation of aorta in adults. *Am Heart J.* 2003;146:920–928.
42. De Giovanni JV, Lip GYH, Osman K, Mohan M, Islim IF, Gupta J, Watson RDS, Singh SP. Percutaneous balloon dilatation of aortic coarctation in adults. *Am J Cardiol.* 1996;77:435–439.
43. Koerselman J, De Vries H, Jaarsma W, Muyldermans L, Ernst JMPG, Plokker HWM. Balloon angioplasty of coarctation of the aorta: A safe alternative for surgery in adults: Immediate and mid-term results. *Catheter Cardiovasc Interv.* 2000;50:28–33.
44. Paddon AJ, Nicholson AA, Ettles DF, Travis SJ, Dyet JF. Long-term follow-Up of percutaneous balloon angioplasty in adult aortic coarctation. *Cardiovasc Interv Radiol.* 2000;23:364–367.
45. Walhout RJ, Suttorp MJ, Mackaij GJ, Ernst JM, Plokker HW. Long-term outcome after balloon angioplasty of coarctation of the aorta in adolescents and adults: Is aneurysm formation an issue? *Catheter Cardiovasc Interv.* 2009;73:549–556.
46. Fawzy ME, Fathala A, Osman A, Badr A, Mostafa MA, Mohamed G, Dunn B. Twenty-two years of follow-up results of balloon angioplasty for discreet native coarctation of the aorta in adolescents and adults. *Am Heart J.* 2008;156:910–917.
47. Schrader R, Bussmann WD, Jacobi V, Kadel C. Long-term effects of balloon coarctation angioplasty on arterial blood pressure in adolescent and adult patients. *Cathet Cardiovasc Diagn.* 1995;36:220–225.
48. Hassan W, Awad M, Fawzy ME, Omrani AA, Malik S, Akhras N, Shoukri M. Long-term effects of balloon angioplasty on left ventricular hypertrophy in adolescent and adult

- patients with native coarctation of the aorta. Up to 18 years follow-up results. *Catheter Cardiovasc Interv.* 2007;70:881–886.
49. Tyagi S, Arora R, Kaul UA, Sethi KK, Gambhir DS, Khalilullah M. Balloon angioplasty of native coarctation of the aorta in adolescents and young adults. *Am Heart J.* 1992;123:674–680.
 50. Biswas PK, Mitra K, De S, Banerjee AK, Roy S, Das Biswas A, Biswas A, Chatterjee SS, Maity AK. Follow-up results of balloon angioplasty for native coarctation of aorta. *Indian Heart J.* 1996;48:673–676.
 51. Forbes TJ, Kim DW, Du W, Turner DR, Holzer R, Amin Z, Hijazi Z, Ghasemi A, Rome JJ, Nykanen D, Zahn E, Cowley C, Hoyer M, Waight D, Gruenstein D, Javois A, Foerster S, Kreutzer J, Sullivan N, Khan A, Owada C, Hagler D, Lim S, Canter J, Zellers T, CCISC Investigators. Comparison of surgical, stent, and balloon angioplasty treatment of native coarctation of the aorta: an observational study by the CCISC (Congenital Cardiovascular Interventional Study Consortium). *J Am Coll Cardiol.* 2011;58:2664–2674.
 52. Harris KC, Du W, Cowley CG, Forbes TJ, Kim DW, On Behalf of the Congenital Cardiac Intervention Study Consortium. A prospective observational multicenter study of balloon angioplasty for the treatment of native and recurrent coarctation of the aorta. *Catheter Cardiovasc Interv.* 2014;83:1116–1123.
 53. Holzer R, Qureshi S, Ghasemi A, Vincent J, Sievert H, Gruenstein D, Weber H, Alday L, Peirone A, Zellers T, Cheatham J, Slack M, Rome J. Stenting of aortic coarctation: acute, intermediate, and long-term results of a prospective multi-institutional registry-- Congenital Cardiovascular Interventional Study Consortium (CCISC). *Catheter Cardiovasc Interv.* 2010;76:553–563.
 54. Kelm M, Goubergrits L, Fernandes JF, Biocca L, Pongiglione G, Muthurangu V, Khushnood A, Secinaro A, Chinali M, Schubert S, Berger F, Kuehne T. MRI as a tool for non-invasive vascular profiling: a pilot study in patients with aortic coarctation. *Expert Rev Med Devices.* 2016;13:103–112.

55. Goubergrits L, Riesenkampff E, Yevtushenko P, Schaller J, Kertzsch U, Berger F, Kuehne T. Is MRI-based CFD able to improve clinical treatment of coarctations of aorta? *Ann Biomed Eng.* 2015;43:168–176.
56. Guyatt GH, Sackett DL, Sinclair JC, Hayward R, Cook DJ, Cook RJ. Users' guides to the medical literature. IX. A method for grading health care recommendations. Evidence-Based Medicine Working Group. *JAMA.* 1995;274:1800–1804.
57. McCrindle BW, Jones TK, Morrow WR, Hagler DJ, Lloyd TR, Nouri S, Latson LA. Acute results of balloon angioplasty of native coarctation versus recurrent aortic obstruction are equivalent. Valvuloplasty and Angioplasty of Congenital Anomalies (VACA) Registry Investigators. *J Am Coll Cardiol.* 1996;28:1810–1817.
58. Vandembroucke JP. Observational research, randomised trials, and two views of medical science. *PLoS Med.* 2008;5:e67.
59. Grimes DA, Schulz KF. Bias and causal associations in observational research. *The Lancet.* 2002;359:248–252.
60. Ioannidis JPA, Patsopoulos NA, Rothstein HR. Reasons or excuses for avoiding meta-analysis in forest plots. *BMJ.* 2008;336:1413–1415.

Figures

Figure 1: PRISMA flow chart of study inclusion and exclusion





Figure 2: Network of evidence



The nodes show different treatment strategies (i.e., stenting, balloon dilatation, and surgery).

The lines connecting the nodes indicate the few studies that directly compared two interventions to each other. The size of the nodes is proportional to the number of patients that received a particular treatment.

Panel A: network of evidence for primary analysis. Panel B: network of evidence for secondary analysis.

Figure 3: Effect of patient baseline characteristics on immediate treatment success




Figure shows the proportion of patients achieving blood pressure gradient ≤ 20 mm Hg in single-arm studies (orange bubbles: stenting; blue bubbles: balloon dilatation). Each bubble represents one study, with bubble size representing study sample size.

Panel A: Studies are ranked by ascending peak pre-treatment blood pressure gradient.

Panel B: Studies ranked by ascending proportion of patients with native aortic coarctation.

Figure 4: Results from single-arm studies





Figure shows pooled estimates and 95% CI of the proportion of patients with any given outcome for balloon dilatation and stenting studies obtained from primary (full circles) and secondary (empty circles) analysis.

Figure 5: Results from comparative studies



OR and 95% CI for comparative studies showing odds of patients with any given outcome in balloon dilatation vs. stenting study arms. Full diamonds represent estimates obtained from primary analysis, and empty diamonds estimates from secondary analysis.

Tables

Table 1: PICOS table

P I C O S	
Patient population	<ul style="list-style-type: none">• Patients with native or recurrent coarctation of the aorta ≥ 10 years of age (primary analysis)• Patients with native or recurrent coarctation of the aorta ≥ 1 month of age (secondary analysis)
Interventions	<ul style="list-style-type: none">• Balloon dilatation• Stenting
Comparators	<ul style="list-style-type: none">• Any comparator
Outcomes	<ul style="list-style-type: none">• Proportion of patients with post-treatment blood pressure gradient ≤ 20mm Hg• Proportion of patients with post-treatment blood pressure gradient ≤ 10mm Hg• Proportion of patients alive 30 days after treatment• Proportion of patients with severe complications during intervention or before discharge• Proportion of patients with minor complications during intervention or before discharge• Proportion of patients with reinterventions at follow-up• Proportion of patients with recurrent coarctation at follow-up• Proportion of patients with aortic wall injuries at follow-up
Study designs	<ul style="list-style-type: none">• Any study design

Table 2: Patient baseline characteristics in all included studies

CoA: coarctation of the aorta; AVD: aortic valve disease; VSD: ventricular septal defect;

PDA: patent ductus arteriosus

*** significant at the 1% level; ** 5% level; * 10% level

	Balloon dilatation		Stenting		<i>t-test for statistical difference</i>
	<i>Studies reporting variable [participants]</i>		<i>Studies reporting variable [participants]</i>		<i>p-value</i>
Patients overall	361		423		
Mean age (years)	26.65	12 [361]	28.82	15 [423]	0.355
Mean weight (kg)	51.80	1 [15]	59.37	5 [163]	0.480
Children patients	1.7%	8 [181]	1.6%	8 [173]	0.930
Adult patients	97.7%	6 [134]	97.9%	6 [130]	0.930
Gradient pre-treatment (mm Hg)	59.47	12 [361]	45.41	13 [339]	0.013**
Patients with native CoA	98.6%	11 [338]	75.6%	14 [364]	0.012**
Patients with recurrent CoA	1.4%	11 [338]	24.4%	14 [364]	0.012**
Patients receiving antihypertensive medication	60.7%	9 [234]	77.7%	9 [198]	0.103
Concomitant heart defects					
Patients with isolated CoA	64.2%	4 [140]	54.7%	3 [60]	0.579
Patients with AVD	37.2%	9 [271]	44.9%	10 [256]	0.616
Patients with VSD	8.7%	7 [225]	7.3%	6 [157]	0.734
Patients with PDA	4.2%	4 [90]	11.3%	4 [117]	0.177
Patients with other concomitant genetic heart defects	3.8%	6 [218]	20.9%	8 [216]	0.093*