

Shanna Marrinan, [Andres Roman-Urrestarazu](#), Declan Naughton, Ermelinda Levari, John Collins, Robert Chilcott, Giuseppe Bersani, Ornella Corazza

Hair analysis for the detection of drug use – is there potential for evasion?

**Article (Accepted version)
(Refereed)**

Original citation:

Marrinan, Shanna, Roman-Urrestarazu, Andres, Naughton, Declan, Levari, Ermelinda, Collins, John, Chilcott, Robert, Bersani, Giuseppe and Corazza, Ornella *Hair analysis for the detection of drug use – is there potential for evasion?* Human Psychopharmacology: Clinical and Experimental . ISSN 1099-1077
DOI: [10.1002/hup.2587](https://doi.org/10.1002/hup.2587)

© 2017 John Wiley & Sons, Inc.

This version available at: <http://eprints.lse.ac.uk/69774/>
Available in LSE Research Online: March 2017

LSE has developed LSE Research Online so that users may access research output of the School. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LSE Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain. You may freely distribute the URL (<http://eprints.lse.ac.uk>) of the LSE Research Online website.

This document is the author's final accepted version of the journal article. There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

Hair analysis for the detection of drug use – is there potential for evasion?

Shanna Marrinan, Andres Roman-Urrestarazu, Declan Naughton, Ermelinda Levari, John Collins, Robert Chilcott, Giuseppe Bersani, Ornella Corazza

Shanna Marrinan¹, Andres Roman-Urrestarazu², Declan Naughton³, Emerlinda Levari⁴, John Collins⁵, Robert Chilcott⁶, Giuseppe Bersani⁷, Ornella Corazza^{5,7}

¹ Dept of Science, Technology, Engineering & Public Policy (STeAPP) University College London, UK

² Institute of Public Health, Cambridge University, UK

³ School of Life Sciences, Pharmacy and Chemistry, Kingston University, London, UK

⁴ Addiction Treatment Unit (SerD), Trento, Italy

⁵ London School of Economics IDEAS International Drug Policy Project, London, UK

⁶ Department of Pharmacy, Pharmacology and Postgraduate Medicine, School of Life and Medical Sciences, University of Hertfordshire, UK

⁷ Sapienza University of Rome, Italy

Correspondence to:

Dr Ornella Corazza
Reader in Substance Addictions and Behaviours
School of Life and Medical Science,
University of Hertfordshire,
College Lane Campus,
Hatfield,
Herts, AL10 9AB
E-mail: o.corazza@herts.ac.uk

ABSTRACT

Background: Hair analysis for illicit substances is widely used to detect chronic drug consumption or abstinence from drugs. Testees are increasingly seeking ways to avoid detection by using a variety of untested adulterant products (e.g. shampoos, cleansers) widely sold online. This study aims to investigate adulteration of hair samples and to assess effectiveness of such methods.

Methods: The literature on hair test evasion was searched for on PubMed/MEDLINE, Psycinfo, and Google Scholar. Given the sparse nature of peer-reviewed data on this subject, results were integrated with a qualitative assessment of online sources, including user-orientated information/commercial websites, drug fora and ‘chat rooms’. Over four million web sources were identified in a Google search by using ‘beat hair drug test’ and the first 86 were monitored on regular basis and considered for further analysis.

Results: Attempts to influence hair test results are widespread. Various ‘shampoos’, ‘cleansers’ among other products, were found for sale, which claim to remove analytes. Often advertised with aggressive marketing strategies, which include discounts, testimonials and unsupported claims of efficacy. However, these products may pose serious health hazards and are also potentially toxic. **In addition, many anecdotal reports suggest that Novel Psychoactive Substances (NPS) are also consumed as an evasion technique**, as these are not easily detectable via standard drug test. Recent changes on NPS legislations such as New Psychoactive Bill in the UK might further challenge the testing process.

Conclusion: Further research is needed by way of chemical analysis and trial of the adulterant products sold online and their effects as well as the development of more sophisticated hair testing techniques.

Key Words: Hair test, hair analysis, drugs, NPS, evasion; Psychoactive Substances Bill

Introduction

Drug testing, including hair analysis, is increasingly utilized in some employment sectors, especially where there are safety concerns around drug use (Karch 2016; Ironmonger 2014; UIC 2008). **There is weak evidence however**, linking drug testing to reducing employee drug use rates, or accident and injury incidence (Pidd & Roche 2014). After ingestion, traces of substances and their metabolites are deposited into hair via the bloodstream and into the growing follicle. A 3cm sample of hair (usually taken from the scalp) is sufficient to test for drug use in the preceding three months, **accepting that variations in growth rate of between 0.7 and 3.6 cm/month are possible (Schultz et al. 1993)**. Despite the existence of very reliable assays, **such as those based on MS techniques, most immunoassay kits are designed to test chronic drug use, meaning a single incident of drug use may escape detection**. Very recent drug use is also usually missed as affected hair has not had sufficient time to grow out of the scalp (Pragst & Balikova 2006). Studies on the appearance of drugs in beard hair reveal variations in incorporation rates with **amphetamines and cocaine appearing within one day of intake whilst morphine was detectable after one week (Wennig, 2000)**. A large number of parameters can affect the appearance of drugs and their metabolites in hair including (i) amount of sweating, (ii) size and structure of the parent drug, (iii) protein binding in blood, and (iv) polarity and charge of the drug (Wennig, 2000). A further consideration is hair colour, with several studies revealing **higher** levels of drug uptake into darker hair (Cone et al. 1993; Joseph et al. 1996; Gygi et al. 1996).

In forensic and clinical toxicology, hair testing is typically used in preference to urine or blood tests, as the former may provide additional information such as indications as to the extent and historical pattern of drug usage (Klein et al. 2000; Friguis et al. 2012). As the use of novel psychoactive substances (NPS) becomes widespread across the globe (Corazza 2013a; Papaseit 2014), developments in the potential of hair testing techniques to detect a wide range of NPS using various chromatographic techniques have gained attention (Smith 2015). Recent advances in analytical methods have improved both hardware and software capabilities, lowering detection limits along with the possibility of increasing the numbers of compounds analysed in a single chromatographic run. The approach of initial screening followed by quantitative analysis of libraries of drugs and their metabolites has opened new horizons for drug detection in hair (Shah et al. 2014). A key feature is the ability to add any newly discovered drug to the method in a few simple steps, thus expanding the library of compounds that can be detected in a 10-minute analytical run. Currently however, the full potential of hair testing is not being routinely used as most immunoassays used are ‘unsuitable for the detection of these compounds’ (Gottardo et al. 2014).

Evasion is a common issue on other types of drug test. Urine samples, which are often provided in private by the testee, can be easily adulterated, substituted or diluted (Beard, 2016; Jaffee, Trucco, Levy & Weiss 2007; Jones et al. 2015; Cody & Valtier 2001; Dasgupta 2007). Blood tests are generally considered tamper-proof (due to the sample being obtained by a professional), but samples must be acquired within much shorter timeframes than urine or hair, especially for drugs with exhibit a short plasma half-life. Testing methods with shorter detection periods see some users temporarily suspending drug use in order to pass the test. They are also considered by many to be highly invasive, and blood samples require specialist extraction and transportation (Koster et al. 2014; Friguis et al. 2012). Hair analysis circumvents many of the issues around detection windows, collection procedures and tampering potential (Deshmukh et al. 2010; Accornero et al. 2002). Recently, however, a dramatic increase in websites selling products to avoid detection have been recorded, supported by a growing literature, suggesting that large numbers of testees are attempting to evade detection (Hill et al. 2014; Polla et al. 2009; Gerada & Gilvarry 2005; Dolan et al. 2004). This study explores the evasion methods discussed in the literature and in Internet fora, discussion threads, online shops and other types of web content. Particular attention is given to the use cleansing shampoos, and NPS, which in some instances, are used in the belief that these will not be tested for in a hair analysis.

Methods

The literature on evasion and adulteration of hair testing was initially searched in PsycInfo and PubMed databases. Considering the sparse nature of peer-reviewed publications on this topic, results were integrated with a qualitative assessment of relevant websites and online fora. Over a three-month period, between February and April 2016, exploratory qualitative searches were carried out in English and Italian. Generic and specific keywords were used, such as ‘pass’/evade/avoid detection/cheat/beat/tamper’ ‘hair drugs/substance test’ along with names of specific substances. The first 86 web results were used for analysis, as the search reached a saturation point, and these were monitored on a regular weekly basis. Any personal data collected from online fora were removed for anonymity. Permission for the study was granted by the School of Pharmacy Ethics Committee, Hatfield, United Kingdom (November 2013 PHAEC/10-42). This study assesses the nature of the products sold claiming to remove drug traces from the hair, the marketing strategies employed by sellers, and the potential side effects and efficacy as reported by users.

Results

A limited amount of academic literature into the deliberate adulteration of hair samples was found (Hill et al. 2014; Polla et al. 2009; Gerada & Gilvarry 2005). Other studies have addressed how bleaching and other cosmetic treatments can affect the quality of hair samples, and the degree to which analytes can be detected, although this might not be done deliberately (ASAM 2013; Martins et al. 2008; Yegles 2005). Most of the emerging evidence on circumvention focused on other specimen types, such as urine and blood (Jaffee, Trucco, Levy & Weiss 2007; Jones et al. 2015; Cody & Valtier 2001; Dasgupta 2007). The range and sophistication of some of those evasion methods (including even the use of a fake penis which delivers warm artificial urine) lends support to the assumption that substance users will seek to evade detection on the hair analysis also.

There is an abundance of journalistic and anecdotal evidence that users are attempting to evade detection (e.g. Featherstone, 2015; Brown 2014; Hill et al. 2014; Wales Online article, 2013). In searching for qualitative user data on this topic, a simple Google search with fewer more colloquial terms, was found to be more effective. The search term ‘beat drug hair test’ for example produced 3,660,000 results at the start of the research period, and by the end (three months after), was producing more than four million, suggesting increasing interest in this area. Most of these appeared to relate to user content (fora, blogs) or evasion-aiding seller sites. Techniques to increase the likelihood of passing the test discussed online include abstaining for a sufficient period, shaving the hair off, using commercially available products (**shampoos, hair cleansers**) or home remedies to ‘strip’ away evidence of drug use, switching to a lesser tested-for substance, and offering excuses for any positive detection (perhaps prompting a later-date retest).

Shampoos and hair cleansers

Whilst academic literature on deliberate adulteration products is limited, a number of papers attest to the idea that certain cosmetic treatments, such as bleaching, colouring, chemically straightening hair, can affect the speed at which drug analytes leave the hair sample (Hill et al. 2014; Hill 2016). Jurado et al. 2007 discovered that ‘in all cases, the drug content in hair that had undergone treatment decreased in comparison with untreated hair’ with mean differences of between approximately 40-60% depending on the substance, type of treatment, and the extent of hair damage. This has been further supported by the Society of Hair testing guidelines for drug testing in hair (Cooper, Kronstrand & Kintz 2012) and the American Society of Addiction Medicine (2013).

Based on the evidence of drug analytes leaching out of hair following some types of treatment, a myriad of commercial products has emerged which claim to strip metabolites from the hair. These are often branded and sold online in brightly coloured boxes as “hair cleansers”, “cleansing systems”, or

“hair detoxifiers” with different names, such as Nexus Aloe Rid, Clear Choice, Two Steps Ahead, Synergy Detox, Supreme Kleen, Test Pass Detox Shampoo, Toxin Wash, Zydol Ultraclean, among others. Many of the sites selling these products utilise misleading marketing strategies by offering apparently neutral and factual information and advice, designed to lend a sense of honesty and trustworthiness. Scientific language is often used **on these sites** to describe how drug metabolites enter the hair shaft, and how these are then tested for as well as medical terminology and specific claims to reinforce the assumption that the products have been clinically researched (DrugRehabAlliance.com; Theweedblog.com, accessed April 2016). Skeptics are reassured with claims such as: ‘500,000 hair tests passed since 1998’ (PassYourTest.com), or with a ‘Talk to a drug test expert’ button to allow for live chat with an expert. Links to the vendors’ Facebook page, Twitter, YouTube, Google+ and even RSS feed, are available to connect to recognizable, trusted brands, and to sustain contact with potential buyers, and even include a money-back guarantee (e.g. Theweedblog.com). Special offers are also common with further unit price reductions in case of multiple buys (e.g. PassYourTest.com).

A small number of sites genuinely appear to be nothing more than information brokers – they do not sell or endorse any commercial product, stating they have not been proven to be effective in clinical trials and they provide home remedies, involving vinegar, salicylic acne treatment, fabric detergent, and hair dye (e.g. hairfollicletest.info; WikiHow). Video online tutorials for these procedures are also available on the YouTube (Youtube 2015) with mix of positives, negatives, and general questions from users/potential users.

The price of these products is very high, ranging from £50 - £827 for a bottle of hair cleansing shampoo and conditioner. Amazon is a particularly interesting vendor of these products, as the reviews left cannot be altered by the seller. For instance, Zydol Ultra Clean has 10 reviews (with an average of 2.5 stars but with only three 5-star and six 1-star reviews – suggesting on balance it hasn’t been successful for these users). The US version of the site has a larger range of hair products aimed at the drug test market, with far larger numbers of reviews, but they too hover around the half-pass mark at best. Variable results were reported on forum discussions (marijuana.com, Reddit, 420 magazine, accessed April 2016), but on independent selling sites, no negative comments or reports of failure were found, suggesting that comments may be moderated, or contain falsified ‘testimonies’ of efficacy.

A selection of user feedback regarding these products is illustrated in Table 1.

(Table 1)

It is problematic to establish effectiveness from user feedback alone, as some drug use may go undetected without any particular method having been employed (Hill et al. 2014). While hair testing demonstrates very high specificity, its sensitivity can be variable (ASAM 2013). Recent studies have reported that Gas Chromatography/Mass Spectrometry techniques detected between 34.4% and 77% of moderate to heavy Cannabis users with lower rates for some other substances (Taylor et al. 2016; Gycznski et al. 2014; Junkuy 2014). In contrast, Musshoff et al. (2012) compared two types of immunoassay procedures for drug testing in hair samples and found much higher sensitivities (65-99% depending on substance tested and method used). The authors advocate for lower cut-off values in order to capture positive results, recommendations for which are set by the European Workplace Drug Testing Society. However, lowering cut off points could raise other potential problems, such as contamination. Confirmatory testing should also be provided for any positive results obtained from screening. The wide variety of hair tests available, and consequent variations in terms of performance, also contribute to these differences (Vincenti et al. 2014; Baciú et al. 2015), although an internationally-recognised standard for accredited methods does exist - ISO/IEC 17025 (Nutt et al. 2014), and the EWDTS have published a common framework to guide those involved in the testing procedure (EWDTS 2015).

This variation in sensitivity means many users will receive an unexpectedly negative result (falling below the minimum cut-off level), inadvertently supporting the claims of any commercial product they may have been used. Much of the ‘success’ therefore, in the claims of these product vendors, could be attributed to these inevitable false negatives.

NPS consumption to avoid drug detection: a growing trend?

Substance substitution - the swapping of a commonly tested-for substance for one less likely to be detected, such as NPS - is another evasion strategy method widely identified in the course of this research (e.g. Topix.com; Drugs Forum; High Existence). NPS have posed significant challenges to policy-makers due to the rapid pace at which chemical structures are altered in response to legislation to control individual substances, creating huge numbers of new NPS (Griffiths, Brown & Sedefov 2013). The introduction of the Psychoactive Substances Bill to the UK in May 2016, however, made any substance that has an effect on normal brain functioning (other than tobacco, alcohol and caffeine) automatically banned, a move which has attracted criticism from scientists and experts due to the inherent difficulties in defining and controlling a limitless number of substances (Reuter & Pardo 2016). A number of studies indicate that NPS are attractive to users regardless of their ‘legal’ status, because they are less often tested for in toxicology (Soussan & Kjellgren 2016; Brown 2015; Gunderson et al. 2014; ASAM 2013; Simmons et al. 2011).

However, it must be noted that hair analysis has the potential to detect any substance, depending on the analytes it has been ‘programmed’ to recognize (Smith et al. 2015; Shah et al. 2014; **Fisichella et al. 2014**). Indeed, a number of hair analysis kit sellers were identified online who claim to sell tests capable of detecting a large number of substances including one who say they can test for over 2000 (e.g. DNA Legal). Research into the practical potential for quantifying NPS in drug testing is limited (Ambach et al. 2015). However, it appears that generally speaking, commercially-available hair tests are only capable of detecting a small selection of the most commonly abused drugs, and employers are often requesting tests for a limited number of better known substances of abuse, leaving room for NPS to go undetected (Gottardo et al. 2014).

Many NPS have effects equal to or greater than traditional drugs, and far less is known about them. However, for some they produce a false ‘sense of security’ largely generated by the marketing strategies of online sellers, which make unsupported claims regarding their safety, efficacy and even health benefits (Corazza et al. 2013b).

As the use of NPS becomes more widespread, the challenges around detection and monitoring of use become more pertinent. Although information is available about individual brands of test and the analytes they are capable of detecting. It is currently not clear to what extent overall current methods are able to detect the use of NPS. For example, there is currently no information available on the market share of each brand of drug test, nor what percentage of currently available NPS may be detected. Indeed, there is no exhaustive list of all NPS; some may trigger a positive result because they have previously been included in the chromatography library of detectable substances. Other NPS might be detectable as they have a similar molecular structure to existing illicit drugs. However, a proportion of NPS are either unknown at this time, **not routinely included, and sufficient** different to other drugs to be currently undetectable.

Numerous conversation threads were discovered where users discussed their use of NPS in order to avoid detection (Topix.com; Drugs Forum; High Existence), **although there** is a high degree of uncertainty within the user community on the detectability of NPS. For example, while synthetic cannabinoids will not react to traditional THC immunoassays (Smith et al. 2015), many users have reported testing positive for traditional substances, possibly as a result of their (often) chemical similarity, but also possibly as a result of substance mislabeling; Smith et al. (2015) and Brandt et al. (2010) discuss the documented presence of controlled psychoactive substances in the so-called ‘legal

highs' being sold before the recent blanket ban. The opposite has also been encountered – A study of 679 nightclub and festival goers by Palamar et al. (2016) detected high levels of NPS via hair analysis in those self-reporting MDMA use but with no lifetime use of NPS, suggesting the contamination goes both ways. Poly drug use is widespread, with many users mixing NPS and more traditional illicit drugs, by choice or by supplier adulteration. Researchers monitoring the presence of NPS in samples previously submitted for drug testing have recorded high prevalence – up to 37% - with rapid growth in recent years (Hondebrink et al. 2015; Salomone et al. 2013; Rust et al. 2012).

Other potential methods for evading drug hair testing

Many of the sites, including those offering commercial preparations, will begin by advising users to consider abstinence from drugs, or shaving their hair in the first instance (e.g. HairFollicleDrugTest.com; hairfollicledrugtest.info, wikihow; AlwaysTestClean.com - accessed April 2016). Cropping hair closely (and shaving body hair) prevents a sufficient sample from being collected for hair testing, which may therefore be substituted by a shorter detection window test, such as those using saliva, blood or urine. Occasional mention was made of whole body cleaners ('consumables'), ingested by mouth, which remove drug traces and metabolites from hair, urine and blood. An example is "Nutracleanse" offered by PassYourTest.com with a full guarantee, while "Total Detoxification System", sold by TestingClean.com, also prescribes a set diet for users to follow in combination with the treatment. No **plausible testimonies** were identified from users achieving a clear test result via the use of consumables only.

A number of information sites (e.g. UKCIA) offer possible 'excuses' that users could give in the event of a positive result. This opens up potential for those with a deservedly positive result to call for a retest. AlwaysStayClean.com for example, makes the dubious claim that 'about 5%' of tests give a false positive and list reasons including the use of prescriptions and over-the-counter medications as well as poppy seeds in bagels. However, studies administering poppy seeds to testees appears have shown that a false positive result was 'highly unlikely' (Hill et al. 2005). Controversially, the ethnicity of testees has also **been offered as** a reason for false positive tests (for example, AlwaysStayClean.com). Kidwell, Smith & Shepherd (2015) demonstrated that some ethnic hair care products can significantly exacerbate the absorption of environmental contaminants and earlier studies measured higher levels of drug uptake into darker hair due to **the higher presence** of melanin (Cone et al. 1993; Joseph et al. 1996; Gygi et al. 1996). However, a comprehensive laboratory analysis failed to find any statistical correlation between ethnicity and incidence of false-positive drug tests (Kelly et al. 2000).

Study limitations

Internet sources: One could wonder about the limitations of carrying out an analysis of online information, **and indeed it** may be inappropriate to trust data obtained from the Internet without some means of independent verification. However, in the absence of relevant peer-reviewed data, this seems to be the only valid method to obtain preliminary information about new and emerging phenomena.

Query: a number of set key words were used for the searches and these did not include street names used to sell the adulterant products.

Search: Similarly, searches were performed with a wide range of search terms, but the language surrounding the use of these substances is often very colloquial and regionally dependent. With more time and resources, interesting directions to continue this research would be to examine the content of the products sold purporting to help secure a negative result, and to assess the extent to which the most widely used hair tests are able to detect NPS.

Conclusions

Overall, the attempted evasion of substance hair testing is a large, rapidly growing and lucrative market. Various ‘shampoos’, ‘cleansers’ among other products, were found for sale, which claim to remove analytes. Evidence surrounding the effectiveness of commercial products sold for this purpose is mixed, but when the quality of sources is considered, generally disputes efficacy. Clearly, there are high numbers of individuals self-reporting success in evading detection on hair tests. Minimum recommended cut-off levels set by the European Workplace Drug Testing Society might not be sufficient to detect light or infrequent drug use and outcomes might thus be interpreted by testees as a false confirmation of a successful evasion attempt. However, a large number of individuals describe their drug use as heavy and/or frequent, and this body of anecdotal evidence, combined with research evidence around the dissipation of drug metabolites from treatment-damaged hair samples, means that these products warrant further investigation. Another important area in need of attention that emerged from this study was the use of NPS to substitute illicit drug use for the purposes of evading detection, especially as it currently unclear which NPS can be detected and by what proportion of available tests. As discussed, the recent introduction of the Psychoactive Substances Bill in the UK and similar legislations further challenge the testing process. Novel testing solutions will need to be found in order to face the unknown and growing number of NPS available in the illicit drug markets as well as a large trial of the adulterant products sold online and their effects.

References

- Accornero VH, Morrow CE, Bandstra ES, Johnson AL, Anthony JC. ‘Behavioral outcome of preschoolers exposed prenatally to cocaine: role of maternal behavioral health’ *J Pediatr Psychol.* 2002;27:259-69. **PMCID: PMC2760334**
- American Society of Addiction Medicine (ASAM, 2013) – *a White Paper ‘Drug testing’* Oct 26.
- Ambach, L. Hernández Redondo, A. König, S. Angerer, V. Schü, S. Weinmann, WW. (2015) ‘Detection and quantification of 56 new psychoactive substances in whole blood and urine by LC–MS/MS’ in *Bioanalysis* May 2015 ,Vol. 7, No. 9, Pages 1119-1136 doi: 10.4155/bio.15.48.
- Baciu, T. Borrull, F. Aguilar, C. Calull, M. (2015) ‘Recent trends in analytical methods and separation techniques for drugs of abuse in hair’ *Analytica Chimica Acta.* Volume 856, 26 January 2015, Pages 1–26 **DOI: 10.1016/j.aca.2014.06.051**
- Beard, CL. (2016) *The Effectiveness of the Alere iScreen® Urine Adulteration Test Strip at Detecting Six Common Urine Adulterants* - Honor Thesis, East Carolina University, <http://thescholarship.ecu.edu/handle/10342/5585>
- Cody, JT. Valtier, S. (2001) ‘Effects of Stealth adulterant on immunoassay testing for drugs of abuse’. In *J Anal Toxicol* 2001 Sep; 25(6):466-70 **PMID:11550822**
- Cone, E.J. Darwin, W.D. Wang, W.L. (1993) ‘The occurrence of cocaine, heroin and metabolites in hair of drug abusers’. *Forensic Science International* 63 (1-3), 55-68. **PMID:8138234**
- Cooper, G. Kronstrand R. Kintz, P (2012) ‘Society of Hair testing guidelines for drug testing in hair’ in *Forensic Science International* Volume 218, issues 1-3 10 May 2012 pages 20-24 doi: **10.1016/j.forsciint.2011.10.024**
- Corazza, O., Demetrovics, Z., van den Brink, W., Schifano, F. (2013a) ‘Legal highs’ an inappropriate term for ‘Novel Psychoactive Drugs’ in drug prevention and scientific debate’, *International Journal of Drug Policy*, 24 (1): 82-3. **DOI:10.1016/j.drugpo.2012.06.005**
- Corazza, O.,** Assi, S., Simonato P, Corkery J, Bersani FS, Demetrovics Z, Stair J, Fergus S, Pezzolesi C, Pasinetti M, Deluca P, Drummond C, Davey Z, Blaszkowicz U, Moskalewicz J, Mervo B, Furia LD,

Farre M, Flesland L, Pisarska A, Shapiro H, Siemann H, Skutle A, Sferrazza E, Torrens M, Sambola F, van der Kreeft P, Scherbaum N, Schifano (2013b). 'Promoting Innovation and excellence to face the rapid diffusion of Novel Psychoactive Substances in the EU: the outcomes of the ReDNet project'. *Human Psychopharmacology: Clinical and Experimental. Special issue*, Jul, 28(4): 317-23. doi: 10.1002/hup.2299.

Dasgupta, A. (2007) 'The effects of adulterants and selected ingested compounds on drug-of-abuse testing in urine' in *Am J Clin Pathol* 2007;128(3): 491-503 DOI:[10.1309/FQY06F8XKTQPM149](https://doi.org/10.1309/FQY06F8XKTQPM149)

Deshmukha, N. Hussaina, I. Barkera, J. Petroczib, A. Naughton, D. (2010) 'Analysis of anabolic steroids in human hair using LC-MS/MS' in *Steroids* Volume 75, Issue 10, October 2010, Pages 710–714 doi: [10.1016/j.steroids.2010.04.007](https://doi.org/10.1016/j.steroids.2010.04.007)

Dolan, K. Rouen², D. Kimber², J (2004) 'An overview of the use of urine, hair, sweat and saliva to detect drug use' in *Drug and Alcohol Review*, Volume 23, Issue 2, pages 213–217, June 2004 DOI: [10.1080/09595230410001704208](https://doi.org/10.1080/09595230410001704208)

European Workplace Drug Testing Society European Guidelines for Workplace Drug and Alcohol Testing in Hair Nov 2015 Version 2 <http://www.ewdts.org/data/uploads/documents/ewdts-guideline-hair-v2.0.pdf>

Friguls, B. Joya, X. Garcia-Serra, J. Gómez-Culebras, M. Pichini, S. Martinez, S. Oriol Vall, O. Garcia-Algar, O. (2012) 'Assessment of exposure to drugs of abuse during pregnancy by hair analysis in a Mediterranean island' in *Addiction* Volume 107, Issue 8, pages 1471–1479, August 2012 doi: [10.1111/j.1360-0443.2012.03828.x](https://doi.org/10.1111/j.1360-0443.2012.03828.x)

[Fisichella M](#), [Morini L](#), [Sempio C](#), [Groppi A](#). (2014) 'Validation of a multi-analyte LC-MS/MS method for screening and quantification of 87 psychoactive drugs and their metabolites in hair' *Anal Bioanal Chem*. 2014 May;406(14):3497-506. doi: 10.1007/s00216-014-7763-2. Epub 2014 Mar 30. doi: [10.1007/s00216-014-7763-2](https://doi.org/10.1007/s00216-014-7763-2).

Gerada, C. & Gilvarry, E. (2005) 'Random drug testing in schools' *BRJ Gen Pract* v.55(516) July 2005 PMID: [PMC1472793](https://pubmed.ncbi.nlm.nih.gov/1472793/)

Gottardo, R. Sorio, D. Musile, G. Trapani, E. Seri, C. Serpelloni, G. Tagliaro, F (2014) 'Screening for synthetic cannabinoids in hair by using LC-QTOF MS: A new and powerful approach to study the penetration of these new psychoactive substances in the population' *Med Sci Law* January 2014 vol. 54 no. 1 22-27 doi: [10.1177/0025802413477396](https://doi.org/10.1177/0025802413477396)

Griffiths, P. Evans-Brown, M. Sedefov, R (2013) 'Getting up to speed with the public health and regulatory challenges posed by new psychoactive substances in the information age' in *Addiction*, Vol 108, issue 10, pages 1700-1703 oct 2013 Doi: [10.1111/add.12287](https://doi.org/10.1111/add.12287)

Gryczynski, J. Schwartz, R. Mitchell, S. O'Grady, K. Ondersma, S. (2014) 'Hair drug testing results and self-reported drug use among primary care patients with moderate-risk illicit drug use'. In *Drug and Alcohol Dependence*, 1st Aug 2014, vol. 141 pp. 44-50 doi: [10.1016/j.drugalcdep.2014.05.001](https://doi.org/10.1016/j.drugalcdep.2014.05.001)

Gygi, S.P. Joseph, R.E. Cone, E.J. Wilkins, D.G. Rollins, D.E. (1996) Incorporation of codeine and metabolites into hair: Role of pigmentation. *Drug Metabolism and Disposition* Volume 24, (4) Pages 495-501. PMID:[8801066](https://pubmed.ncbi.nlm.nih.gov/8801066/)

Hill, V. Schaffer, M.I. Loni, E. Stowe, G.N. (2016) *Integrity testing of hair samples* Patent US 9261522 B2

Hill, V., Loni, E., Cairns, T., Sommer, J. and Schaffer, M. (2014) 'Identification and analysis of damaged or porous hair' *Drug Test Analysis* 6: 42–54. doi: [10.1002/dta.1652](https://doi.org/10.1002/dta.1652).

Hill, V. (2005) 'Multiple Aspects of Hair Analysis for Opiates' - *Journal of Analytical Toxicology* Oct;**29(7):696-703**. PMID:16419403

Hondebrink, L.I. Nugteren-van Lonkhuyzen, JJ2. Van Der Gouwe, D3. Brunt, TM4. 'Monitoring new psychoactive substances (NPS) in The Netherlands: data from the drug market and the Poisons Information Centre.' In *Drug Alcohol Depend.* 2015 Feb 1;147:109-15. doi: 10.1016/j.drugalcdep.2014.11.033. Epub 2014 Dec 13. DOI:[10.1016/j.drugalcdep.2014.11.033](https://doi.org/10.1016/j.drugalcdep.2014.11.033)

Jaffee, WB. Trucco, E. Levy S. & Weiss, RD. (2007) 'Is this urine really negative? A systematic review of tampering methods in urine drug screening and testing'. In *J Subst Abuse* 2007 Jul;33(1):33-42 DOI:[10.1016/j.jsat.2006.11.008](https://doi.org/10.1016/j.jsat.2006.11.008)

Jones, JD. Atchinson, JJ. Madera, G. Metz VE. Comer, SD. (2015) Need and Utility of a polyethylene glycol marker to ensure against urine falsification among heroin users in *Drug Alcohol Depend* 2015 Aug 1; 153:201-6 DOI:[10.1016/j.drugalcdep.2015.05.021](https://doi.org/10.1016/j.drugalcdep.2015.05.021)

Joseph, R.E. Su, T.P. Cone, E.J. (1996) In vitro binding studies of drugs to hair: influence of melanin and lipids on cocaine binding to Caucasoid and Africoid hair. *Journal of Analytical Toxicology* Volume 20, Pages 338-344. PMID:8889667

[Junkuy, A.](#) Aramrattana, A. Sribanditmongkol, P. (2014) 'A comparative study of self-report, urinalysis and hair analysis in the detection of methamphetamine in Yaba users' in *Journal of the Medical Association of Thailand Thangphaet* 97(7):776-784 PMID:25265778

Jurado C1, Kintz P, Menéndez M, Repetto M. (1997) 'Influence of the cosmetic treatment of hair on drug testing' in *Int J Legal Med.* 1997;110(3):159-63. PMID: 9228567

Karch, SB (2016) *Workplace Drugs Testing*, CRC Press.

Kelly RC1, Mieczkowski T, Sweeney SA, Bourland JA. (2000) 'Hair analysis for drugs of abuse. Hair color and race differentials or systematic differences in drug preferences?' *Forensic Sci Int.* 2000 Jan 10;107(1-3):63-86. PMID:10689562

Kidwell, D. Smith, F. Shepherd, P (2015) 'Ethnic hair care products may increase false positives in hair drug testing' in *Forensic science international*, Volume 257, December 2015, Pages 160–164 DOI:[10.1016/j.forsciint.2015.07.023](https://doi.org/10.1016/j.forsciint.2015.07.023)

Klein J, Karaskov T, Koren G. (2000) 'Clinical applications of hair testing for drugs of abuse--the Canadian experience' in *Forensic Sci Int*;107:281-8. PMID: 10689580

Koster, R. Alffenaar, J. Greijdanus, B. VanDerNagel, J. Uges, D. (2014) 'Fast and Highly Selective LC-MS/MS Screening for THC and 16 Other Abused Drugs and Metabolites in Human Hair to Monitor Patients for Drug Abuse' in *Therapeutic Drug Monitoring* April 2014 - Volume 36 - Issue 2 - p 234–243 Doi: [10.1097/FTD.0b013e3182a377e8](https://doi.org/10.1097/FTD.0b013e3182a377e8)

Martins, L. Yegles, M. Thieme, D. Wennig, R. (2008) 'Influence of bleaching on the enantiomeric disposition of amphetamine-type stimulants in hair'. *Forensic Sci. Int.* 2008, 176, 38. DOI: [10.1016/j.forsciint.2007.06.023](https://doi.org/10.1016/j.forsciint.2007.06.023)

Musshoff, F. Kirschbaum, K. M. Graumann, K. Herzfeld, C. Sachs, H. Madea, B. (2012) 'Evaluation of two immunoassay procedures for drug testing in hair samples' Doi: [10.1016/j.forsciint.2011.03.030](https://doi.org/10.1016/j.forsciint.2011.03.030)

Nutt, J. Tsanaclis, L. Bevan, S. Bagley, K. Blackwell, K. Wicks, J. (2014) 'Hair analysis in the workplace: global harmonisation required' paper presented at the *Society of Hair Testing* in Bordeaux, June 2014. DOI:10.13140/RG.2.2.11233.92005

Palamar, JJ. Salomone A. Vincenti M. Cleland CM. (2016) 'Detection of "bath salts" and other novel psychoactive substances in hair samples of ecstasy/MDMA/"Molly" users.' *Drug and Alcohol Dependence* 2016, 161:200-205 DOI:10.1016/j.drugalcdep.2016.02.001

Papaseit, E. Farré, M. Schifano, F. Torrens, M. (2014) 'Emerging drugs in Europe' *Current Opinion in Psychiatry*. July 2014 - Volume 27 - Issue 4 - p 243–250 Doi: 10.1097/YCO.0000000000000071

Pidd, K. Roche, A (2014) 'How effective is drug testing as a workplace safety strategy? A systematic review of the evidence' in *Accident Analysis and Prevention* Oct 2014 vol. 71: 154-165 Doi: 10.1016/j.aap.2014.05.012.

Polla, M. Stramesi, C. Pichini, S. Palmi, I. Vignali, C. Dall'Olio, G. (2009) 'Hair testing is superior to urine to disclose cocaine consumption in driver's licence regranting' in *Forensic Science International* August 10, 2009 Volume 189, Issues 1-3, Pages e41–e43 Doi:http://dx.doi.org/10.1016/j.forsciint.2009.04.020

Pragsta, F. Balikovab, M (2006) 'State of the art in hair analysis for detection of drug and alcohol abuse' in *Clinica Chimica Acta* Volume 370, Issues 1–2, August 2006, Pages 17–49 DOI:10.1016/j.cca.2006.02.019

Reuter, P. Pardo, B. (2016) 'Can new psychoactive substances be regulated effectively? An assessment of the British Psychoactive Substances Bill' in *Addiction* 2016 May 24 doi:10.1111/add.13439

Rust, KY1. Baumgartner, MR. Dally, AM. Kraemer, T. (2012) 'Prevalence of new psychoactive substances: A retrospective study in hair'. *Drug Test Anal.* 2012 Jun;4(6):402-8. doi: 10.1002/dta.1338.

Salamone, A. Luciano, C. Di Corcia, D. Gerace, E. Vincenti, M. (2013) 'Hair analysis as a tool to evaluate the prevalence of synthetic cannabinoids in different populations of drug consumers' in *Drug Testing and Analysis* Volume 6, issues 1-2, pages 126-134, Jan-Feb 2014. DOI: 10.1002/dta.1556

Schütz, H. Ahrens, B. Erdmann, F. Rochholz G. (1993) Nachweis von Arznei- und anderen Fremdstoffen in Haaren' in *Pharmazie i.u. Zeit*, 22, pp. 65–77

Shah, I. Petroczi, A. Novacek, M. Ránky, M. Naughton, DP. (2014) 'Hair-based rapid analyses for multiple drugs in forensics and doping: application of dynamic multiple reaction monitoring with LC-MS/MS'. *Chemistry Central Journal* Dec 13;8(1):73. Doi: 10.1186/s13065-014-0073-0.

Simmons, J. Cookman, L. Kang, C. Skinner, C. (2011) 'Three cases of "spice" exposure' *Clin. Toxicol.*, 2011, 49, 431–433 Doi: 10.3109/15563650.2011.584316

Smith, JP. Oliver, S. Sutcliffe, B. Banks, C. (2015) 'An overview of recent developments in the analytical detection of new psychoactive substances (NPS)' *Analyst* 2015, 140, 4932-4948 DOI: 10.1039/C5AN00797F

Soussan, C. & Kjellgren, A. (2016) The users of Novel Psychoactive Substances: Online survey about their characteristics, attitudes and motivations *International Journal of Drugs Policy* June 2016 Volume 32, Pages 77–84 http://dx.doi.org/10.1016/j.drugpo.2016.03.007

Taylor, M. Lees, R. Henderson, G. Lingford-Hughes, A. Macleod, J. Sullivan, J. Hickman, M. (2016) 'Comparison of cannabinoids in hair with self-reported cannabis consumption in heavy, light and non-

cannabis users' *Drug and Alcohol Review* Jun 2016. Open Access DOI: <http://dx.doi.org/10.1016/j.drugpo.2016.03.007>

UIC (2008) *Occupational Health and Safety Group 2008: Developing management arrangements for the control of safety risks related to the influence of alcohol, drugs, and/or psychoactive medication. International Union of Railways, Paris* pp. 1-17

Yegles, M. (2005) 'Pitfalls in hair analysis: Cosmetic treatment'. *Ann. Toxicol. Anal.* 2005, 17, 275.

Vincenti, M. Salomone, A. Gerace, E. Pirro, V. (2013) 'Application of mass spectrometry to hair analysis for forensic toxicological investigations' *Mass Spectrometry Reviews*. Volume 32, Issue 4, pages 312–332, July/August 2013 **doi: 10.1002/mas.21364**

Wales Online (2013) 'Drug Addicts try to fool hair test' 21 march 2013 <http://www.walesonline.co.uk/news/wales-news/drug-addicts-try-fool-hair-1861374> August 10, 2009 Volume 189, Issues 1-3, Pages e41–e43

Wennig, R. (2000) 'Potential problems with the interpretation of hair analysis results'. *Forensic Science International* Volume 107, Issues 1-3, Pages 5-12. DOI: [http://dx.doi.org/10.1016/S0379-0738\(99\)00146-2](http://dx.doi.org/10.1016/S0379-0738(99)00146-2)

Non-literature sources

AlwaysTestClean.com <http://alwaystestclean.com/pass-hair-drug-test/>

Amazon (UK) listing for Zydor Ultraclean https://www.amazon.co.uk/Zydor-Ultra-Clean-Shampoo-Conditioner/dp/B003ZUW8EI/ref=sr_1_cc_1_a_it?s=aps&ie=UTF8&qid=1462909031&sr=1-1-catcorr&keywords=hair+detox+drugs

Amazon (UK) listing for Test Pass https://www.amazon.co.uk/Test-Pass-Detox-Shampoo-Single/dp/B00GMQUPO2/ref=pd_sim_194_5?ie=UTF8&dpID=418IHt1U8sL&dpSrc=sims&preST=_AC_UL160_SR160%2C160_&refRID=7TQAF5PB97FPFZDXAKW0

Amazon (UK) listing for Nexxus Aloe Rid https://www.amazon.co.uk/NEXXUSS-ALOE-GENTLE-CLARIFYING-SHAMPOO/dp/B000GFZW8Y/ref=pd_sim_sbs_121_11?ie=UTF8&dpID=31zWc6iae9L&dpSrc=sims&preST=_AC_UL160_SR160%2C160_&refRID=93ESTJR9FA5N0TPS3F9Z

DNA Legal 'Designer Drugs and Legal Highs' <https://www.dnalegal.com/hair-drug-test>

Drugs Forum - <https://drugs-forum.com/forum/showthread.php?t=193927>

DrugRehabAlliance.com 'The 2016 guide – how to pass the hair drug test' <http://drugrehaballiance.com/how-to-pass-hair-follicle-test/>

European Workplace Drug Testing Society (EWDTS 2015) 'Guidelines' <http://www.ewdts.org/data/uploads/documents/ewdts-guideline-hair-v2.0.pdf>

HairFollicleDrugTest.info 'How the hair follicle drug test works' <http://hairfollicledrugtest.info>

High Existence 'Synthetic Marijuana – DO NOT DO IT' <http://highexistence.com/topic/synthetic-marijuana-do-not-do-it/>

Ironmonger, J. (2014) 'Workplace drug testing "on the rise", say providers' (internet) BBC News 3rd Oct 2014 <http://www.bbc.co.uk/news/uk-29465755>

Marijuana.com 'What you need to know about hair testing'

<http://marijuana.com/community/threads/what-you-need-to-know-about-hair-testing.286845/>

PassYourTest.com - <http://passyourtest.com>

PassYourDrugTest.com - <http://passyourdrugtest.com>

420 Magazine. Thread: 'I passed the psychmedics hair follicle test with six days clean'
<https://www.420magazine.com/forums/420-drug-testing/214509-i-passed-psychemedics-hair-follicle-drug-test-6-days-clean.html>

420 Magazine. Thread: 'Methods to beat a hair drug test' <https://www.420magazine.com/forums/420-drug-testing/135614-methods-beat-hair-drug-test.html>

Reddit https://www.reddit.com/r/trees/comments/1ccp3s/how_i_passed_a_hair_follicle_test/

topix.com 'Does legal incense show up on a drugs test?' <http://www.topix.com/forum/city/herkimer-ny/T6NV0B8G56ML47F5D/p10>

UKCIA 'Taking the piss - drug testing at work' <http://www.ukcia.org/culture/pisstest.php>

TestingClean.com – 'How to Pass a Drug test' <http://testingclean.com>

The Weed Blog – 'How to pass a hair follicle drug test for marijuana'
<http://www.theweedblog.com/how-to-pass-a-hair-follicle-drug-test-for-marijuana/>

WikiHow 'How to Pass a Hair Follicle Test' <http://www.wikihow.com/Pass-a-Hair-Follicle-Drug-Test>

Youtube (2015) Macujo method – does it help pass a hair drug test?
<https://www.youtube.com/watch?v=EhUu80X7MY0>