

The scale of predictability*

F.M. Bandi,[†] B. Perron,[‡] A. Tamoni,[§] C. Tebaldi[¶]

September 21, 2017

Abstract

We introduce a new stylized fact: the hump-shaped behavior of slopes and coefficients of determination as a function of the aggregation horizon when running (forward/backward) predictive regressions of *future* excess market returns onto *past* economic uncertainty (as proxied by market variance, consumption variance, or economic policy uncertainty). To justify this finding formally, we propose a novel modeling framework in which predictability is specified as a property of low-frequency components of both excess market returns and economic uncertainty. We dub this property *scale-specific predictability*. We show that classical predictive systems imply restricted forms of *scale-specific predictability*. We conclude that for certain predictors, like economic uncertainty, the restrictions imposed by classical predictive systems may be excessively strong.

JEL classification: C22, E32, E44, G12, G17

Keywords: long run, predictability, aggregation, risk-return trade-off

*This paper was briefly circulated under the title “Economic uncertainty and predictability.” We are grateful to the Editor, Zhengjun Zhang, two anonymous reviewers, Yanqin Fan (the Toulouse discussant), Lars P. Hansen, L. Mancini (the EFA discussant), A. Neuberger, F. Ortú, A. Patton and A. P. Taylor for their helpful comments. We thank seminar participants at the 69th European Meeting of the Econometric Society in Geneva, the Australian Conference for Economists 2015 (Brisbane, 2015), the 2014 NBER-NSF Time Series Conference at the FRB of St. Louis, the 41st EFA in Lugano, the Workshop “Measuring and Modelling Financial Risk with High-Frequency Data” (Florence, 2014), the 2014 SoFiE Annual Conference in Toronto, the 2014 Financial Econometrics Conference at the Toulouse School of Economics, the 2014 Workshop on Uncertainty and Economic Forecasting at University College of London, the 2013 NBER Summer Institute, the 2013 CIREQ Econometrics Conference on Time Series and Financial Econometrics, the Ninth CSEF-IGIER Symposium on Economics and Institutions, the 7th International Conference on Computational and Financial Econometrics, Aspect Capital, U Penn Economics, Duke Economics, LSE Finance, Johns Hopkins Economics, Tilberg University, Erasmus University, Tinbergen Institute, Rutgers Business School, CK Graduate School of Business and Tsinghua University.

[†]Johns Hopkins University and Edhec-Risk Institute. Address: 100 International Drive, Baltimore 21202, USA. E-mail: fbandi1@jhu.edu

[‡]Université de Montréal, Department of Economics, CIRANO and CIREQ. E-mail: benoit.perron@umontreal.ca.

[§]London School of Economics, Department of Finance. E-mail: a.g.tamoni@lse.ac.uk.

[¶]Università Bocconi. E-mail: claudio.tebaldi@unibocconi.it.

1 Introduction

The introduction to the 2013 Nobel for Economic Sciences states: “*There is no way to predict whether the price of stocks and bonds will go up or down over the next few days or weeks. But it is quite possible to foresee the broad course of the prices of these assets over longer time periods, such as the next three to five years...*”

Hard-to-detect predictability over short horizons is generally viewed as the result of a low signal-to-noise problem. The magnitude of shocks to returns swamps predictable variation in expected stock returns. The aggregation of stock returns over longer horizons, however, operates as a signal extraction process uncovering predictability.

Existing work has highlighted the empirical usefulness of aggregating *both* the regressand (excess market returns) and the regressor (the market return predictor). Specifically, Bandi and Perron (2008) have suggested running adapted (to time t information) regressions of *forward* aggregated returns (i.e., long-run *future* returns) on *backward* aggregated predictors (long-run *past* market variance, in their case), rather than on raw predictors, as common in the literature. The use of forward/backward aggregation was shown to lead to a strengthening of variance-induced predictability over the long-run, the 10-year horizon being the longest prediction horizon considered in that paper.¹

We make three contributions. First, we show that the relation between *future* excess market returns and *past* uncertainty, as proxied by market variance (Bandi and Perron, 2008), consumption variance (Tamoni, 2011) and economic policy uncertainty (EPU, henceforth; see Baker, Bloom and Davis, 2016), is *hump-shaped*. The forward/backward regressions are conducted in this paper over horizons of aggregation reaching 20 years, thereby doubling the 10-year horizon spanned in the existing work. The peak of predictability is around 16 years. Estimated slopes and R^2 s feature increasing (resp. decreasing) dynamics before (resp. after) the 16-year mark. Around 16 years, the reported R^2 s may reach a value of about 55%.

¹The long-run predictability of past variance has been reported to be robust to the use of alternative variance notions (Tamoni, 2011, uses consumption variance) and the dynamics of the variance process (Sizova, 2013, assumes long memory in variance). Among other stylized facts regarding stock returns, such predictability has also been justified in the context of an asset pricing model with loss aversion (see Bonomo, Garcia, Meddahi, and Tedongap, 2015).

Second, we show that a traditional predictive system in which excess market returns are predicted by a persistent uncertainty process would find it hard to replicate the structure and magnitude of the reported hump-shaped behavior upon two-way aggregation. Theory and simulations lead to this conclusion. If a traditional predictive system is an unlikely data generating process for the reported result, what would a more likely data generating process look like?

In its third contribution, the paper uses a data generating process in which returns and uncertainty are interpreted as linear aggregates of components operating over different frequencies. Predictability is not modeled directly on the raw series. Rather, it is modeled as a property of individual return and uncertainty components. We dub this property *scale-specific predictability*. Theoretically, we show that should components with cycles of suitable lengths be linked by a predictability relation, then two-way (forward/backward) aggregation would yield hump-shaped patterns in estimated slopes and R^2 s. Empirically, after filtering excess returns and uncertainty components, we find predictability between components with cycles between 8 and 16 years. In agreement with theory, this type of *scale-specific predictability* should yield, upon two-way aggregation, a hump-shaped pattern with a peak at 16 years. This outcome is, as discussed above, consistent with data.

We show that the notion of scale-specific predictability is general in the sense that it would also apply to classical predictive systems. However, in the case of classical predictive systems, scale-specific predictability would be tightly parametrized across scales. Our proposed approach can, therefore, be viewed as freeing up the nature of predictability across scales and, as a result, for the raw series.

We conclude with an asset pricing model with Epstein-Zin recursive preferences which begins with component-wise structures for consumption growth, dividend growth and their conditional variance process. Consistent with our empirical findings, the model yields scale-specific risk-return trade-offs, i.e., linear dependencies between future values of specific components of the (excess) market return process and past values of the *same* components of the variance process, as a feature of its equilibrium. While the model provides an economic channel which justifies our empirical results regarding the existence of scale-wise dependencies between returns and variances, we view

the paper’s contribution as being methodological and, as such, broadly applicable to predictors other than variance. In our final remarks in Section 9, we comment further on this issue.

The evaluation of low-frequency contributions to economic and financial time series has a long history, one which we cannot attempt to review here. Barring fundamental methodological and conceptual differences having to do with our assumed data generating process, the approach adopted in this paper shares features with successful existing approaches.

Beveridge and Nelson (1981) popularized time series decompositions into stochastic trends and transitory components. We too operate with component-wise decompositions in which the components (more than two, in our framework) feature different levels of (calendar-time) persistence. Comin and Gertler (2006) argue that the common practice, in business-cycle research, of including longer than 8-year oscillations into the trend (see e.g., Baxter and King, 1999) may be associated with significant loss of information. Comin and Gertler (2006) decompose a series into a “high-frequency” component with cycles between 2 and 32 quarters and a “medium-frequency” component with cycles between 32 and 200 quarters. As emphasized, we focus on a number of components larger than two, thereby capturing cycles of different length (8 years being the upper bound of one of these cycles). News arrivals at various frequencies also characterize the multifractal regime switching approach of Calvet and Fisher (2001, 2007). As we will show below, modeling (through Wold representations, c.f. Section 3) and identification (through multiresolution filters, c.f. Section 5) are conducted differently in the present paper. As in Hansen and Scheinkman (2009), we employ operators to extract low-frequency information (c.f. Section 5). In our case, it is the low-frequency information embedded in the components. Finally, we show that essential scale-wise information in the extracted components can be summarized by a finite number of non-overlapping, “fundamental” points, the result of an econometric process called “decimation” (c.f., again, Section 5). These points can be viewed as being akin to “the small number of data averages” used by Müller and Watson (2008) to identify low-frequency information in the raw data. In our framework, these “fundamental” points are scale-specific and their locations correspond to the support of uncorrelated innovations over a specific scale. As such, they are particularly useful to formalize our notion of frequency-specific, or scale-specific, predictability, as shown in Section 6 and in Section 7.

Scale-wise specifications have proven successful in consumption models to explain the market risk premium (Ortu, Tamoni and Tebaldi, 2013, and Tamoni, 2011) and define granular notions of systematic risk through scale-wise betas (Bandi and Tamoni, 2016). This paper formalizes the role of scale-wise specifications in the context of a new methodological approach to predictability.

The work on stock return predictability is broad. The literature documents return forecastability induced by financial ratios, see e.g. Campbell and Shiller (1988), Lamont (1998), Kelly and Pruitt (2013), interest rate variables, see e.g. Fama and Schwert (1977), Fama and French (1989) and macroeconomic variables, see e.g. Lettau and Ludvigson (2001), Menzly et al. (2004), Nelson (1976) and Campbell and Vuolteenaho (2004). The notion of return predictability has led to controversy (e.g., Welch and Goyal, 2008, for a critique, and Cochrane, 2008, for a well-known defense).

We contribute to this extensive literature by emphasizing that traditional predictive systems may be viewed as restricted versions of the data generating process we propose. We free up *implicit* (given standard approaches) links across frequencies and provide a parsimonious framework to capture predictive relations at different frequencies.

Before continuing, we emphasize that the paper has an Online Supplement containing extensive simulations, supporting proofs for the pricing model, and additional results. In what follows, when we refer to the simulations, it should always be understood that they can be found in the Supplement. Should other results (mentioned in the main text) also be in the Supplement, we will explicitly state it.

2 Low-frequency humps

The empirical analysis in this paper is based on yearly data from 1930 to 2014. Appendix B describes the data and the construction of the variables.

We begin by running forward/backward regressions of long-run future excess market returns on long-run past *market* variance:

$$r_{t+1,t+h} = \alpha_h + \beta_h v_{t-h+1,t} + u_{t+1,t+h}, \quad (1)$$




Figure 1: R^2 values obtained by regressing forward-aggregated excess market returns on backward aggregated market variance (blue line, with square), consumption variance (red line, with circles), and (squared) economic policy uncertainty (black line, with diamonds) for different levels of aggregation (on the horizontal axis).

where $r_{t+1,t+h}$ and $v_{t-h+1,t}$ are non-overlapping sums of logarithmic excess market returns and return variances over an horizon of length h years.

Empirical results are displayed in Table 1-Panel A1 (horizons 1 to 10 years) and Table 1-Panel A2 (horizons 11 to 20 years). Panel A1 and A2 report estimated regression coefficients, adjusted R^2 statistics (in square brackets) and heteroskedasticity and autocorrelation-consistent t -statistics for the hypothesis that the regression coefficients are zero (in parentheses). The table also reports, in curly brackets, the rescaled t -statistics recommended by Valkanov (2003).²

[Insert Table 1 about here]

In Table 1-Panel A1, we report horizons of aggregations up to 10 years: *future* excess market returns are correlated with *past* market variance. Dependence increases with the horizon, and is strong in the long run, with R^2 values between 8 and 10 years ranging between 14.7% and 36.1%.

In Table 1-Panel A2 we extend the two-way regressions to horizons between 11 years and 20 years. The R^2 values reach their peak (around 55%) at 16 years. The structure of the R^2 s, before and after, is roughly tent-shaped (c.f., Fig. 1). Using Valkanov's rescaled t -statistics as a metric, past

²Valkanov's methods have become standard tools in the predictability literature. We use them here to evaluate robustness. We recall, however, that they are justifiable under a classical data generating process (as in Eqs. (2) and (3)), regressors near unity, and aggregation of the regressand, of the regressor, or both.

market variance is a powerful predictor of future excess returns (leading to statistically significant slope estimates at the 2.5% level) for horizons ranging between 11 and 16 years.³

Replacing market variance with *consumption* variance and EPU does not modify the previous results in any meaningful way (see Table 2 and 3). We will therefore not comment on these measures separately.

[Insert Table 2 and 3 about here]

We argue that classical predictive systems would find it hard to replicate the observed hump-shaped behavior. This is easy to see in theory. A traditional predictive system (on demeaned variables) would write:

$$r_{t+1} = \beta v_t + u_{t+1}, \tag{2}$$

$$v_{t+1} = \rho v_t + \varepsilon_{t+1}, \tag{3}$$

where u_{t+1} and ε_{t+1} are possibly cross-correlated, white noise shocks and $0 < \rho < 1$.

When aggregating r_{t+1} forward and v_t backward over an horizon h , the theoretical slope of the regression on forward/backward aggregates becomes $\beta\rho^h$, but $\beta\rho^h \rightarrow 0$ as $h \rightarrow \infty$.⁴ Similarly, the R^2 should go to zero with the horizon of aggregation. Fig. 1 shows, instead, that the R^2 increases steeply to about 55% before decreasing equally sharply.⁵

Leaving theoretical considerations aside, in the simulations we conduct an experiment similar in

³By generating stochastic trends, forward/backward aggregation could lead to spurious (in the sense of Granger and Newbold, 1974, and Phillips, 1986) predictability. If the reported predictability were induced mechanically by aggregation, however, *contemporaneous* (i.e., forward/forward) aggregation should also lead to patterns that are similar to those found with forward/backward aggregation. In all cases above, one could show that this is not the case. The corresponding tables are not reported for conciseness but can be provided by the authors upon request. Simulations confirm the statement. In addition, spurious behavior would prevent a tent-shape pattern from arising in the slopes, t -statistics and R^2 s from predictive regressions on the aggregated series because it would likely lead to upward trending behavior in them. We will later show that tent-shaped patterns are, instead, a natural by-product of an alternative data generating process.

⁴The reported “slope” should be intended as the resulting slope from direct forward/backward iterations of the model. In light of the dependence between the regression residuals obtained by iteration and the backward-aggregated regressors, this slope does not coincide with the one associated with the true conditional mean of forward-aggregated regressands onto backward-aggregated regressors. Such slope, for a large aggregation horizon h , would be approximately $\frac{\beta}{1+\rho} \frac{1}{h}$. Hence, it would also vanish as $h \rightarrow \infty$. We thank Nour Meddahi for discussions about this point.

⁵As shown by Sizova (2013), a “large” ρ , captured by long memory in her framework, would help over horizons over which the statistics are reported as being monotonically increasing (1 to about 16 years). A long-memory variance process would, however, find it difficult to capture the hump-shaped dynamics illustrated above and further discussed below.

spirit to that in Boudoukh et al. (2008) and show that classical predictive systems are, in any finite sample, unlikely to generate the tent-shaped dynamics detected in the data. Given carefully-chosen parameter values for Eq. (2) and Eq. (3), the experiment finds that the percentage of simulated paths delivering *both* hump-shaped R^2 values and hump-shaped slope values, as well as R^2 's in excess of 50%, is about 1.15.

In the next section, we express excess market returns and economic uncertainty as linear combinations of $J > 1$ uncorrelated, mean-zero components $r^{(j)}$ and $v^{(j)}$ with $1 \leq j \leq J$. Each component will be shown to operate over a specific scale or frequency. We will discuss how a predictive system analogous to Eq. (2) and Eq. (3), but applied to specific component(s) (i.e., a *scale-specific predictive system* as in Eq. (12) and Eq. 13 of Section 4), may yield the reported hump-shaped pattern(s) upon two-way aggregation of the raw series. Proposition I in Section 6 will then link scale-specific predictability to two-way aggregation.

We will view scale-wise predictability as a “spectral” feature of the series. Predictability upon two-way aggregation will, instead, be interpreted as a way to translate scale-specific predictability into return predictability for the long haul, with all of its applied implications, including long-run asset allocation.

3 Scale-specific components

In the following subsection we summarize the basic steps to separate a covariance-stationary process into orthogonal, scale-specific components. A formalization of this decomposition based on Hilbert space theory is given in Ortu et al. (2017) (OSTT henceforth). Importantly, since the procedure is applicable to any process for which a classical Wold representation applies, it can be regarded as general.

3.1 A scale-wise representation of a covariance-stationary scalar process

Let $\mathbf{x} = \{x_t\}_{t \in \mathbb{Z}}$ be a covariance-stationary scalar process defined onto the space $L^2(\Omega, \mathcal{F}, \mathbb{P})$. For simplicity, but without loss of generality, assume the process has a zero mean.⁶

⁶Adding the mean would, of course, simply add a constant to its Wold representation.

The Wold representation of \mathbf{x} states that there exists a unit variance white noise process $\boldsymbol{\varepsilon} = \{\varepsilon_t\}_{t \in \mathbb{Z}}$ such that, for any t in \mathbb{Z} ,

$$x_t = \sum_{k=0}^{+\infty} \alpha_k \varepsilon_{t-k}, \quad (4)$$

where the equality is in the L^2 -norm and $\{\alpha_k\}_{k \in \mathbb{N}_0}$ is a square-summable sequence of real coefficients with $\alpha_k = \mathbb{E}(x_t \varepsilon_{t-k})$.

Let us now introduce scales and define the innovation process at scale j with $j \in \mathbb{N}$. If $j = 1$, the innovation process at scale 1, denoted by $\boldsymbol{\varepsilon}^{(1)} = \{\varepsilon_t^{(1)}\}_{t \in \mathbb{Z}}$, is defined as the process whose terms are

$$\varepsilon_t^{(1)} = \frac{\varepsilon_t - \varepsilon_{t-1}}{\sqrt{2}}.$$

We observe that $\varepsilon_t^{(1)}$ has a zero mean and its variance is equal to 1 for all t . More generally, we define the innovation process at scale j as the process $\boldsymbol{\varepsilon}^{(j)} = \{\varepsilon_t^{(j)}\}_{t \in \mathbb{Z}}$ with

$$\varepsilon_t^{(j)} = \frac{1}{\sqrt{2^j}} \left(\sum_{i=0}^{2^{j-1}-1} \varepsilon_{t-i} - \sum_{i=0}^{2^{j-1}-1} \varepsilon_{t-2^{j-1}-i} \right). \quad (5)$$

The structure of the shocks $\boldsymbol{\varepsilon}^{(j)}$ is that of a Discrete Haar Transform (DHT). Once a scale level j is set, one may consider the sub-series of $\boldsymbol{\varepsilon}^{(j)}$ defined on the support $S_t^{(j)} = \{t - k2^j : k \in \mathbb{Z}\}$. The process $\boldsymbol{\varepsilon}^{(j)}$ is a moving average of order $2^j - 1$ with respect to the Wold innovations of \mathbf{x} . In calendar time, there is correlation between the variables $\varepsilon_{t-2^j}^{(j)}$ and $\varepsilon_{\tau-2^j}^{(j)}$ with $|t - \tau| \leq 2^j - 1$. Nonetheless, each sub-series $\{\varepsilon_{t-2^j}^{(j)}\}_{k \in \mathbb{Z}}$ is a unit variance white noise process on the support $S_t^{(j)}$.

Technically, we are decomposing the space $\mathcal{H}_t(\boldsymbol{\varepsilon})$ of infinite moving averages whose underlying white noise process is $\boldsymbol{\varepsilon}$, i.e.,

$$\mathcal{H}_t(\boldsymbol{\varepsilon}) = \left\{ \sum_{k=0}^{+\infty} a_k \varepsilon_{t-k} : \sum_{k=0}^{+\infty} a_k^2 < +\infty \right\},$$

into orthogonal subspaces⁷

$$\mathcal{W}_t^{(j)} = \left\{ \sum_{k=0}^{+\infty} b_k^{(j)} \varepsilon_{t-k2j}^{(j)} \in \mathcal{H}_t(\varepsilon) : b_k^{(j)} \in \mathbb{R} \right\},$$

with $j = 1, \dots, +\infty$.

We now turn to the Wold coefficients at scale j . The process \mathbf{x} is contained in $\mathcal{H}_t(\varepsilon)$. Therefore, the decomposition of the space $\mathcal{H}_t(\varepsilon)$ induces the following representation of \mathbf{x} :

$$x_t = \sum_{j=1}^{+\infty} x_t^{(j)}, \quad (6)$$

where each $x_t^{(j)}$ is the projection of x_t on the subspace $\mathcal{W}_t^{(j)}$, with $j \in \mathbb{N}$, and the equality is, again, in the L^2 -norm. By construction, for a fixed t , the components $x_t^{(j)}$ are orthogonal to each other. Since each $x_t^{(j)}$ belongs to $\mathcal{W}_t^{(j)}$, we have that

$$x_t^{(j)} = \sum_{k=0}^{+\infty} \psi_k^{(j)} \varepsilon_{t-k2j}^{(j)}$$

for some square-summable sequence of real coefficients $\{\psi_k^{(j)}\}_{k \in \mathbb{N}_0}$.

Each coefficient $\psi_k^{(j)}$ is obtained by projecting \mathbf{x} on the linear subspace generated by the (scale-specific) innovation $\varepsilon_{t-k2j}^{(j)}$:

$$\psi_k^{(j)} = \mathbb{E} \left[x_t \varepsilon_{t-k2j}^{(j)} \right]. \quad (7)$$

Substituting the expression of $x_t^{(j)}$ into Eq. (6), we arrive at the *extended* Wold representation of x_t , that is⁸

$$x_t = \sum_{j=1}^{+\infty} \sum_{k=0}^{+\infty} \psi_k^{(j)} \varepsilon_{t-k2j}^{(j)}. \quad (8)$$

Eq. (8) is a (type of) Wold representation describing any weakly-stationary time series of interest as a linear combination of shocks classified on the basis of their arrival time as well as their scale.

It is now interesting to discuss the connection between the coefficients $\psi_k^{(j)}$ of the *extended* Wold

⁷The orthogonal decomposition into subspaces is provided in Theorem 1 of OSTT (2017).

⁸See Theorem 2 in OSTT (2017).

representation and the coefficients α_k of the *classical* Wold representation. Exploiting Eq. (7) after expressing $\varepsilon_t^{(j)}$ as a finite linear combination of variables ε_t as in Eq. (5) and using Eq. (4), we obtain

$$\psi_k^{(j)} = \frac{1}{\sqrt{2^j}} \left(\sum_{i=0}^{2^{j-1}-1} \alpha_{k2^j+i} - \sum_{i=0}^{2^{j-1}-1} \alpha_{k2^j+2^{j-1}+i} \right). \quad (9)$$

Just like the shocks $\varepsilon_t^{(j)}$ s are Discrete Haar Transforms (DHTs) of the high-frequency shocks in the classical Wold (see Eq. (5)), the coefficients $\psi_k^{(j)}$ s are DHTs of the coefficients in the classical Wold. Since the Wold coefficients α_k are unique, it follows that the coefficients $\psi_k^{(j)}$ are unique, and time-invariant, functions of the coefficients α_h given the Haar structure.

3.2 The multivariate case

Consider the mean-zero, covariance stationary, bi-variate process $\mathbf{X} = \{(y_t, x_t)'\}_{t \in \mathbb{Z}}$. Its Wold representation is, again,

$$\mathbf{X}_t = \begin{pmatrix} y_t \\ x_t \end{pmatrix} = \sum_{k=0}^{+\infty} \boldsymbol{\alpha}_k \boldsymbol{\varepsilon}_{t-k},$$

where $\boldsymbol{\varepsilon} = \{(u_t, e_t)'\}_{t \in \mathbb{Z}}$ is a bi-variate vector of possibly cross-correlated white noise shocks and the Wold coefficients $\boldsymbol{\alpha}_k$ are, for all $k \in \mathbb{N}_0$, 2×2 matrices.

By analogy with the scalar case, we can now write

$$\begin{pmatrix} y_t \\ x_t \end{pmatrix} = \sum_{j=1}^{+\infty} \sum_{k=0}^{+\infty} \boldsymbol{\Psi}_k^{(j)} \boldsymbol{\varepsilon}_{t-k2^j}^{(j)}. \quad (10)$$

For any $j \in \mathbb{N}$, the 2×2 matrices $\boldsymbol{\Psi}_k^{(j)}$ are unique Haar transforms of the original Wold coefficients:

$$\boldsymbol{\Psi}_k^{(j)} = \frac{1}{\sqrt{2^j}} \left(\sum_{i=0}^{2^{j-1}-1} \boldsymbol{\alpha}_{k2^j+i} - \sum_{i=0}^{2^{j-1}-1} \boldsymbol{\alpha}_{k2^j+2^{j-1}+i} \right).$$

Similarly,

$$\boldsymbol{\varepsilon}_t^{(j)} = \frac{1}{\sqrt{2^j}} \left(\sum_{i=0}^{2^{j-1}-1} \boldsymbol{\varepsilon}_{t-i} - \sum_{i=0}^{2^{j-1}-1} \boldsymbol{\varepsilon}_{t-2^{j-1}-i} \right). \quad (11)$$

We emphasize that the components of the *extended* Wold in Eq. (10) are uncorrelated. As an example, take the first and the second scale and notice that the corresponding shocks are defined as follows:

$$\boldsymbol{\varepsilon}_t^{(1)} = \begin{pmatrix} \frac{u_t - u_{t-1}}{\sqrt{2}} \\ \frac{e_t - e_{t-1}}{\sqrt{2}} \end{pmatrix}, \quad \boldsymbol{\varepsilon}_{t-2}^{(1)} = \begin{pmatrix} \frac{u_{t-2} - u_{t-3}}{\sqrt{2}} \\ \frac{e_{t-2} - e_{t-3}}{\sqrt{2}} \end{pmatrix}, \quad \boldsymbol{\varepsilon}_{t-4}^{(1)} = \begin{pmatrix} \frac{u_{t-4} - u_{t-5}}{\sqrt{2}} \\ \frac{e_{t-4} - e_{t-5}}{\sqrt{2}} \end{pmatrix}, \dots$$

and

$$\boldsymbol{\varepsilon}_t^{(2)} = \begin{pmatrix} \frac{(u_t + u_{t-1}) - (u_{t-2} + u_{t-3})}{\sqrt{4}} \\ \frac{(e_t + e_{t-1}) - (e_{t-2} + e_{t-3})}{\sqrt{4}} \end{pmatrix}, \quad \boldsymbol{\varepsilon}_{t-4}^{(2)} = \begin{pmatrix} \frac{(u_{t-4} + u_{t-5}) - (u_{t-6} + u_{t-7})}{\sqrt{4}} \\ \frac{(e_{t-4} + e_{t-5}) - (e_{t-6} + e_{t-7})}{\sqrt{4}} \end{pmatrix}, \dots$$

It is immediate to verify that the first and the second components are orthogonal at all leads and lags on the supports $S_t^{(1)}$ and $S_t^{(2)}$ defined in Subsection 3.1. Formally, for generic components j and l , we have

$$\mathbb{E} \left[\mathbf{g}_{t-m2^j}^{(j)} \mathbf{g}_{t-n2^l}^{(l)'} \right] = \mathbf{0} \quad \forall j \neq l, \quad \forall m, n \in \mathbb{N}_0, \quad \forall t \in \mathbb{Z},$$

where $\mathbf{g}^{(j)} = \left\{ \left(y_t^{(j)}, x_t^{(j)} \right)' \right\}_{t \in \mathbb{Z}}$.⁹

We now turn to the focus of our analysis, i.e., scale-specific predictability.

4 Scale-specific predictive systems

Consider a regressand y and a predictor x . Assume y and x are covariance-stationary and, therefore, admit an *extended* Wold representation, see Eq. (10). Assume, also, that for some scale $1 \leq j^* \leq J$, the components of the processes y and x (written in scale-specific times) are such that

$$y_{k2^{j^*} + 2^{j^*}}^{(j^*)} = \beta_{j^*} x_{k2^{j^*}}^{(j^*)} + u_{k2^{j^*} + 2^{j^*}}^{(j^*)} \quad (12)$$

$$x_{k2^{j^*} + 2^{j^*}}^{(j^*)} = \rho_{j^*} x_{k2^{j^*}}^{(j^*)} + \varepsilon_{k2^{j^*} + 2^{j^*}}^{(j^*)}, \quad (13)$$

where $\varepsilon_{k2^{j^*} + 2^{j^*}}^{(j^*)}$ is a white noise process in scale time with a zero mean and a variance $\sigma_{\varepsilon_{j^*}}^2$ and $u_{k2^{j^*} + 2^{j^*}}^{(j^*)}$ is a white noise (again, in scale time) forecast error with a zero mean and a variance $\sigma_{u_{j^*}}^2$.

⁹The orthogonality between components is also a property of the multivariate Wold representation, see Cerreia-Vioglio et al. (2017) and Theorem 3 in OSTT (2017).

The shocks $u_{k2^{j^*}+2j^*}^{(j^*)}$ and $\varepsilon_{k2^{j^*}+2j^*}^{(j^*)}$ are possibly cross-correlated. Assume all other components $\{y_{k2^j+2j}^{(j)}, x_{k2^j+2j}^{(j)}\}$ with $j \neq j^*$ are mean-zero, finite variance, white noise processes, uncorrelated with each other.¹⁰

Eqs. (12)-(13) define a predictive system on individual layers of the $\{(y_t, x_t)'\}_{t \in \mathbb{Z}}$ process to be contrasted with the traditional system written on the raw series directly.

There is an understanding in the predictability literature that slow-moving predictors should drive slow-moving conditional means. This relation is, however, hidden by short-term noise. The noise leads to the appearance of low predictability in the short-run, large long-run predictability being the outcome of noise reduction through return aggregation. There is also an understanding that predictors may be imperfect (Pastor and Stambaugh, 2009). In the context of a different conceptual framework, Eqs. (12) and (13) account for both effects: the link between slow-moving components (for a large j) and, by being defined on components rather than on noisier (i.e., “imperfect”) raw series, the “imperfection” of predictors (and regressands).

4.1 Classical predictive systems and scale-specific predictability

Can a typical predictive system yield scale-specific predictability? The answer is positive. However, classical predictive systems impose tight restrictions on the nature of predictability across scales. To see this, consider a vector autoregressive process of order 1 (VAR(1)) for the bi-variate process $(y_t, x_t)'$, i.e.,

$$\begin{pmatrix} y_t \\ x_t \end{pmatrix} = A \begin{pmatrix} y_{t-1} \\ x_{t-1} \end{pmatrix} + \begin{pmatrix} u_t \\ e_t \end{pmatrix}.$$

As in Subsection 3.2, the vector $(u_t, \varepsilon_t)' = \varepsilon_t$ is a vector of possibly cross-correlated white noise shocks. The standard Wold representation yields

$$\begin{pmatrix} y_t \\ x_t \end{pmatrix} = \underbrace{I_2}_{\alpha_0} \begin{pmatrix} u_t \\ \varepsilon_t \end{pmatrix} + \underbrace{A}_{\alpha_1} \begin{pmatrix} u_{t-1} \\ \varepsilon_{t-1} \end{pmatrix} + \underbrace{A^2}_{\alpha_2} \begin{pmatrix} u_{t-2} \\ \varepsilon_{t-2} \end{pmatrix} + \dots$$

¹⁰Needless to say, we are not excluding the possibility of multiple scale-wise predictive systems at different frequencies. Our emphasis on a single frequency is simply meant to illustrate the potential of a simple scale-specific predictive structure for yielding patterns found in the data.

with I_2 defining the 2×2 identity matrix. Hence, in terms of the notation in Subsection 3.2, we have $\alpha_k = A^k$.

We can now obtain the same *extended* Wold representation as in Eq. (10) with the following coefficients $\Psi_k^{(j)}$:

$$\begin{aligned}\Psi_0^{(1)} &= \frac{I_2 - A}{\sqrt{2}} \\ \Psi_1^{(1)} &= \frac{A^2 - A^3}{\sqrt{2}} = \frac{I_2 - A}{\sqrt{2}} A^2, \\ &\dots \\ \Psi_k^{(1)} &= \frac{A^{2k} - A^{2k+1}}{\sqrt{2}} = \frac{I_2 - A}{\sqrt{2}} A^{2k},\end{aligned}$$

for $j = 1$. Similarly, for a generic $j = 1, 2, \dots$, and for $k = 0$:

$$\begin{aligned}\Psi_0^{(j)} &= \frac{\underbrace{1 + A + \dots + A^{2^{(j-1)}-1}}_{2^{(j-1)} \text{ terms}} - \left(\underbrace{A^{2^{(j-1)}} + \dots + A^{2^j-1}}_{2^{(j-1)} \text{ terms}} \right)}{\sqrt{2^j}} \\ &= \frac{\left(I_2 - A^{2^{(j-1)}} \right) (I_2 - A)^{-1} - \left(I_2 - A^{2^{(j-1)}} \right) (I_2 - A)^{-1} A^{2^{(j-1)}}}{\sqrt{2^j}} \\ &= \frac{\left(I_2 - A^{2^{(j-1)}} \right)^2 (I_2 - A)^{-1}}{\sqrt{2^j}}.\end{aligned}$$

For $j = 1, 2, \dots$, and for $k > 0$:

$$\begin{aligned}\Psi_k^{(j)} &= \frac{\underbrace{1 + A + \dots + A^{2^{(j-1)}-1}}_{2^{(j-1)} \text{ terms}} - \left(\underbrace{A^{2^{(j-1)}} + \dots + A^{2^j-1}}_{2^{(j-1)} \text{ terms}} \right)}{\sqrt{2^j}} \times A^{k2^j} \\ &= \frac{\left(I_2 - A^{2^{(j-1)}} \right)^2 (I_2 - A)^{-1}}{\sqrt{2^j}} \times A^{k2^j}.\end{aligned}$$

Let us now be explicit about the components. Since $\Psi_k^{(j)} = \Psi_0^{(j)} \times A^{k2^j}$, we obtain

$$\begin{pmatrix} y_t^{(j)} \\ x_t^{(j)} \end{pmatrix}' = \sum_{k=0}^{+\infty} \Psi_k^{(j)} \varepsilon_{t-k2^j}^{(j)} = \widehat{\varepsilon}_t^{(j)} + A^{2^j} \widehat{\varepsilon}_{t-2^j}^{(j)} + A^{2 \times 2^j} \widehat{\varepsilon}_{t-2 \times 2^j}^{(j)} + \dots,$$

where $\widehat{\varepsilon}_t^{(j)} = \Psi_0^{(j)} \varepsilon_t^{(j)}$. Hence, as expected, the structure of the $\Psi_k^{(j)}$ coefficients is that of a VAR(1) defined on the support $S_t^{(j)} = \{t - k2^j : k \in \mathbb{Z}\}$. However, this implies

$$\begin{pmatrix} y_t^{(j)} \\ x_t^{(j)} \end{pmatrix} = A^{2^j} \begin{pmatrix} y_{t-2^j}^{(j)} \\ x_{t-2^j}^{(j)} \end{pmatrix} + \widehat{\varepsilon}_t^{(j)},$$

which is, in our jargon, a *scale-wise predictive system*.

The above derivations lead to two observations. First, we can rewrite the VAR(1) as an infinite sum of VAR(1)s with time steps 2^j and autoregressive matrix given by A^{2^j} . We are, of course, interested in the case

$$A = \begin{pmatrix} 0 & \beta \\ 0 & \rho \end{pmatrix},$$

which yields

$$A^{2^j} = \begin{pmatrix} 0 & \beta \times \rho^{2^j-1} \\ 0 & \rho^{2^j} \end{pmatrix}.$$

Thus, a classical predictive system implies a tightly parametrized form of scale-specific predictability. For each scale j , the predictive slope should be $\beta \times \rho^{2^j-1}$. Our approach frees up this restriction and, therefore, allows for rich dependence structures in the raw series without dispensing with parsimony.

Second, the standard predictive system only allows for scale-specific predictability between identical scales of the regressand and the regressor, cross-predictability (between two generic scales, say j and l) being excluded. Just like in the standard predictive system, even in our unrestricted framework lack of cross-predictability is *not* an assumption. As discussed in Subsection 3.2, it readily rests on the Haar structure of the scale-specific shocks.

5 Filtering the scale-specific components

Because the components $x_t^{(j)}$ are needed to validate scale-specific predictability in the data (c.f. Section 7), their extraction - the topic of this section - is empirically important.

Let, again, $\{x_{t-i}\}_{i \in \mathbb{Z}}$ be the time series of interest. Consider the case $J = 1$. We have

$$x_t = \underbrace{\frac{x_t - x_{t-1}}{2}}_{\widehat{x}_t^{(1)}} + \underbrace{\left[\frac{x_t + x_{t-1}}{2} \right]}_{\pi_t^{(1)}},$$

which effectively amounts to breaking the series down into a “transitory” component $\widehat{x}_t^{(1)}$ and a “persistent” component $\pi_t^{(1)}$. Set, now, $J = 2$. We obtain

$$x_t = \underbrace{\frac{x_t - x_{t-1}}{2}}_{\widehat{x}_t^{(1)}} + \underbrace{\left[\frac{x_t + x_{t-1} - x_{t-2} - x_{t-3}}{4} \right]}_{\widehat{x}_t^{(2)}} + \underbrace{\left[\frac{x_t + x_{t-1} + x_{t-2} + x_{t-3}}{4} \right]}_{\pi_t^{(2)}},$$

which further separates the “persistent” component $\pi_t^{(1)}$ into another “transitory” component $\widehat{x}_t^{(2)}$ and an another “persistent” component $\pi_t^{(2)}$.

The procedure can, of course, be iterated yielding a general expression for the generic component $\widehat{x}_t^{(j)}$, i.e.,

$$\widehat{x}_t^{(j)} = \frac{\sum_{i=0}^{2^{(j-1)}-1} x_{t-i}}{2^{(j-1)}} - \frac{\sum_{i=0}^{2^j-1} x_{t-i}}{2^j} = \pi_t^{(j-1)} - \pi_t^{(j)},$$

where the element $\pi_t^{(j)}$ satisfies the recursion

$$\pi_t^{(j)} = \frac{\pi_t^{(j-1)} + \pi_{t-2^{j-1}}^{(j-1)}}{2},$$

with $\pi_t^{(0)} \equiv x_t$.

In essence, for every t , $\{x_{t-i}\}_{i \in \mathbb{Z}}$ can be written as a collection of components $\widehat{x}_t^{(j)}$ with different calendar-time persistence along with a low-frequency trend $\pi_t^{(J)}$. Equivalently, it can be written as

a telescopic sum

$$x_t = \sum_{j=1}^J \underbrace{\left\{ \pi_t^{(j-1)} - \pi_t^{(j)} \right\}}_{\widehat{x}_t^{(j)}} + \pi_t^{(J)} = \pi_t^{(0)}, \quad (14)$$

in which the filtered components are naturally viewed as changes in information between scale 2^{j-1} and scale 2^j . The scales are dyadic¹¹ and, therefore, enlarge with j . The higher j , the lower the frequency. In particular, the innovations $\widehat{x}_t^{(j)} = \pi_t^{(j-1)} - \pi_t^{(j)}$ become smoother, and more persistent in calendar time, as j increases. Eq. (14) is the filtered analogue of Eq. (6).

One may also write a convenient representation of the filter using a suitable projection operator. To illustrate how the operator works, set $J = 2$. In this case,

$$\pi_t^{(2)} = \frac{x_t + x_{t-1} + x_{t-2} + x_{t-3}}{4}. \quad (15)$$

and

$$\begin{aligned} \widehat{x}_t^{(2)} &= \frac{\pi_t^{(1)} - \pi_{t-1}^{(1)}}{2} = \frac{1}{2} \left(\frac{x_t + x_{t-1}}{2} - \frac{x_{t-2} + x_{t-3}}{2} \right) \\ \widehat{x}_t^{(1)} &= \frac{\pi_t^{(0)} - \pi_{t-1}^{(0)}}{2} = \left(\frac{x_t - x_{t-1}}{2} \right) \\ \widehat{x}_{t-2}^{(1)} &= \frac{\pi_{t-2}^{(0)} - \pi_{t-3}^{(0)}}{2} = \left(\frac{x_{t-2} - x_{t-3}}{2} \right). \end{aligned} \quad (16)$$

Stacking now Eq. (15) on top of Eq. (16), we obtain:

$$\begin{pmatrix} \pi_t^{(2)} \\ \widehat{x}_t^{(2)} \\ \widehat{x}_t^{(1)} \\ \widehat{x}_{t-2}^{(1)} \end{pmatrix} = \begin{pmatrix} \frac{1}{4} & \frac{1}{4} & \frac{1}{4} & \frac{1}{4} \\ \frac{1}{4} & \frac{1}{4} & -\frac{1}{4} & -\frac{1}{4} \\ \frac{1}{2} & -\frac{1}{2} & 0 & 0 \\ 0 & 0 & \frac{1}{2} & -\frac{1}{2} \end{pmatrix} \begin{pmatrix} x_t \\ x_{t-1} \\ x_{t-2} \\ x_{t-3} \end{pmatrix}. \quad (17)$$

Denoting by $\mathcal{T}^{(2)}$ the (4×4) matrix in Eq. (17), we notice that $\mathcal{T}^{(2)}$ is orthogonal, that is $\Lambda^{(2)} \equiv \mathcal{T}^{(2)} (\mathcal{T}^{(2)})^\top$ is diagonal. Moreover, the diagonal elements of $\Lambda^{(2)}$ are non-vanishing so that $(\mathcal{T}^{(2)})^{-1} = (\mathcal{T}^{(2)})^\top (\Lambda^{(2)})^{-1}$ is well-defined. Importantly, the matrix $\mathcal{T}^{(2)}$ permits to construct the filtered components by virtue of a simple matrix multiplication.

¹¹The term ‘‘dyadic’’ refers to the fact that the horizon of the scales, i.e. $(2^{j-1}, 2^j)$, increases like powers of 2.

Similarly, by matrix inversion, one could reconstruct the original process given the filtered components. This is, again, easy to see:

$$\begin{pmatrix} x_t \\ x_{t-1} \\ x_{t-2} \\ x_{t-3} \end{pmatrix} = \left(\mathcal{T}^{(2)}\right)^{-1} \begin{pmatrix} \pi_t^{(2)} \\ \hat{x}_t^{(2)} \\ \hat{x}_t^{(1)} \\ \hat{x}_{t-2}^{(1)} \end{pmatrix}. \quad (18)$$

For an extension of this procedure to any $J \geq 2$ and a recursive algorithm for the construction of the operator matrix $\mathcal{T}^{(J)}$, associated to an arbitrary level of persistence J , we refer the reader to Mallat (1989). The matrix $\mathcal{T}^{(J)}$ will be referred to in what follows as the Haar transform (or the Haar matrix).¹²

As in Beveridge and Nelson (1981), who initiated the filtering of a time series into a stochastic trend and a transitory component, we can view the adopted filter as a nonparametric method to separate the time series into components (more than two, in our case) with different levels of (calendar-time) persistence operating at different frequencies. In our framework, the components' shocks are functions of both time and scale.

5.1 Decimation

Decimation is the process of defining *non-redundant* information, as contained in a suitable number of *non-overlapping* “typical” points, in the observed components. Equivalently, it is the process of sampling in scale time, the type of sampling used in Section 6 and in Section 7 below.

For clarity, let us now return to the case $J = 2$, as above, with the understanding that identical considerations apply more generally. Define the vector

$$X_t = [x_t, x_{t-1}, x_{t-2}, x_{t-3}]^\top.$$

By letting time t vary in the set $\{t = k2^2 \text{ with } k \in \mathbb{Z}\}$ one can now define (from $\mathcal{T}^{(2)}X_t$) the *dec-*

¹²For any dilation index j , one can also view the components $\hat{x}_t^{(j)}$ as projections of the original series x_t onto the space spanned by a system of Haar basis with translation parameter $k \in \mathbb{Z}$ (see Mallat, 1989, and Dijkerman and Mazumdar, 1994).

imated counterparts of the calendar-time components, namely $\{\widehat{x}_t^{(j)}, t = k2^j \text{ with } k \in \mathbb{Z}\}$ for $j = 1, 2$ and $\{\pi_t^{(2)}, t = k2^2 \text{ with } k \in \mathbb{Z}\}$.

The decimated points have a similar intuition as the “the small number of data averages” advocated by Müller and Watson (2008) to identify low-frequency information in the raw data. In our framework, however, these points are scale-specific and useful to formalize our notion of frequency-specific predictability. Implicitly, they were used to define scale-wise predictive systems in Eq. (12) and in Eq. (13).

In multiresolution analysis¹³ the Haar matrix is routinely used to *filter* $\{x_{t-k2^j}^{(j)}\}_{j=1,\dots,J, k \in \mathbb{Z}}$ from $\{x_{t-n}\}_{n \in \mathbb{Z}}$. While we also filter the components using the Haar matrix, one methodological novelty of this work is to also operate in the opposite direction, i.e., to propose a data generating process which specifies the law of motion of the components $\{x_{t-k2^j}^{(j)}\}_{j=1,\dots,J, k \in \mathbb{Z}}$ (see Eqs. (12)-(13)) and, only then, obtains each observation x_t as a linear combination of the components themselves (see Eq. (18)). This is, for instance, how we generate observations from a data generating process allowing for scale-specific predictability in the simulations.

Before turning to an empirical evaluations of Eqs. (12)-(13), we discuss the implications of scale-wise predictability for two-way aggregation.

6 The mapping between two-way aggregation and scale-wise predictive systems

We showed in Section 2 that forward-backward regressions uncover a strong risk-return trade-off beyond business cycle frequencies. The logic for aggregating both the regressand and the regressor resides in the intuition according to which equilibrium implications of economic models may hold for specific, highly persistent components of the variables $\{y, x\}$ while being hidden by short-term noise. Aggregation provides a natural way to extract these components, filter out the noise, and generate a cleaner signal. Using the assumed data generating process, we now formalize this logic.

¹³See, e.g., Mallat (1989). For applications to economic and financial time series, we refer to the comprehensive treatments in Ramsey (1999), Percival and Walden (2000), Gençay et al. (2001), and Crowley (2007).

Proposition I. *Assume that, for some $j = j^*$, we have*

$$\begin{aligned} y_{k2^{j^*}+2^{j^*}}^{(j^*)} &= \beta_{j^*} x_{k2^{j^*}}^{(j^*)}, \\ x_{k2^{j^*}+2^{j^*}}^{(j^*)} &= \rho_{j^*} x_{k2^{j^*}}^{(j^*)} + \varepsilon_{k2^{j^*}+2^{j^*}}^{(j)}, \end{aligned}$$

whereas $\{y_{k2^j}^{(j)}, x_{k2^j}^{(j)}\} = 0$ for $j \neq j^*$. We map scale-time (or decimated) observations into calendar-time observations by using the inverse Haar transform. Then, the forward-backward regressions

$$y_{t+1,t+h} = \beta_h x_{t-h+1,t} + \epsilon_{t+1,t+h}$$

reach the maximum R^2 of 1 over the horizon $h = 2^{j^*}$ and, at that horizon, $\beta_h = \beta_{j^*}$.

Proof. See Appendix A.

For simplicity, in Proposition I we dispense with the forecasting shocks u_t . Predictability applies to a specific j^* component. All other components are set equal to zero.

Proposition I shows that predictability on the components implies predictability upon suitable aggregation of both the regressand and the regressor. More explicitly, economic relations which apply to specific, low-frequency components will be revealed by two-way averaging. Adding short-term or long-term shocks in the form of uncorrelated components $\{y_{k2^j}^{(j)}, x_{k2^j}^{(j)}\}$, for $j < j^*$ or for $j > j^*$, or forecast errors different from zero, would solely lead to a blurring of the resulting relation upon two-way aggregation. We add uncorrelated components $\{y_{k2^j}^{(j)}, x_{k2^j}^{(j)}\}$ for $j \neq j^*$ in the simulations.

The proposition also makes explicit the fact that the optimal level of (forward/backward) aggregation should coincide with the horizon over which (frequency-specific) predictability applies. More specifically, under the above assumptions, if predictability applies to a specific detail with fluctuations between 2^{j^*-1} and 2^{j^*} periods, an R^2 of 1 would be achieved for a level of (forward/backward) aggregation corresponding to 2^{j^*} periods. Before and after, the R^2 s should display a *tent-like* behavior.

Next, we broaden the scope of classical predictability relations in the literature. We turn to regressions on the extracted components and illustrate the consistency of their findings with those

obtained, in Section 2, from two-way aggregation. This consistency, which is an implication of Proposition I, is further confirmed by simulation.

7 Predictability on the components

We focus on the component-wise relation between equity returns and market variance (Table 1). The case of consumption variance and EPU is completely analogous (Table 2 and 3) and, as earlier, we will not comment on it independently.

The filtered components are shown in Figs. 2 and 3. For an explicit interpretation of the j -th scale in terms of yearly time spans, we refer to Table 4.

[Insert Table 4 about here]




Figure 2: Decomposition into components for market returns: the calendar-time observations are solid blue lines, the scale-time or “decimated” observations are red diamonds.

Table 5 presents pair-wise correlations between individual components, for both series. Virtually all correlations are small and statistically insignificant. This evidence is suggestive of the fact that




Figure 3: Decomposition into components for market realized variance: the calendar-time observations are solid blue lines, the scale-time or “decimated” observations are red diamonds.

our proposed Haar-based filtering procedure described in Section 5 is effective in extracting the orthogonal Wold components.¹⁴

[Insert Table 5 about here]

We run scale-wise predictive regressions as in Eqs. (12)-(13). The results are reported in Table 1-Panel B.¹⁵ The strongest predictability is for $j = 4$, which corresponds to economic fluctuations between 8 and 16 years. For $j = 4$, the R^2 of the scale-wise predictive regression is a considerable 58.2%.

Consistent with Proposition I, the R^2 values upon two-way aggregation reach their peak (around

¹⁴The pair-wise correlations are obtained by using overlapping, calendar-time, or *redundant* data on the components rather than the non-overlapping, scale-time or *decimated* counterparts described in Subsection 5.1. This is, of course, due to the need of having the same number of observations for each scale. There could also be leakage between adjacent time scales. It is possible to reduce the impact of leakage by replacing the Haar filter with alternative filters with superior robustness properties (the Daubechies filter is one example). The investigation of which filter is the most suitable for the purpose of studying predictability on the scales is beyond the scopes of the present paper. As pointed out earlier, also, the use of the Haar filter is particularly helpful to relate scale-wise predictability to aggregation, a core aspect of our treatment, and yield components with cycles whose length is easily interpretable.

¹⁵Given a scale j , we work with an effective sample size of $\lceil T/2^j \rceil$ observations, where $\lceil \cdot \rceil$ denotes the smallest integer near $T/2^j$. In this empirical analysis, we consider a sample spanning the period 1930 to 2014 and $J = 4$.

55%) between 14 and 16 years ($2^j = 16$ years). Remarkably, the structure of the R^2 s, before and after, is roughly tent-shaped (see Fig. 1).

The study of low frequency relations is made difficult by the limited availability of observations over certain, long horizons. We do not believe that this difficulty detracts from the importance of inference at low frequency, provided such inference is conducted carefully. Importantly for the purposes of this paper, however, here we do not solely focus on low-frequency dynamics. A crucial implication of our conceptual framework is, in fact, the existence of a tent-shaped behavior in R^2 values as a by-product of scale-wise predictability. Tent-shaped behavior, an implication of Proposition I, requires the dynamics at *all* frequencies to cooperate effectively, i.e., even at those high frequencies for which data availability would not be put forward as a statistical concern. In sum, we are not just drawing conclusions from frequencies associated with high statistical uncertainty, we are relying on *all* frequencies.

Standard economic theory views the presence of a market risk-return trade-off as compensation for variance risk. Given this logic, for past variance to affect future expected returns, higher past variance should predict higher future variance. Importantly, this is not the case when simply aggregating the raw data over the long run.¹⁶

However, at the scale over which we report predictability (i.e., the 8 to 16 year scale), we find a positive dependence between past values of the variance component and future values of the same component. In other words, consistent with an autoregressive (of order 1) specification of the components, the $j = 4$ variance component has a positive autocorrelation with uncorrelated residuals. The R^2 of the scale-wise autoregression on market variance is a rather substantial 43.28% with a positive slope of 0.12. As explained earlier, it is unsurprising to find a low *scale-wise* (for $j = 4$) autocorrelation. While the autocorrelation value appears small, we recall that it is a measure of correlation on the dilated time of a scale designed to capture economic fluctuations with 8-to-16 year cycles. As shown in the Online Supplement (see Section “Fitting an AR(1) process to the regressor”), the corresponding autocorrelation in calendar-time would naturally be higher.

Interestingly, the documented low dependence between past and future uncertainty dynamics at frequencies over which predictability applies differentiates our inferential problem from classical

¹⁶The corresponding results are available upon request.

assessments of predictability. High first-order autocorrelation of the predictor, in particular, has been put forward as a leading cause of inaccurate inference in predictability (e.g., Stambaugh (1999), Valkanov (2003), Lewellen (2004), Campbell and Yogo (2006), and Boudoukh et al. (2008)). In our framework, however, the autocorrelation ρ_{j^*} is scale-specific. At the low frequencies over which we identify scale-wise forecastability, ρ_{j^*} is estimated to be small.

In essence, we show that, at scale $j = 4$, a very slow-moving component of the uncertainty process predicts itself as well as the corresponding component in future excess market returns. Said differently, higher past values of the uncertainty component appear to predict higher future values of the same uncertainty component and, consequently, higher future values of the corresponding component in excess market returns, as required by conventional logic behind compensations for uncertainty (i.e., variance) risk. While this logic applies to a specific level of resolution in our framework, it translates - upon forward/backward aggregation - into predictability for long-run returns as shown formally in Proposition I and in the data. Extensive simulations from a data generating process allowing for scale-wise predictability provide further support.

8 Scale-wise risk-return trade-offs in equilibrium

The structure of the model follows the long-run risk model of Bansal and Yaron (2004) duly modified to account for heterogeneity in persistence as in Ortu et al. (2013). We introduce component-wise representations for both consumption growth and dividend growth. However, we shut down long-run risk and set the expectation of both growth rates equal to zero. As in Bollerslev, Tauchen, and Zhou (2009), who also make this assumption in their model specification, this is solely done to simplify matters and focus on the substantive core of our analysis, namely the role of scales and their equilibrium implications.

We begin with component-wise decompositions for log consumption growth, g_t , and log dividend growth, gd_t , taking the form:

$$g_t = \sum_{j=1}^J g_t^{(j)}, \quad gd_t = \sum_{j=1}^J gd_t^{(j)}.$$

We assume that each component of the consumption growth process, $g_t^{(j)}$, and of the dividend growth process, $gd_t^{(j)}$, is driven by its own scale-specific stochastic variance, $\sigma_t^{(j)2}$, i.e.,

$$g_{t+2j}^{(j)} = \sigma_t^{(j)} e_{g,t+2j}^{(j)}, \quad \text{where } e_{g,t+2j}^{(j)} \sim N(0, 1) \quad (19)$$

$$gd_{t+2j}^{(j)} = \varphi_d^{(j)} \sigma_t^{(j)} e_{gd,t+2j}^{(j)}, \quad \text{where } e_{gd,t+2j}^{(j)} \sim N(0, 1) \quad (20)$$

with the shocks $e_{g,t+2j}^{(j)}$ and $e_{gd,t+2j}^{(j')}$ being correlated for $j = j'$ (with scale-specific correlation denoted by $\rho_{dg}^{(j)}$) and uncorrelated otherwise.¹⁷ For parsimony, as in Bansal and Yaron (2004), the variance of consumption growth and the variance of dividend growth are driven by a common time-varying component (the terms $\varphi_d^{(j)}$ are, therefore, just scaling factors).

To complete the dynamics of the model, we assume that each of the stochastic variance components $\left\{ \sigma_t^{(j)2} \right\}_{j=1}^J$ follows an autoregressive process, i.e.,

$$\sigma_{t+2j}^{(j)2} = \rho_j \sigma_t^{(j)2} + \varepsilon_{t+2j}^{(j)}, \quad \text{with } \varepsilon_{t+2j}^{(j)} \sim N\left(0, \sigma^{(j)2}\right). \quad (21)$$

We operate in a pure exchange economy with a representative agent endowed with Epstein-Zin recursive preferences. The representative agent's Euler equation is well known:

$$\mathbb{E}_t \left[e^{m_{t+1} + r_{t+1}^i} \right] = 1, \quad (22)$$

where m_{t+1} is the log stochastic discount factor given by

$$m_{t+1} = \theta \log \beta - \frac{\theta}{\psi} g_{t+1} + (\theta - 1) r_{t+1}^a, \quad (23)$$

r_{t+1}^a is the (log) return on an asset yielding a dividend equal to aggregate consumption and r_{t+1}^i is the (log) return on any asset i . The parameter β denotes the discount factor. The preference

¹⁷If, as in Section 3, we were to assume that the scale- j shocks are DHTs of white noise shocks, then $e_{g,t}^{(j)}$ and $e_{g,s}^{(j)}$ would be uncorrelated for all t and s so that $|t - s| > 2^j - 1$. The same would, of course, apply to $e_{gd,t}^{(j)}$ and $e_{gd,s}^{(j)}$. In addition, assuming that the scale- j shocks are DHTs of white noise shocks implies that the components of the variance of consumption growth coincide with the variances of the components of the consumption growth process (see the Online Supplement for a proof). Since we use the former (i.e., the components of the variance of the consumption growth process) in Section 7 and we directly model the latter (i.e., the variances of the components of the consumption growth process) in this section, this equivalence is interesting.

parameter ψ measures the intertemporal elasticity of substitution (IES), γ measures risk aversion and $\theta = (1 - \gamma) / (1 - 1/\psi)$.

We now discuss the market risk premium.¹⁸ Recall, first, that the standard Campbell and Shiller (1988) log-linear approximation gives:

$$r_{a,t+1} = \kappa_0 + \kappa_1 z_{t+1}^a - z_t^a + g_{t+1}, \quad (24)$$

$$r_{m,t+1} = \kappa_{0,m} + \kappa_{1,m} z_{t+1}^m - z_t^m + g d_{t+1},$$

where z_t^a , z_t^m , denote the log price-consumption ratio and the log price-dividend ratio, respectively.

Because of our decompositions of consumption growth and dividend growth into components with different levels of persistence, representing by $z_{a,t}^{(j)}$, $z_{m,t}^{(j)}$ the components with persistence j of the (log) price-consumption ratio and the (log) price-dividend ratio, respectively, it is natural to conjecture that there exists a *component-by-component* linear relation between the financial ratios and the state variables $\sigma_t^{(j)2}$, i.e.,

$$z_{a,t}^{(j)} = A_{0,j} + A_j \sigma_t^{(j)2}, \quad (25)$$

$$z_{m,t}^{(j)} = A_{0,j}^m + A_j^m \sigma_t^{(j)2}.$$

As long as A_j and A_j^m are not zero, these relations tell us that the variation in the valuation ratios can be attributed to fluctuations in economic uncertainty.

The values of $A_{0,j}$, A_j , $A_{0,j}^m$, A_j^m in terms of the parameters of the model are obtained from the Euler condition in Eq. (22) after expressing the log stochastic discount factors and the returns in terms of the factors $\{\sigma_t^{(j)2}\}_{j=1}^J$ and the innovations $\{e_{g,t+2j}^{(j)}\}_{j=1}^J$ and $\{\varepsilon_{t+2j}^{(j)}\}_{j=1}^J$. One can, in fact, show that, plugging these expressions into the Euler equation and using the method of undetermined coefficients, one obtains a set of equations for the coefficients $A_{0,j}$, A_j , $A_{0,j}^m$, A_j^m whose solutions are

¹⁸Derivations and technical details are contained in the Online Supplement.

given by the following vectors of sensitivities:

$$\begin{aligned}\underline{A} &= 0.5 \frac{\left(\theta - \frac{\theta}{\psi}\right)^2}{\theta} \underline{1} (\mathbb{I}_J - \kappa_1 M)^{-1}, \\ \underline{A}_m &= \left(\frac{H_m}{2} - .5(1 - \gamma) \left(\frac{1}{\psi} - \gamma \right), \right) (\mathbb{I}_J - \kappa_{1,m} M)^{-1},\end{aligned}$$

where $M \equiv -diag(\rho_1, \dots, \rho_J)$, $\lambda_g \equiv \left(\frac{\theta}{\psi} - \theta + 1\right)$, $\underline{\lambda}_\epsilon \equiv \kappa_1(1 - \theta) \underline{A}$, $H_m \equiv \left[\lambda_g^2 + \varphi_d^{(1)2}, \dots, \lambda_g^2 + \varphi_d^{(J)2}\right]$ and \underline{A} , \underline{A}_m denote the row vectors with entries, A_1, \dots, A_J , A_1^m, \dots, A_J^m , respectively.

Two features of this model specification are noteworthy. First, if the IES and risk aversion are larger than 1, then θ is negative, and a volatility increase lowers the price-consumption ratio. Similarly, an increase in economic uncertainty will make consumption more volatile, thereby lowering asset valuations and increasing the risk premia on all assets. This highlights that a IES larger than 1 is critical for capturing the negative correlation between the price-dividend ratio and consumption variance. Second, an increase in the persistence of the variance shocks, namely M , magnifies their effects on valuation ratios, as changes in economic uncertainty are perceived as being long lasting.

Now, we can show that the innovation in the returns' component at level of persistence j are given by:

$$\begin{aligned}r_{a,t+2j}^{(j)} - \mathbb{E}_t[r_{a,t+2j}^{(j)}] &= \sigma_t^{(j)} e_{g,t+2j}^{(j)} + \kappa_1 A_j \varepsilon_{t+2j}^{(j)}, \\ r_{m,t+2j}^{(j)} - \mathbb{E}_t[r_{m,t+2j}^{(j)}] &= \varphi_d^{(j)} \sigma_t^{(j)} e_{d,t+2j}^{(j)} + \kappa_{1,m} A_j^m \varepsilon_{t+2j}^{(j)}.\end{aligned}$$

Similarly, the innovations in the stochastic discount factor's components are given by:

$$m_{t+2j}^{(j)} - \mathbb{E}_t[m_{t+2j}^{(j)}] = -\lambda_g \sigma_t^{(j)} e_{g,t+2j}^{(j)} - \underline{\lambda}_\epsilon^{(j)} \varepsilon_{t+2j}^{(j)}. \quad (26)$$

Given scale-wise risk premia equal to $\mathbb{E}_t[r_{m,t+2j}^{(j)} - r_{f,t+2j}^{(j)}] + 0.5\sigma_{r_{m,t+2j}^{(j)}}^2 = -\text{Cov}_t(m_{t+2j}^{(j)}, r_{i,t+2j}^{(j)})$, this leads to:

$$\mathbb{E}_t[r_{m,t+2j}^{(j)} - r_{f,t+2j}^{(j)}] + 0.5\sigma_{r_{m,t+2j}^{(j)}}^2 = \lambda_g \varphi_d^{(j)} \sigma_t^{(j)2} \rho_{dg}^{(j)} + \kappa_{1,m} [\underline{\lambda}_\epsilon \mathbf{Q}]_j A_j^m, \quad (27)$$

where

$$\boldsymbol{\varepsilon}_{t+1}^T \equiv [\varepsilon_{t+1}^{(1)}, \dots, \varepsilon_{t+1}^{(J)}], \quad \mathbf{Q} \equiv \mathbb{E}_t [\boldsymbol{\varepsilon}_{t+1} \boldsymbol{\varepsilon}_{t+1}^T].$$

Eq. (27) represents a risk-return trade-off on the generic scale j obtained as the result of a simple equilibrium argument based on a classical change of measure (in Eq. (23)) and scale-wise decompositions for consumption growth and dividend growth. Section 2 and Section 7 provided empirical content to it.

9 Conclusions and further discussion

Economic relations may apply to individual low-frequency components of the series of interest and be hidden by transient effects at high frequencies. To capture this idea parsimoniously, this paper models stock market returns and their predictors (economic uncertainty, in our case) as aggregates of uncorrelated components operating over different frequencies and introduces a notion of *scale-specific predictability*, i.e., predictability on the components.

We conclude with two observations. First, the use of variance as a predictor of asset returns is appealing in our framework because the corresponding backward-aggregated measure does not lose its economic interpretation. Backward-aggregated variance has, in fact, a natural interpretation in terms of long-run past variance. Having made this point, all alternative predictors, like the dividend-yield and other financial ratios, may also be employed. While their long-run past averages are not as easily interpretable, the role played by aggregation in the extraction of low-frequency information contained in the components and its translation into long-run properties of the data, as laid out in Proposition I, apply generally. To the extent that market return data and the dividend-yield - for instance - contain relevant information about long-run cash-flow and discount rate risks, regressions on their components and on properly-aggregated data appear very well-suited to uncover this information.

Some have argued that the dividend yield may be nonstationary (e.g., Cai and Wang, 2014,

and the references therein).¹⁹ Our decompositions apply to all processes for which a classical Wold representation apply and, therefore, to all covariance-stationary process. Thus, even though extreme forms of persistence can be accommodated, unit root type behavior would fall outside of our framework. However, given a Beveridge-Nelson decomposition of the dividend yield (dp) into a transitory (covariance-stationary) component and a stochastic trend, i.e.,

$$dp_t = x_{dp,t} + \psi(1) \sum_{k=0}^{\infty} \epsilon_{t-k},$$

the extended Wold representation will still be valid for the transitory component $x_{dp,t}$. In particular, when interpreting the stochastic trend ($\psi(1) \sum_{k=0}^{\infty} \epsilon_{t-k}$) as one of the components of the raw series dp , scale-wise predictability may apply to it as well as to individual components of the transitory component $x_{dp,t}$. We leave this issue for future work.

Second, we have shown that a classical predictive system is not incompatible with scale-specific predictability. On the contrary, it imposes predictability at all scales with tight parametrizations on the predictive slopes. Our data generating process, instead, frees up predictability at different scales and reconstructs the original series from unrestricted, yet parsimonious, scale-wise relations. An interesting question to ask is, then: what would be the *implied* nature of predictability on the raw series? Our conjecture is that one could obtain interesting predictive relations which go well beyond the typical autoregressive structure of order 1 of classical predictive systems and give an important role to *long lags*, something that appears to be a recurrent theme in recent macro-finance work (see, e.g., Fuster et al., 2010). A formalization of predictability on the raw series given scale-specific predictability represents the next step in this agenda.

¹⁹Nonstationarity in the dividend yield would induce nonstationarity in returns upon predictability and, therefore, raise economic issues. This said, well-accepted tests would generally fail to reject nonstationarity.

References

- Andersen, T. G., Bollerslev, T., 1997. Heterogeneous information arrivals and return volatility dynamics: Uncovering the long-run in high frequency returns. *Journal of Finance* 52 (3), 975–1005.
- Baker, S. R., Bloom, N., Davis, S. J., 2016. Measuring Economic Policy Uncertainty. *The Quarterly Journal of Economics* 131 (4), 1593–1636.
- Bandi, F. M., Perron, B., 2008. Long-run risk-return trade-offs. *Journal of Econometrics* 143 (2), 349–374.
- Bandi, F. M., Tamoni, A., 2016. The horizon of systematic risk: a new beta representation. SSRN eLibrary.
- Bansal, R., Khatchatrian, V., Yaron, A., 2005. Interpretable asset markets? *European Economic Review* 49 (3), 531 – 560.
- Baxter, M., King, R. G., 1999. Measuring business cycles: Approximate band-pass filters for economic time series. *The Review of Economics and Statistics* 81 (4), pp. 575–593.
- Beveridge, S., Nelson, C. R., 1981. A new approach to decomposition of economic time series into permanent and transitory components with particular attention to measurement of the ‘business cycle’. *Journal of Monetary Economics* 7 (2), 151–174.
- Bollerslev, T., Tauchen, G., Zhou, H., 2009. Expected stock returns and variance risk premia. *Review of Financial Studies* 22 (11), 4463–4492.
- Bonomo, M., Garcia, R., Meddahi, N., Tedongap, R., 2015. The long and the short of the risk-return trade-off. *Journal of Econometrics* 187 (2), 580–592.
- Boudoukh, J., Richardson, M., Whitelaw, R. F., 2008. The myth of long-horizon predictability. *Review of Financial Studies* 21 (4), 1577–1605.
- Cai, Z., Wang, Y., 2014. Testing predictive regression models with nonstationary regressors. *Journal of Econometrics* 178 (P1), 4–14.

- Calvet, L. E., Fisher, A. J., 2001. Forecasting multifractal volatility. *Journal of Econometrics* 105 (1), 27–58.
- Calvet, L. E., Fisher, A. J., 2007. Multifrequency news and stock returns. *Journal of Financial Economics* 86 (1), 178–212.
- Campbell, J. Y., Shiller, R., 1988. The dividend-price ratio and expectations of future dividends and discount factors. *Review of Financial Studies* 1 (3), 195–228.
- Campbell, J. Y., Vuolteenaho, T., 2004. Inflation illusion and stock prices. *American Economic Review* 94 (2), 19–23.
- Campbell, J. Y., Yogo, M., 2006. Efficient tests of stock return predictability. *Journal of Financial Economics* 81 (1), 27–60.
- Cerreia-Vioglio, S., Ortu, F., Severino, F., Tebaldi, C., 2017. Multivariate Wold Decompositions. Working Papers 606, IGIER, Bocconi University.
URL <https://ideas.repec.org/p/igi/igierp/606.html>
- Cochrane, J. H., 2008. The dog that did not bark: A defense of return predictability. *Review of Financial Studies* 21 (4), 1533–1575.
- Comin, D., Gertler, M., 2006. Medium-term business cycles. *American Economic Review* 96 (3), 523–551.
- Crowley, P. M., 2007. A Guide To Wavelets For Economists. *Journal of Economic Surveys* 21 (2), 207–267.
- Dijkerman, R., Mazumdar, R., 1994. Wavelet representations of stochastic processes and multiresolution stochastic models. *Signal Processing, IEEE Transactions on* 42 (7), 1640–1652.
- Fama, E. F., French, K. R., 1989. Business conditions and expected returns on stocks and bonds. *Journal of Financial Economics* 25 (1), 23–49.
- Fama, E. F., Schwert, G. W., 1977. Asset returns and inflation. *Journal of Financial Economics* 5 (2), 115–146.

- Fuster, A., Hebert, B., Laibson, D., 2010. Natural Expectations and Macroeconomic Fluctuations. *Journal of Economic Perspectives* 24 (4), 67–84.
- Gençay, R., Selçuk, F., Whitcher, B., 2001. *An Introduction to Wavelets and Other Filtering Methods in Finance and Economics*, 1st Edition. Academic Press, New York.
- Granger, C. W. J., Newbold, P., 1974. Spurious regressions in econometrics. *Journal of Econometrics* 2 (2), 111–120.
- Hansen, L. P., Scheinkman, J. A., 2009. Long-term risk: An operator approach. *Econometrica* 77 (1), 177–234.
- Kelly, B., Pruitt, S., 2013. Market expectations in the cross-section of present values. *The Journal of Finance* 68 (5), 1721–1756.
- Lamont, O., 1998. Earnings and expected returns. *The Journal of Finance* 53 (5), pp. 1563–1587.
- Lettau, M., Ludvigson, S., 2001. Consumption, aggregate wealth, and expected stock returns. *Journal of Finance* 56 (3), 815–849.
- Lewellen, J., 2004. Predicting returns with financial ratios. *Journal of Financial Economics* 74 (2), 209–235.
- Mallat, S. G., 1989. A theory for multiresolution signal decomposition: The wavelet representation. *IEEE Trans. Pattern Anal. Mach. Intell.* 11, 674–693.
- Menzly, L., Santos, T., Veronesi, P., 2004. Understanding predictability. *Journal of Political Economy* 112 (1), 1–47.
- Muller, U. A., Dacorogna, M. M., Dave, R. D., Olsen, R. B., Pictet, O. V., von Weizsacker, J. E., 1997. Volatilities of different time resolutions – analyzing the dynamics of market components. *Journal of Empirical Finance* 4 (2), 213–239.
- Müller, U. K., Watson, M. W., 2008. Testing models of low-frequency variability. *Econometrica* 76 (5), 979–1016.

- Nelson, C. R., 1976. Inflation and rates of return on common stocks. *The Journal of Finance* 31 (2), 471–483.
- Ortu, F., Severino, F., Tamoni, A., Tebaldi, C., 2017. A persistence-based wold-type decomposition for stationary time series. SSRN eLibrary.
- Ortu, F., Tamoni, A., Tebaldi, C., 2013. Long-run risk and the persistence of consumption shocks. *Review of Financial Studies* 26 (11), 2876–2915.
- Pastor, L., Stambaugh, R. F., 2009. Predictive systems: Living with imperfect predictors. *Journal of Finance* 64 (4), 1583–1628.
- Percival, D. B., Walden, A. T., 2000. *Wavelet Methods for Time Series Analysis* (Cambridge Series in Statistical and Probabilistic Mathematics). Cambridge University Press.
- Phillips, P. C. B., 1986. Understanding spurious regressions in econometrics. *Journal of Econometrics* 33 (3), 311–340.
- Ramsey, J. B., 1999. The contribution of wavelets to the analysis of economic and financial data. *Phil. Trans. R. Soc. Lond. A* 357 (1760), 2593–2606.
- Sizova, N., 2013. Long-horizon return regressions with historical volatility and other long-memory variables. *Journal of Business and Economic Statistics* 31 (4), 546–559.
- Stambaugh, R. F., 1999. Predictive regressions. *Journal of Financial Economics* 54 (3), 375–421.
- Tamoni, A., 2011. The multi-horizon dynamics of risk and returns. SSRN eLibrary.
- Valkanov, R., 2003. Long horizon regressions: Theoretical results and applications. *Journal of Financial Economics* 68, 201–232.
- Welch, I., Goyal, A., 2008. A comprehensive look at the empirical performance of equity premium prediction. *Review of Financial Studies* 21 (4), 1455–1508.

Panel A1: $(r_{t+1,t+h} - r f_t) = \alpha_h + \beta_h v_{t-h+1,t} + \epsilon_{t+h}$

	Horizon h (in years)									
	1	2	3	4	5	6	7	8	9	10
$v_{t-h+1,t}$	0.42 (0.86) {0.07}	0.55 (0.83) {0.11}	0.12 (0.19) {0.03}	0.32 (0.44) {0.07}	0.89 (1.28) {0.20}	1.10 (1.36) {0.25}	1.35 (1.60) {0.03}	1.86 (2.24) {0.41}	2.80 (3.68) {0.62*}	3.36 (5.02) {0.74*}
$R^2(\%)$ [5 th , 95 th]	[0.52] [0.00, 3.74]	[1.18] [0.01, 7.10]	[0.08] [0.01, 8.17]	[0.54] [0.01, 9.64]	[3.82] [0.03, 20.92]	[5.82] [0.11, 32.60]	[8.70] [0.15, 38.30]	[14.75] [2.27, 50.55]	[28.17] [11.24, 61.52]	[36.12] [20.81, 65.43]

Panel A2: $(r_{t+1,t+h} - r f_t) = \alpha_h + \beta_h v_{t-h+1,t} + \epsilon_{t+h}$

	Horizon h (in years)									
	11	12	13	14	15	16	17	18	19	20
$v_{t-h+1,t}$	3.79 (6.29) {0.84**}	3.94 (6.34) {0.91**}	3.93 (6.34) {0.95**}	3.79 (6.56) {0.99**}	3.71 (7.44) {1.00*}	3.70 (8.61) {1.09**}	3.11 (7.57) {0.90}	2.41 (5.73) {0.65}	1.38 (3.69) {0.34}	0.77 (2.08) {0.19}
$R^2(\%)$ [5 th , 95 th]	[42.22] [24.59, 67.43]	[46.35] [26.70, 73.01]	[48.46] [31.12, 77.42]	[50.44] [36.46, 78.03]	[51.11] [35.45, 76.56]	[55.24] [33.64, 83.37]	[45.83] [21.67, 86.63]	[30.64] [7.48, 76.41]	[10.62] [0.65, 48.81]	[3.50] [0.11, 36.72]

Panel B: $r_{k2^j+2^j}^{(j)} - r f_{k2^j+2^j}^{(j)} = \beta_j v_{k2^j}^{(j)} + \epsilon_{k2^j+2^j}$

	Time-scale j		
	1	2	3
$v_t^{(j)}$	-0.68 (-0.69)	0.87 (0.43)	-0.34 (-0.26)
$R^2(\%)$	[1.16]	[0.96]	[0.77]

Table 1: Market risk. Panel A: We run linear regressions of h -period continuously compounded market returns on the CRSP value-weighted index in excess of a 1-year Treasury bill rate on h -period past market variance, $v_{t-h,t}$. For each regression, the table reports OLS estimates of the regressors, Newey-West t -statistics with 2 * (horizon - 1) lags (in parentheses), the t/\sqrt{T} test suggested in Valkanov (2003) (in curly brackets), and R^2 (in square brackets). Ninety percent confidence intervals for the true R^2 's are reported in brackets below the sample values. Significance at the 5%, 2.5%, and 1% level of the t/\sqrt{T} test using Valkanov's (2003) critical values is indicated by *, **, and ***, respectively. **Panel B:** Component-wise predictive regressions of the components of excess stock market returns on the components of market variance. For each regression (in Panel B), the table reports OLS estimates of the regressors, t -stats in parentheses and R^2 statistics (and bootstrapped confidence intervals) in square brackets. The sample is annual and spans the period 1930-2014.

Panel A1: $(r_{t+1,t+h} - r f_t) = \alpha_h + \beta_h v_{t-h+1,t} + \epsilon_{t+h}$

	Horizon h (in years)									
	1	2	3	4	5	6	7	8	9	10
$v_{t-h+1,t}$	1.80 (1.12) {0.14}	2.04 (2.31) {0.22}	0.50 (0.60) {0.07}	-0.04 (-0.05) {-0.01}	-0.19 (-0.17) {-0.03}	1.07 (0.89) {0.16}	2.71 (2.28) {0.45}	3.05 (2.31) {0.49}	3.89 (2.94) {0.62*}	5.00 (4.11) {0.84**}
$R^2(\%)$ [5 th , 95 th]	[1.94] [0.03, 8.54]	[4.83] [0.09, 15.12]	[0.48] [0.01, 10.74]	[0.00] [0.01, 15.78]	[0.08] [0.01, 26.83]	[2.70] [0.03, 32.83]	[17.44] [6.55, 44.24]	[19.78] [8.05, 49.43]	[28.43] [12.97, 56.32]	[42.18] [25.21, 61.04]

Panel A2: $(r_{t+1,t+h} - r f_t) = \alpha_h + \beta_h v_{t-h+1,t} + \epsilon_{t+h}$

	Horizon h (in years)									
	11	12	13	14	15	16	17	18	19	20
$v_{t-h+1,t}$	5.54 (4.44) {0.98***}	5.84 (5.20) {1.10***}	5.61 (5.32) {1.09***}	5.43 (6.80) {1.08***}	4.87 (8.01) {0.95*}	4.48 (9.34) {0.91*}	3.97 (9.29) {0.82}	3.14 (6.14) {0.63}	2.08 (3.38) {0.39}	0.93 (1.34) {0.18}
$R^2(\%)$ [5 th , 95 th]	[50.01] [32.65, 69.62]	[55.68] [38.15, 71.14]	[55.06] [38.18, 71.03]	[54.69] [31.30, 72.59]	[48.41] [19.49, 69.39]	[46.13] [11.88, 71.55]	[41.15] [4.85, 71.52]	[29.03] [0.76, 62.01]	[13.95] [0.42, 42.73]	[3.29] [0.12, 26.83]

Panel B: $r_{k2^j+2^j}^{(j)} - r f_{k2^j+2^j}^{(j)} = \beta_j v_{k2^j}^{(j)} + \epsilon_{k2^j+2^j}^{(j)}$

	Time-scale j		
	1	2	3
$v_t^{(j)}$	-6.01 (-1.24)	-9.64 (-1.55)	-3.24 (-0.85)
$R^2(\%)$	[3.70]	[25.83]	[7.41]
	[61.62]		

Table 2: Consumption risk. Panel A: We run linear regressions of h -period continuously compounded market returns on the CRSP value-weighted index in excess of a 1-year Treasury bill rate on h -period past consumption variance, $v_{t-h,t}$. For each regression, the table reports OLS estimates of the regressors, Newey-West t -statistics with 2 * (horizon - 1) lags (in parentheses), the t/\sqrt{T} test suggested in Valkanov (2003) (in curly brackets), and R^2 (in square brackets). Ninety percent confidence intervals for the true R^2 's are reported in brackets below the sample values. Significance at the 5%, 2.5%, and 1% level of the t/\sqrt{T} test using Valkanov's (2003) critical values is indicated by *, **, and ***, respectively. **Panel B:** Component-wise predictive regressions of the components of excess stock market returns on the components of consumption variance $v_{j,t}^2$. For each regression (in Panel B), the table reports OLS estimates of the regressors, t -stats in parentheses and R^2 statistics (and bootstrapped confidence intervals) in square brackets. The sample is annual and spans the period 1930-2014.

Panel A1: $(r_{t+1,t+h} - r f_t) = \alpha_h + \beta_h v_{t-h+1,t} + \epsilon_{t+h}$

	Horizon h (in years)									
	1	2	3	4	5	6	7	8	9	10
$v_{t-h+1,t}$	0.03 (1.94) {0.19}	0.02 (1.56) {0.17}	0.01 (1.05) {0.16}	0.01 (0.72) {0.13}	0.01 (0.75) {0.15}	0.01 (0.99) {0.21}	0.02 (1.64) {0.37}	0.03 (2.42) {0.52}	0.03 (3.07) {0.60*}	0.03 (3.51) {0.70*}
$R^2(\%)$ [5 th , 95 th]	[3.39] [0.08, 11.81]	[2.94] [0.04, 12.91]	[2.49] [0.03, 15.86]	[1.68] [0.02, 18.85]	[2.21] [0.02, 21.09]	[4.52] [0.05, 29.56]	[12.63] [0.29, 43.24]	[21.91] [2.65, 53.49]	[26.83] [4.05, 56.10]	[33.32] [5.81, 63.18]

Panel A2: $(r_{t+1,t+h} - r f_t) = \alpha_h + \beta_h v_{t-h+1,t} + \epsilon_{t+h}$

	Horizon h (in years)									
	11	12	13	14	15	16	17	18	19	20
$v_{t-h+1,t}$	0.04 (3.70) {0.78*}	0.04 (3.87) {0.90**}	0.04 (3.54) {0.95**}	0.04 (3.27) {1.00**}	0.03 (3.07) {1.00**}	0.03 (3.00) {1.04*}	0.03 (2.92) {1.00*}	0.03 (2.92) {0.88}	0.02 (3.00) {0.69}	0.02 (3.35) {0.52}
$R^2(\%)$ [5 th , 95 th]	[38.77] [7.85, 70.50]	[45.34] [13.59, 77.32]	[48.41] [15.11, 82.42]	[50.91] [15.50, 84.17]	[51.07] [13.68, 81.64]	[52.76] [15.51, 79.83]	[51.04] [12.93, 76.72]	[44.75] [9.15, 70.75]	[33.43] [4.27, 61.60]	[21.78] [1.12, 58.55]

Panel B: $r_{k2^j+2^j}^{(j)} - r f_{k2^j+2^j}^{(j)} = \beta_j v_{k2^j}^{(j)} + \epsilon_{k2^j+2^j}$

	Time-scale j		
	1	2	3
$v_t^{(j)}$	-0.02 (-0.74)	0.04 (1.11)	0.05 (1.65)
$R^2(\%)$	[1.39]	[6.04]	[23.32]
	[72.00]		

Table 3: Economic policy uncertainty (Baker, Bloom and Davis, 2015). **Panel A:** We run linear regressions of h -period continuously compounded market returns on the CRSP value-weighted index in excess of a 1-year Treasury bill rate on h -period past (squared) economic policy uncertainty (EPU), $v_{t-h,t}$. For each regression, the table reports OLS estimates of the regressors, Newey-West t -statistics with $2*$ (horizon -1) lags (in parentheses), the t/\sqrt{T} test suggested in Valkanov (2003) (in curly brackets), and R^2 (in square brackets). Ninety percent confidence intervals for the true R^2 s are reported in brackets below the sample values. Significance at the 5%, 2.5%, and 1% level of the t/\sqrt{T} test using Valkanov's (2003) critical values is indicated by *, **, and ***, respectively. **Panel B:** Component-wise predictive regressions of the components of excess stock market returns on the components of EPU. For each regression (in Panel B), the table reports OLS estimates of the regressors, t -stats in parentheses and R^2 statistics (and bootstrapped confidence intervals) in square brackets. The sample is annual and spans the period 1930-2014.

	Annual calendar time
Time-scale	Frequency resolution
$j = 1$	1 – 2 years
$j = 2$	2 – 4 years
$j = 3$	4 – 8 years
$j = 4$	8 – 16 years
$\pi_t^{(4)}$	> 16 years

Table 4: Interpretation of the time-scale (or persistence level) j in terms of time spans in the case of annual time series. Each scale corresponds to a frequency interval, or conversely an interval of periods, and thus each scale is associated with a range of time horizons.

Scales $j =$	Panel A: Market excess returns				Panel B: Consumption risk			
	1	2	3	4	1	2	3	4
1		-0.03 (0.09)	-0.04 (0.07)	0.09 (0.05)		0.33 (0.18)	0.17 (0.10)	-0.09 (0.09)
2			-0.13 (0.09)	0.15 (0.10)			-0.09 (0.26)	-0.12 (0.15)
3				0.14 (0.13)				-0.06 (0.13)

Scales $j =$	Panel C: Market risk				Panel D: Economic policy uncertainty			
	1	2	3	4	1	2	3	4
1		0.27 (0.12)	-0.13 (0.09)	0.00 (0.06)		0.08 (0.09)	0.09 (0.11)	0.06 (0.05)
2			-0.03 (0.15)	-0.00 (0.09)			0.22 (0.09)	0.09 (0.08)
3				0.33 (0.12)				0.32 (0.15)

Table 5: **Pairwise correlations.** We report the pair-wise correlations between the individual components of excess market returns (Panel A), consumption variance (Panel B), market variance (Panel C), and (squared) economic policy uncertainty (Panel D). The pair-wise correlations are obtained by using redundant data on the components rather than the decimated counterparts. Standard errors for the correlation between $x_t^{(j)}$ and $x_t^{(j')}$, $j \neq j'$, are Newey-West with $2^{\max(j,j')}$ lags.

Appendix

A Proof of Proposition I

We begin with some matrix manipulations. Consider the following component dynamics for $j = j^*$, where $j^* \in \{1, \dots, J\}$:

$$y_{t+2j}^{(j)} = \beta_j x_t^{(j)} \quad (\text{A.1})$$

$$x_{t+2j}^{(j)} = \rho_j x_t^{(j)} + \sigma_j \epsilon_{t+2j} \quad (\text{A.2})$$

For $j = 1, \dots, J$, with $j \neq j^*$, we have

$$y_t^{(j)} = 0, \quad (\text{A.3})$$

$$x_t^{(j)} = 0. \quad (\text{A.4})$$

Assume that $T = 16$, $j^* = 2$, and $J = 3$. Arrange the components of x as follows:

$$\begin{pmatrix} \pi_8^{(3)} & \pi_{16}^{(3)} \\ x_8^{(3)} & x_{16}^{(3)} \\ x_8^{(2)} & x_{16}^{(2)} \\ x_4^{(2)} & x_{12}^{(2)} \\ x_8^{(1)} & x_{16}^{(1)} \\ x_6^{(1)} & x_{14}^{(1)} \\ x_4^{(1)} & x_{12}^{(1)} \\ x_2^{(1)} & x_{10}^{(1)} \end{pmatrix} \quad (\text{A.5})$$

and, analogously, for the components of y . To reconstruct the time series x_t , we run through each column of the matrix in Eq. (A.5) and, for each column, we perform the following operation:

$$X_8^{(3)} = \begin{pmatrix} x_8 \\ x_7 \\ x_6 \\ x_5 \\ x_4 \\ x_3 \\ x_2 \\ x_1 \end{pmatrix} = (\mathcal{T}^{(3)})^{-1} \begin{pmatrix} \pi_8^{(3)} \\ x_8^{(3)} \\ x_8^{(2)} \\ x_4^{(2)} \\ x_8^{(1)} \\ x_6^{(1)} \\ x_4^{(1)} \\ x_2^{(1)} \end{pmatrix} \quad (\text{A.6})$$

and

$$X_{16}^{(3)} = \begin{pmatrix} x_{16} \\ x_{15} \\ x_{14} \\ x_{13} \\ x_{12} \\ x_{11} \\ x_{10} \\ x_9 \end{pmatrix} = (\mathcal{T}^{(3)})^{-1} \begin{pmatrix} \pi_{16}^{(3)} \\ x_{16}^{(3)} \\ x_{16}^{(2)} \\ x_{12}^{(2)} \\ x_{16}^{(1)} \\ x_{14}^{(1)} \\ x_{12}^{(1)} \\ x_{10}^{(1)} \end{pmatrix}. \quad (\text{A.7})$$

We do the same for the components of y_t . The matrix $(\mathcal{T}^{(3)})^{-1}$ takes the following form:

$$(\mathcal{T}^{(3)})^{-1} = \begin{pmatrix} \frac{1}{\sqrt{8}} & \frac{1}{\sqrt{8}} & \frac{1}{2} & 0 & \frac{1}{\sqrt{2}} & 0 & 0 & 0 \\ \frac{1}{\sqrt{8}} & \frac{1}{\sqrt{8}} & \frac{1}{2} & 0 & -\frac{1}{\sqrt{2}} & 0 & 0 & 0 \\ \frac{1}{\sqrt{8}} & \frac{1}{\sqrt{8}} & -\frac{1}{2} & 0 & 0 & \frac{1}{\sqrt{2}} & 0 & 0 \\ \frac{1}{\sqrt{8}} & \frac{1}{\sqrt{8}} & -\frac{1}{2} & 0 & 0 & -\frac{1}{\sqrt{2}} & 0 & 0 \\ \frac{1}{\sqrt{8}} & -\frac{1}{\sqrt{8}} & 0 & \frac{1}{2} & 0 & 0 & \frac{1}{\sqrt{2}} & 0 \\ \frac{1}{\sqrt{8}} & -\frac{1}{\sqrt{8}} & 0 & \frac{1}{2} & 0 & 0 & -\frac{1}{\sqrt{2}} & 0 \\ \frac{1}{\sqrt{8}} & -\frac{1}{\sqrt{8}} & 0 & -\frac{1}{2} & 0 & 0 & 0 & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{8}} & -\frac{1}{\sqrt{8}} & 0 & -\frac{1}{2} & 0 & 0 & 0 & -\frac{1}{\sqrt{2}} \end{pmatrix}. \quad (\text{A.8})$$

Using the dynamics of the state in Eq. (A.2), together with Eqs. (A.6) and (A.7), we obtain

$$X_{16}^{(3)} = \begin{pmatrix} x_{16} = x_{16}^{(2)}/2 \\ x_{15} = x_{16}^{(2)}/2 \\ x_{14} = -x_{16}^{(2)}/2 \\ x_{13} = -x_{16}^{(2)}/2 \\ x_{12} = x_{12}^{(2)}/2 \\ x_{11} = x_{12}^{(2)}/2 \\ x_{10} = -x_{12}^{(2)}/2 \\ x_9 = -x_{12}^{(2)}/2 \end{pmatrix} \quad \text{and} \quad X_8^{(3)} = \begin{pmatrix} x_8 = x_8^{(2)}/2 \\ x_7 = x_8^{(2)}/2 \\ x_6 = -x_8^{(2)}/2 \\ x_5 = -x_8^{(2)}/2 \\ x_4 = x_4^{(2)}/2 \\ x_3 = x_4^{(2)}/2 \\ x_2 = -x_4^{(2)}/2 \\ x_1 = -x_4^{(2)}/2 \end{pmatrix}. \quad (\text{A.9})$$

We now turn to forward/backward aggregation. Let us construct the temporally-aggregated series

$$y_{t+1,t+h} = \sum_{i=1}^h y_{t+i}$$

and run the forward/backward regression

$$y_{t+1,t+h} = \tilde{\beta} x_{t-h+1,t} + \epsilon_{t+1,t+h},$$

where $x_{t+1,t+h}$ is defined like $y_{t+1,t+h}$. For $h = 4$, and using Eqs. (A.1) and (A.3) together with Eq. (A.9),

we have

$$\begin{aligned}
y_{13,16} &= 0 & x_{13,16} &= 0 \\
y_{12,15} &= (-y_{16}^{(2)} + y_{12}^{(2)})/2 = \beta \left(-x_{12}^{(2)} + x_8^{(2)} \right) / 2 & x_{12,15} &= (-x_{16}^{(2)} + x_{12}^{(2)})/2 \\
y_{11,14} &= -y_{16}^{(2)} + y_{12}^{(2)} = \beta \left(-x_{12}^{(2)} + x_8^{(2)} \right) & x_{11,14} &= -x_{16}^{(2)} + x_{12}^{(2)} \\
y_{10,13} &= (-y_{16}^{(2)} + y_{12}^{(2)})/2 = \beta \left(-x_{12}^{(2)} + x_8^{(2)} \right) / 2 & x_{10,13} &= (-x_{16}^{(2)} + x_{12}^{(2)})/2 \\
y_{9,12} &= 0 & x_{9,12} &= 0 \\
y_{8,11} &= (-y_{12}^{(2)} + y_8^{(2)})/2 = \beta \left(-x_8^{(2)} + x_4^{(2)} \right) / 2 & x_{8,11} &= (-x_{12}^{(2)} + x_8^{(2)})/2 \\
y_{7,10} &= -y_{12}^{(2)} + y_8^{(2)} = \beta \left(-x_8^{(2)} + x_4^{(2)} \right) & x_{7,10} &= -x_{12}^{(2)} + x_8^{(2)} \\
y_{6,9} &= (-y_{12}^{(2)} + y_8^{(2)})/2 = \beta \left(-x_8^{(2)} + x_4^{(2)} \right) / 2 & x_{6,9} &= (-x_{12}^{(2)} + x_8^{(2)})/2 \\
y_{5,8} &= 0 & x_{5,8} &= 0 \\
y_{4,7} &= (-y_8^{(2)} + y_4^{(2)})/2 = \beta \left(-x_4^{(2)} + x_0^{(2)} \right) / 2 & x_{4,7} &= (-x_8^{(2)} + x_4^{(2)})/2 \\
y_{3,6} &= -y_8^{(2)} + y_4^{(2)} = \beta \left(-x_4^{(2)} + x_0^{(2)} \right) & x_{3,6} &= -x_8^{(2)} + x_4^{(2)} \\
y_{2,5} &= (-y_8^{(2)} + y_4^{(2)})/2 = \beta \left(-x_4^{(2)} + x_0^{(2)} \right) / 2 & x_{2,5} &= (-x_8^{(2)} + x_4^{(2)})/2 \\
y_{1,4} &= 0 & x_{1,4} &= 0.
\end{aligned}$$

Thus, regressing $y_{t+1,t+4}$ on $x_{t-3,t}$ yields $\tilde{\beta} = \beta$ with $R^2 = 100\%$. Hence, when scale-wise predictability applies to a scale operating between 2^{j^*-1} and 2^j , maximum predictability upon two-way aggregation arises over an horizon $h = 2^{j^*}$. In our case, $j^* = 2$ and $h = 4$. Consider, for example, an alternative aggregation level: $h = 2$. We have

$$\begin{aligned}
y_{15,16} &= y_{16}^{(2)} = \beta x_{12}^{(2)} & x_{15,16} &= x_{16}^{(2)} \\
y_{14,15} &= 0 & x_{14,15} &= 0 \\
y_{13,14} &= -y_{16}^{(2)} = -\beta x_{12}^{(2)} & x_{13,14} &= -x_{16}^{(2)} \\
y_{12,13} &= (-y_{16}^{(2)} + y_{12}^{(2)})/2 = \beta(-x_{12}^{(2)} + x_8^{(2)})/2 & x_{12,13} &= (-x_{16}^{(2)} + x_{12}^{(2)})/2 \\
y_{11,12} &= y_{12}^{(2)} = \beta x_8^{(2)} & x_{11,12} &= x_{12}^{(2)} \\
y_{10,11} &= 0 & x_{10,11} &= 0 \\
y_{9,10} &= -y_{12}^{(2)} = -\beta x_8^{(2)} & x_{9,10} &= -x_{12}^{(2)} \\
y_{8,9} &= (-y_{12}^{(2)} + y_8^{(2)})/2 = \beta(-x_8^{(2)} + x_4^{(2)})/2 & x_{8,9} &= (-x_{12}^{(2)} + x_8^{(2)})/2 \\
y_{7,8} &= y_8^{(2)} = \beta x_4^{(2)} & x_{7,8} &= x_8^{(2)} \\
y_{6,7} &= 0 & x_{6,7} &= 0 \\
y_{5,6} &= -y_8^{(2)} = -\beta x_4^{(2)} & x_{5,6} &= -x_8^{(2)} \\
y_{4,5} &= (-y_8^{(2)} + y_4^{(2)})/2 = \beta(-x_4^{(2)} + x_0^{(2)})/2 & x_{4,5} &= (-x_8^{(2)} + x_4^{(2)})/2 \\
y_{3,4} &= y_4^{(2)} = \beta x_0^{(2)} & x_{3,4} &= x_4^{(2)} \\
y_{2,3} &= 0 & x_{2,3} &= 0 \\
y_{1,2} &= -y_4^{(2)} = -\beta x_0^{(2)} & x_{1,2} &= -x_4^{(2)},
\end{aligned}$$

where we use the implied dynamics for x , see equations (A.9), and the equivalent ones for y together with (A.1) and (A.2). The regression of $y_{t+1,t+2}$ on $x_{t-1,t}$ yields (based on a fundamental block of four elements):

$$\begin{aligned}\tilde{\beta} &= \frac{Cov(y_{15,16}, x_{13,14}) + Cov(y_{13,14}, x_{11,12})}{Var(x_{10,11}) + Var(x_{11,12}) + Var(x_{12,13}) + Var(x_{13,14})} \\ &= \frac{-\beta Var(x_{12}^{(2)}) \rho - \beta Var(x_{12}^{(2)})}{Var(x_{12}^{(2)}) + Var\left(\frac{x_{16}^{(2)}}{2}\right) + Var\left(\frac{x_{12}^{(2)}}{2}\right) - \frac{Cov(x_{16}^{(2)}, x_{12}^{(2)})}{2} + Var(x_{16}^{(2)})} \\ &= -2\beta \frac{(1 + \rho_j)}{5 - \rho_j}\end{aligned}$$

and, hence, an inconsistent slope estimate. This estimate could have a changed sign (with respect to β) and be drastically attenuated. In fact, $\tilde{\beta} = 0$ if $\rho_j = -1$ and $\tilde{\beta} = -\beta$ if $\rho_j = 1$.

B Data

The empirical analysis in Sections 2, is conducted using annual data on consumption and stock returns from 1930 to 2014. We take the view that this sample is the most representative of the overall high/medium/low-frequency variation in asset prices and macroeconomic data.

Aggregate consumption is from the Bureau of Economic Analysis, series 7.1, and is defined as consumer expenditures on non-durables and services. Our measure of consumption volatility is based on modeling consumption growth as following an AR(1) with an error variance evolving as an heterogeneous ARCH model (see Muller et al. (1997)). The HARCh dynamics accommodates numerous heterogeneous information arrival processes, see, e.g., Andersen and Bollerslev (1997). Similar results are obtained by modeling consumption growth as following an AR(1)-GARCh(1,1), as in Bansal et al. (2005).

We use the NYSE/Amex value-weighted index with dividends as our market proxy, R_{t+1} . Return data on the value-weighted market index are obtained from the Chicago Center for Research in Security Prices (CRSP). The annual return series is constructed from monthly data under the assumption of reinvestment at a zero-rate. The nominal short-term rate ($R_{f,t+1}$) is the yield on the 1-year Treasury bill.

The h -horizon continuously-compounded excess market return is calculated as $r_{t+1,t+h} = r_{t+1}^e + \dots + r_{t+h}^e$, where $r_{t+j}^e = \ln(R_{t+j}) - \ln(R_{f,t+j})$ is the 1-year excess logarithmic market return between dates $t + j - 1$ and $t + j$, R_{t+j} is the simple gross market return, and $R_{f,t+j}$ is the gross risk-free rate.

The market's realized variance between the end of period t and the end of period $t + n$, a measure of integrated volatility, is obtained by computing

$$v_{t,t+n}^2 = \sum_{d=t_1}^{t_D} r_d^2,$$

where $[t_1, t_D]$ denotes the sample of available daily returns between the end of period t and the end of period $t + n$, and r_d is the market's logarithmic return on day d .

The measure of economic policy uncertainty (EPU) is based on Baker et al. (2016).