

Jiwook Jang, [Angelos Dassios](#), Hongbiao Zhao
Moments of renewal shot-noise processes
and their applications

Article (Accepted version)
(Refereed)

Original citation:

Jang, Jiwook and Dassios, Angelos and Zhao, Hongbiao (2018) *Moments of renewal shot-noise processes and their applications*. [Scandinavian Actuarial Journal](#). ISSN 0346-1238

DOI: [10.1080/03461238.2018.1452285](https://doi.org/10.1080/03461238.2018.1452285)

© 2018 [Informa UK Limited, trading as Taylor & Francis Group](#)

This version available at: <http://eprints.lse.ac.uk/87428/>

Available in LSE Research Online: April 2018

LSE has developed LSE Research Online so that users may access research output of the School. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LSE Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain. You may freely distribute the URL (<http://eprints.lse.ac.uk>) of the LSE Research Online website.

This document is the author's final accepted version of the journal article. There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

Moments of Renewal Shot-noise Processes and their Applications*

Jiwook Jang[†]

Macquarie University

Angelos Dassios[‡]

London School of Economics

Hongbiao Zhao[§]

Shanghai University of Finance and Economics

12th March 2018

Abstract

In this paper, we study the family of renewal shot-noise processes. The Feynmann-Kac formula is obtained based on the piecewise deterministic Markov process theory and the martingale methodology. We then derive the Laplace transforms of the conditional moments and asymptotic moments of the processes. In general, by inverting the Laplace transforms, the asymptotic moments and the first conditional moments can be derived explicitly, however, other conditional moments may need to be estimated numerically. As an example, we develop a very efficient and general algorithm of Monte Carlo exact simulation for estimating the second conditional moments. The results can be then easily transformed to the counterparts of discounted aggregate claims for insurance applications, and we apply the first two conditional moments for the actuarial net premium calculation. Similarly, they can also be applied to credit risk and reliability modelling. Numerical examples with four distribution choices for interarrival times are provided to illustrate how the models can be implemented.

Keywords: Renewal shot-noise processes; Discounted aggregate claims; Actuarial net premium; Piecewise-deterministic Markov processes; Martingale method; Monte Carlo exact simulation; Credit risk; Reliability

Mathematics Subject Classification (2010): Primary: 91B30; Secondary: 60G55; 60K20

* An edited version to be published by *Scandinavian Actuarial Journal*

[†]Department of Applied Finance and Actuarial Studies, Faculty of Business and Economics, Macquarie University, Sydney NSW 2109, Australia, E-mail: jiwook.jang@mq.edu.au

[‡]Department of Statistics, London School of Economics, Houghton Street, London WC2A 2AE, United Kingdom. Email: a.dassios@lse.ac.uk

[§]*Corresponding author*, School of Statistics and Management, Shanghai University of Finance and Economics, No. 777 Guoding Road, Shanghai 200433, China; Shanghai Institute of International Finance and Economics, No. 777 Guoding Road, Shanghai 200433, China. Email: h.zhao1@lse.ac.uk

1 Introduction

Since the beginning of the 20th century, *shot-noise processes* have been extensively used to model a very wide variety of natural phenomena, with numerous applications in electronics, optics, biology and many other fields in natural science, see early literature in Campbell (1909a,b), Schottky (1918), Picinbono et al. (1970) and Verveen and DeFelice (1974). More recent applications extended to insurance and actuarial science in particular can be found in Klüppelberg and Mikosch (1995), Brémaud (2000), Dassios and Jang (2003, 2005), Jang (2004), Jang and Kravavych (2004), Torrisi (2004), Albrecher and Asmussen (2006), Macci and Torrisi (2011), Zhu (2013) and Schmidt (2014). Mostly, they adopted the classical *Poisson shot-noise process* (Cox and Isham, 1980, p.88), where the arrivals of claims are simply assumed to follow a Poisson process. However, an exponential distribution could be not appropriate for modelling claim interarrival times in practice when the likelihood of a claim given the time elapsed since the previous one is not constant over time. There has been a significant volume of literature that questions the appropriateness of a Poisson process in insurance modelling (Seal, 1983; Beard et al., 1984) such as the rainfall modelling (Cox and Isham, 1980; Smith, 1980). For catastrophic events (e.g. floods, storms, hails, bushfires, earthquakes and terrorist attacks), the assumption that resulting claims occur in terms of a Poisson process is inadequate as it has a deterministic intensity, i.e., it has the same claim frequency rate between the same time interval of duration.

A natural generalisation of Poisson process is the family of renewal processes (Cox, 1962; Cox and Miller, 1965; Karlin and Taylor, 1975; Grandell, 1991; Ross, 1996; Rolski et al., 2008), which could offer more flexible model choices and are versatile enough to capture different styles of claim interarrival times in reality. Using ordinary, delayed and stationary renewal processes to derive the moments and moment generating functions of compound renewal sums with discounted claims can be found in Lévêillé and Garrido (2001a,b) and Lévêillé et al. (2010). Since Andersen (1957) proposed to use a compound renewal risk model and Gerber and Shiu (1998) introduced the so-called discounted penalty function, the delayed and stationary renewal risk models and their extensions for modelling insurers' surplus processes can also be noticed in Willmot and Dickson (2003), Gerber and Shiu (2005), Li and Garrido (2005), Willmot (2007) and Woo (2010).

In this paper, we mainly study *renewal shot-noise processes*, the generalised family of Poisson shot-noise process. They are shot-noise processes driven by ordinary renewal processes, so that the interarrival times could be any positive independent identically distributed random variables. This paper can be considered as the generalisation of Jang (2004) from the classical Poisson process

to a rather general renewal process for the underlying point process. However, this generalisation is technically nontrivial, since the renewal components lead our new models beyond the affine framework in general, and several new approaches have been adopted or developed to investigate the properties of moments. Based on the *piecewise deterministic Markov process theory* (Davis, 1984, 1993) and the martingale methodology (Dassios and Embrechts, 1989), we first obtain the Feynmann-Kac formula. We then derive the Laplace transforms of the conditional moments and asymptotic moments of the processes. In general, by inverting the Laplace transforms of these moments, any asymptotic moments as well as the first conditional moments can be derived explicitly, however, other conditional moments may need to be estimated numerically. As an example, we develop a very efficient and general algorithm of Monte Carlo exact simulation for estimating the second conditional moments. The results then can be easily transformed to the counterparts of discounted aggregate claims in insurance, and we apply the first two conditional moments for the actuarial net premium calculation. Similarly, they can also be applied to credit risk and reliability modelling. Numerical examples with four different distributions for modelling interarrival times are provided, and the implementation details are also discussed.

This paper is structured as follows. Section 2 introduces renewal shot-noise processes and the associated processes of discounted aggregate claims in insurance. In Section 3, based on the piecewise deterministic Markov process theory and the martingale methodology, we present the Feynmann-Kac formula. It is then used in Section 4 to derive the Laplace transforms of the moments of renewal shot-noise processes and discounted aggregate claims. Afterwards, in Section 5, we apply the results of the means and variances to the actuarial context for calculating net insurance premiums as well as credit risk and reliability modelling, for which we specify exponential, gamma, inverse Gaussian and folded normal distributions for modelling interarrival times, respectively. Section 6 contains concluding remarks.

2 Renewal Shot-noise Processes and Discounted Aggregate Claims

Claims arising from catastrophic events could be different from the different time interval of duration, and they could also depend on the time elapsed since the previous claim. Therefore, improved models beyond the Poisson process to predict claims arising from catastrophic events are required. For this purpose, let us start with a compound model of insurance risk with the additional economic assumption of a positive interest rate, and the accumulated value of aggregate claims up to time t

in continuous time on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ is

$$L_t = \sum_{i=1}^{N_t} X_i e^{r(t-T_i)}, \quad t \geq 0,$$

where

- $r > 0$ is the risk-free force of interest rate;
- $\{X_i\}_{i=1,2,\dots}$ are claim sizes (or jump sizes), which are assumed to be independent and identically distributed (*i.i.d.*) with cumulative distribution function (CDF) $H(x), x > 0$;
- $\{T_i\}_{i=1,2,\dots}$ are the claim occurrence times (or, *renewal epochs*), which follows a renewal point process $N_t = \sum_i \mathbf{1}_{\{T_i \leq t\}}$ with $N_0 = 0$.

\mathcal{F}_t is the associated natural filtration of L_t . Setting $L_t^0 = e^{-rt} L_t$, we have the discounted value at time 0 of aggregate claims (or, *discounted aggregate claims*) up to time t , i.e.,

$$L_t^0 = \sum_{i=1}^{N_t} X_i e^{-rT_i}. \quad (2.1)$$

As Jang (2004) and Jang and Krvavych (2004) noted the duality property between the process of discounted aggregate claims and the shot-noise process, we now introduce a *renewal shot-noise process* (or, shot-noise process driven by an ordinary renewal process)

$$S_t = S_0 e^{-\delta t} + \sum_{i=1}^{N_t} X_i e^{-\delta(t-T_i)}, \quad (2.2)$$

where δ is a constant. Setting $S_0 = 0$ and $\delta = -r$ in (2.2), the processes of S_t and L_t become identical. S_t was also discussed in Rice (1977) and was used as the stochastic intensity of a *double stochastic Poisson process* (or *Cox process*) in Møller and Torrisi (2005) and Dassios et al. (2015). Simulated sample paths of the renewal shot-noise process S_t and the underlying renewal process N_t are provided in Figure 1, where we assume the interarrival times follow an inverse Gaussian distribution and jump sizes follow an exponential distribution.

Note that, this process S_t is no longer within the usual framework of *affine processes* (Duffie et al., 2000, 2003) or a Markov process due to the additional renewal components. In order to establish a Markovian framework, we need to further include a supplementary stochastic process U_t , the *backward recurrence time* (Cox, 1962, p.27) (or *the time elapsed since the last jump arrived*) in the process S_t , i.e.,

$$U_t := U_0 + t - \sum_{i=1}^{N_t} \tau_i,$$




Figure 1: Simulated paths of renewal shot-noise process S_t and renewal process N_t when the interarrival times follow an inverse Gaussian distribution and jump sizes follow an exponential distribution

where $U_0 \geq 0$ is the initial value of U_t ; $\{\tau_i\}_{i=1,2,\dots}$ are interarrival times of claim arrivals, i.e.,

$$\tau_i := T_i - T_{i-1}, \quad i = 1, 2, \dots, \quad T_0 = 0,$$

and they are *i.i.d.* with the CDF $P(u)$, $u > 0$, which is assumed to be absolutely continuous with the associated density function $p(u)$. The idea of adding this supplementary variable U_t to make the underlying process Markovian can be found as early as in Cox (1955). U_t increases at unit rate till a jump arrives, then it goes back to 0. Note that, if $\rho(u)$ is denoted as the *failure rate* of the distribution, we have

$$P(u) = 1 - \exp\left(-\int_0^u \rho(v)dv\right), \quad p(u) = \rho(u) \exp\left(-\int_0^u \rho(v)dv\right),$$

where $\rho(u) = \frac{p(u)}{\bar{P}(u)}$, and $\bar{P}(u) := 1 - P(u)$ is denoted as the tail probability or the *survivor function* (Cox, 1962, p.3). For notation simplification, we denote the first mean and the Laplace transform respectively by

$$\gamma_1 := \int_0^\infty up(u)du < \infty, \quad \hat{p}(\theta) := \int_0^\infty e^{-\theta u}p(u)du < \infty.$$

We denote the m^{th} moment of S_t conditional on S_0 and U_0 and the associated Laplace transform with respect to time t respectively by

$$\begin{aligned} e_m(t; S_0, U_0; \delta) &:= \mathbb{E}[S_t^m \mid S_0, U_0], \quad m \in \mathbb{N}^+, \\ \hat{e}_m(\theta; S_0, U_0; \delta) &:= \int_0^\infty e^{-\theta t} \mathbb{E}[S_t^m \mid S_0, U_0] dt, \end{aligned}$$

and the moments of claim amounts by

$$\mu_k := \int_0^\infty x^k dH(x), \quad k = 0, 1, 2, \dots$$

The Laplace transform of any given function $f(t)$ in general is denoted by

$$\hat{f}(\theta) := \mathcal{L}_\theta\{f(t)\} := \int_0^\infty e^{-\theta t} f(t) dt.$$

All moments and Laplace transforms above are assumed to be finite.

3 Martingales

Let us define a process

$$Z_t = \int_0^t e^{-\theta u} \sum_{k=1}^m \kappa_k S_u^k du, \quad (3.1)$$

where $\theta \geq 0$ and $\{\kappa_k\}_{k=1,2,\dots,m}$ are all constants. The *infinitesimal generator* of (Z_t, S_t, U_t, t) acting on any function $g(z, s, u, t)$ belonging to its domain is given by

$$\begin{aligned} \mathcal{A}g(z, s, u, t) &= \left(e^{-\theta t} \sum_{k=1}^m \kappa_k s^k \right) \frac{\partial g}{\partial z} + \frac{\partial g}{\partial t} + \frac{\partial g}{\partial u} - \delta s \frac{\partial g}{\partial s} \\ &\quad + \frac{p(u)}{P(u)} \left[\int_0^\infty g(z, s+x, 0, t) dH(x) - g(z, s, u, t) \right], \quad (3.2) \end{aligned}$$

where $g : (0, \infty) \times (0, \infty) \times (0, \infty) \times \mathbb{R}^+ \rightarrow (0, \infty)$. It is sufficient that $g(z, s, u, t)$ is differentiable with respect to z, s, u, t for any z, s, u, t and that

$$\left| \int_0^\infty g(\cdot, s+x, \cdot, \cdot) dH(x) - g(\cdot, s, \cdot, \cdot) \right| < \infty \quad (3.3)$$

for $g(z, s, u, t)$ to belong to the domain of the (extended) generator \mathcal{A} . For the details on finding the generator of (Z_t, S_t, U_t, t) using the *piecewise deterministic Markov process theory* (Davis, 1984,

1993), see Dassios and Embrechts (1989), Dassios and Jang (2003), Rolski et al. (2008), Dassios and Zhao (2011, 2012, 2014) and many others.

Let us first provide a proposition as below which will be used very often in this paper.

Proposition 3.1. *The ordinary differential equation (ODE) of $A(u)$,*

$$a - \xi A(u) + A'(u) + \frac{p(u)}{\bar{P}(u)} [b + A(0) - A(u)] = 0, \quad (3.4)$$

has the solution

$$A(u) := \frac{a}{\xi} + \frac{b}{1 - \hat{p}(\xi)} \frac{e^{\xi u}}{\bar{P}(u)} \int_u^\infty e^{-\xi v} p(v) dv, \quad (3.5)$$

where a, b, ξ are all constants, and $\xi \geq 0$.

Now, in order to derive the m^{th} moment of S_t conditional on S_0 and U_0 at time $t = 0$ in the next section, we have to first find a suitable martingale with respect to the filtration \mathcal{F}_t , which is given in Theorem 3.1.

Theorem 3.1. *We have a \mathcal{F}_t -martingale*

$$Z_t + e^{-\theta t} \sum_{k=0}^m S_t^k A_k(U_t), \quad (3.6)$$

where

$$A_k(u) := \frac{\kappa_k}{\theta + \delta k} + \frac{\sum_{n=k+1}^m A_n(0) \binom{n}{k} \mu_{n-k}}{1 - \hat{p}(\theta + \delta k)} \frac{e^{(\theta + \delta k)u}}{\bar{P}(u)} \int_u^\infty e^{-(\theta + \delta k)v} p(v) dv, \quad k = 0, 1, \dots, m-1, \quad (3.7)$$

and

$$A_m(u) := \frac{\kappa_m}{\theta + \delta m}. \quad (3.8)$$

Proof. To find a \mathcal{F}_t -martingale, we assume a function in form of

$$g(z, s, u, t) = z + e^{-\theta t} \sum_{k=0}^m s^k A_k(u). \quad (3.9)$$

Setting $\mathcal{A}g = 0$ in (3.2), we obtain the equation

$$\begin{aligned} & \sum_{k=1}^m \kappa_k s^k - \theta \sum_{k=0}^m s^k A_k(u) + \sum_{k=0}^m s^k A'_k(u) - \delta \sum_{k=0}^m k s^k A_k(u) \\ & + \frac{p(u)}{\bar{P}(u)} \left[\int_0^\infty \sum_{k=0}^m (s+x)^k A_k(0) dH(x) - \sum_{k=0}^m s^k A_k(u) \right] = 0. \end{aligned}$$

Note that, based on $(s+x)^k = \sum_{j=0}^k \binom{k}{j} s^j x^{k-j}$ where

$$\binom{k}{j} := \frac{k!}{j!(k-j)!}, \quad j = 0, 1, \dots, k,$$

we have

$$\begin{aligned} 0 &= \sum_{k=1}^m \kappa_k s^k - \theta \sum_{k=0}^m s^k A_k(u) + \sum_{k=0}^m s^k A'_k(u) - \delta \sum_{k=0}^m k s^k A_k(u) \\ &\quad + \frac{p(u)}{\bar{P}(u)} \left[\int_0^\infty \sum_{k=0}^m \sum_{j=0}^k \binom{k}{j} s^j x^{k-j} A_k(0) dH(x) - \sum_{k=0}^m s^k A_k(u) \right] \\ &= \sum_{k=1}^m \kappa_k s^k - \theta \sum_{k=0}^m s^k A_k(u) + \sum_{k=0}^m s^k A'_k(u) - \delta \sum_{k=0}^m k s^k A_k(u) \\ &\quad + \frac{p(u)}{\bar{P}(u)} \left[\sum_{k=0}^m \sum_{j=0}^k \binom{k}{j} s^j \mu_{k-j} A_k(0) - \sum_{k=0}^m s^k A_k(u) \right], \end{aligned}$$

where

$$\mu_{k-j} = \int_0^\infty x^{k-j} dH(x), \quad j = 0, 1, 2, \dots, k.$$

Then, setting $\kappa_0 = 0$, we can rewrite it by

$$\begin{aligned} &\sum_{k=0}^m s^k \left[\kappa_k - (\theta + \delta k) A_k(u) + A'_k(u) \right] + \frac{p(u)}{\bar{P}(u)} \sum_{k=0}^m \left[A_k(0) \sum_{j=0}^k \binom{k}{j} \mu_{k-j} s^j - s^k A_k(u) \right] \\ &= \sum_{k=0}^m c_k(u) s^k = 0, \end{aligned} \tag{3.10}$$

where $c_k(u)$ is the coefficient of s^k , i.e.,

$$c_k(u) := \kappa_k - (\theta + \delta k) A_k(u) + A'_k(u) + \frac{p(u)}{\bar{P}(u)} \left[\sum_{n=k}^m A_n(0) \binom{n}{k} \mu_{n-k} - A_k(u) \right], \quad k = 0, 1, \dots, m.$$

Since (3.10) should hold for any s^k where $\forall k \in \{0, 1, 2, \dots, m\}$, each coefficient should be equal to zero, i.e., we have the ODEs

$$c_k(u) = 0, \quad k = 0, 1, \dots, m.$$

Using Proposition 3.1, we have the solutions

$$A_k(u) = \frac{\kappa_k}{\theta + \delta k} + \frac{\sum_{n=k}^m A_n(0) \binom{n}{k} \mu_{n-k}}{1 - \hat{p}(\theta + \delta k)} \frac{e^{(\theta + \delta k)u}}{\bar{P}(u)} \int_u^\infty e^{-(\theta + \delta k)v} p(v) dv,$$

with the boundary conditions $A_k(0) = 0$ for $k = 0, 1, \dots, m$. More specifically, they are equivalent to (3.7) for $k = 0, 1, 2, \dots, m - 1$ and (3.8) for $k = m$. Finally, it is easy to see that, this function (3.9) is differentiable with respect to all its arguments z, s, u, t , and also the expectation of the associated jumps is bounded, i.e., it satisfies (3.3). Hence, it belongs to the domain of the (extended) generator \mathcal{A} . It is based on the *piecewise deterministic Markov process theory*, which was developed by Davis (1984, Theorem 5.5, p.367), see also more details on this theory and its conditions in the book by Davis (1993, p.69). \square

Proposition 3.2. *We have the Feynmann-Kac formula*

$$\mathbb{E} \left[\int_0^\infty e^{-\theta t} \sum_{k=1}^m \kappa_k S_t^k dt \mid S_0, U_0 \right] = \sum_{k=0}^m S_0^k A_k(U_0). \quad (3.11)$$

Proof. Using the \mathcal{F}_t -martingale (3.6) provided in Theorem 3.1 and the martingale property, we have the expectation conditional on S_0 and U_0 at time $t = 0$ by

$$\mathbb{E} \left[Z_t + e^{-\theta t} \sum_{k=0}^m S_t^k A_k(U_t) \mid S_0, U_0 \right] = \sum_{k=0}^m S_0^k A_k(U_0). \quad (3.12)$$

Setting $t = \infty$ in (3.12), (3.11) follows immediately. \square

Applications of the Feynmann-Kac formula in general can be found in Karatzas and Shreve (1991). Its applications to financial mathematics can be noticed in Linetsky (1997, 2004, 2007) and the refereed papers therein. More recently, Goovaerts et al. (2012) constructed a recursive scheme for the Laplace transform of the transition density function of a diffusion process using the Feynmann-Kac formula, also see Shang et al. (2011).

4 Moments

In this section, we first derive Laplace transforms of the conditional moments and asymptotic moments of renewal shot-noise processes and discounted aggregate claims, respectively. Then, by inverting the Laplace transforms, we obtain the asymptotic moments and the first conditional moments in explicit forms. They are the main contribution of this paper. As examples, the associated first two moments and variances are discussed in more details.

4.1 Moments of Renewal Shot-noise Processes

Theorem 4.1. *The Laplace transform (with respect to time t) of the m^{th} moment of S_t conditional on S_0 and U_0 is given by*

$$\hat{e}_m(\theta; S_0, U_0; \delta) = \sum_{k=0}^m S_0^k A_k^*(U_0), \quad (4.1)$$

where the series of functions $\{A_k^*(u)\}_{k=0,1,\dots,m}$ can be iteratively solved from the system of equations

$$A_m^*(u) := \frac{1}{\theta + m\delta}, \quad (4.2)$$

$$A_k^*(u) := \frac{\sum_{n=k+1}^m A_n^*(0) \binom{n}{k} \mu_{n-k}}{1 - \hat{p}(\theta + \delta k)} \frac{e^{(\theta+\delta k)u}}{\bar{P}(u)} \int_u^\infty e^{-(\theta+\delta k)v} p(v) dv, \quad k = m-1, m-2, \dots, 1, 0, \quad (4.3)$$

with

$$\begin{aligned} A_m^*(0) &= \frac{1}{\theta + m\delta}, \\ A_k^*(0) &= \frac{\hat{p}(\theta + \delta k)}{1 - \hat{p}(\theta + \delta k)} \sum_{n=k+1}^m A_n^*(0) \binom{n}{k} \mu_{n-k}, \quad k = m-1, m-2, \dots, 1, 0. \end{aligned}$$

Proof. Firstly, we express (3.11) in terms of Laplace transforms by

$$\sum_{k=1}^m \kappa_k \hat{e}_k(\theta; S_0, U_0; \delta) = \sum_{k=0}^m S_0^k A_k(U_0). \quad (4.4)$$

Setting $\kappa_m = 1$ and $\kappa_k = 0$ for all $k = 1, \dots, m-1$ in (4.4), (3.7) and (3.8), we have (4.1), (4.2) and (4.3), respectively. Further setting $u = 0$ in (4.3), we have

$$A_k^*(0) = \frac{\sum_{n=k+1}^m A_n^*(0) \binom{n}{k} \mu_{n-k}}{1 - \hat{p}(\theta + \delta k)} \int_0^\infty e^{-(\theta+\delta k)v} p(v) dv = \frac{\hat{p}(\theta + \delta k)}{1 - \hat{p}(\theta + \delta k)} \sum_{n=k+1}^m A_n^*(0) \binom{n}{k} \mu_{n-k}.$$

□

Based on Theorem 4.1, it is straightforward to obtain the Laplace transform for any conditional moment of S_t . For example, the Laplace transforms of the first two moments are specified as below.

Corollary 4.1. *The Laplace transform of the first moment of S_t conditional on S_0 and U_0 is given by*

$$\hat{e}_1(\theta; S_0, U_0; \delta) = \frac{S_0}{\theta + \delta} + \frac{\mu_1}{1 - \hat{p}(\theta)} \frac{1}{\theta + \delta} \frac{e^{\theta U_0}}{\bar{P}(U_0)} \int_{U_0}^\infty e^{-\theta v} p(v) dv. \quad (4.5)$$

Proof. Set $\kappa_2 = 0$ and $\kappa_1 = 1$ in (3.7) and (3.8), then, we have

$$A_2^*(u) = 0, \quad A_1^*(u) = \frac{1}{\theta + \delta}, \quad A_0^*(u) = \frac{\mu_1}{1 - \hat{p}(\theta)} \frac{1}{\theta + \delta} \frac{e^{\theta u}}{\bar{P}(u)} \int_u^\infty e^{-\theta v} p(v) dv.$$

From (4.4), the result follows. \square

Corollary 4.2. *The Laplace transform of the second moment of S_t conditional on S_0 and U_0 is given by*

$$\begin{aligned} \hat{e}_2(\theta; S_0, U_0; \delta) &= \frac{S_0^2}{\theta + 2\delta} + \frac{S_0}{\theta + 2\delta} \frac{2\mu_1}{1 - \hat{p}(\theta + \delta)} \frac{e^{(\theta + \delta)U_0}}{\bar{P}(U_0)} \int_{U_0}^\infty e^{-(\theta + \delta)v} p(v) dv \\ &\quad + \frac{\frac{\mu_2}{\theta + 2\delta} + \frac{1}{\theta + 2\delta} \frac{2\mu_1^2 \hat{p}(\theta + \delta)}{1 - \hat{p}(\theta + \delta)}}{1 - \hat{p}(\theta)} \frac{e^{\theta U_0}}{\bar{P}(U_0)} \int_{U_0}^\infty e^{-\theta v} p(v) dv. \end{aligned} \quad (4.6)$$

Proof. Setting $\kappa_2 = 1$ and $\kappa_1 = 0$ in (3.7) and (3.8), we have

$$\begin{aligned} A_2^*(u) &= \frac{1}{\theta + 2\delta}, \\ A_1^*(u) &= \frac{1}{\theta + 2\delta} \frac{2\mu_1}{1 - \hat{p}(\theta + \delta)} \frac{e^{(\theta + \delta)u}}{\bar{P}(u)} \int_u^\infty e^{-(\theta + \delta)v} p(v) dv, \\ A_0^*(u) &= \frac{\frac{\mu_2}{\theta + 2\delta} + \frac{1}{\theta + 2\delta} \frac{2\mu_1^2 \hat{p}(\theta + \delta)}{1 - \hat{p}(\theta + \delta)}}{1 - \hat{p}(\theta)} \frac{e^{\theta u}}{\bar{P}(u)} \int_u^\infty e^{-\theta v} p(v) dv. \end{aligned}$$

From (4.4), the result follows. \square

The distribution converges pretty fast, and we can easily observe how the distribution converges to its asymptotic distribution via its mean.

Corollary 4.3. *We have the asymptotics of the first moment,*

$$e_1(t; S_0, U_0; \delta) = d_0 + d_1 e^{-\delta t} + o(e^{-\delta t}), \quad (4.7)$$

where

$$d_0 := \frac{\mu_1}{\delta \gamma_1}, \quad d_1 := S_0 + \frac{\mu_1}{\bar{P}(U_0)} \frac{e^{-\delta U_0}}{1 - \hat{p}(-\delta)} \int_{U_0}^\infty e^{\delta v} p(v) dv.$$

Proof. The Laplace transform of the first moment of S_t conditional on S_0 and U_0 is given by (4.5).

We know that the limit $\lim_{t \rightarrow \infty} e_1(t; S_0, U_0; \delta)$ exists, more precisely,

$$\begin{aligned} d_0 &:= \lim_{t \rightarrow \infty} e_1(t; S_0, U_0; \delta) \\ &= \lim_{\theta \rightarrow 0} \theta \hat{e}_1(\theta; S_0, U_0; \delta) \end{aligned}$$

$$\begin{aligned}
&= \frac{\mu_1}{\bar{P}(U_0)} \int_{U_0}^{\infty} \lim_{\theta \rightarrow 0} \left[\frac{1}{\theta + \delta} \frac{\theta}{1 - \hat{p}(\theta)} e^{\theta(U_0 - v)} \right] p(v) dv \\
&= \frac{1}{\delta} \frac{1}{\gamma_1} \frac{\mu_1}{\bar{P}(U_0)} \int_{U_0}^{\infty} p(v) dv \\
&= \frac{\mu_1}{\delta \gamma_1}.
\end{aligned}$$

Define the function

$$g(t) := e^{\delta t} \left[e_1(t; S_0, U_0; \delta) - d_0 \right],$$

and its Laplace transform

$$\hat{g}(\theta) := \int_0^{\infty} e^{-\theta t} g(t) dt.$$

Then, we have

$$\begin{aligned}
d_1 &:= \lim_{t \rightarrow \infty} g(t) \\
&= \lim_{\theta \rightarrow 0} \theta \hat{g}(\theta) \\
&= \lim_{\theta \rightarrow 0} \theta \left[\hat{e}_1(\theta - \delta; S_0, U_0; \delta) - \frac{d_0}{\theta - \delta} \right] \\
&= \lim_{\theta \rightarrow 0} \theta \left[\frac{1}{\theta} S_0 + \frac{1}{\theta} \frac{\mu_1}{1 - \hat{p}(\theta - \delta)} \frac{e^{(\theta - \delta)U_0}}{\bar{P}(U_0)} \int_{U_0}^{\infty} e^{-(\theta - \delta)v} p(v) dv \right] \\
&= S_0 + \frac{\mu_1}{\bar{P}(U_0)} \int_{U_0}^{\infty} \lim_{\theta \rightarrow 0} \left[\frac{e^{(\theta - \delta)(U_0 - v)}}{1 - \hat{p}(\theta - \delta)} \right] p(v) dv \\
&= S_0 + \frac{\mu_1}{\bar{P}(U_0)} \frac{e^{-\delta U_0}}{1 - \hat{p}(-\delta)} \int_{U_0}^{\infty} e^{\delta v} p(v) dv.
\end{aligned}$$

Therefore, we have (4.7). □

We can see from (4.7) that the conditional moment converges at an exponential rate with respect to time t , and the asymptotic results could provide reasonable approximations to their moments and distributions.

The initial value U_0 is usually unknown in practice. To calculate the first two conditional moments of S_t for actuarial applications, we assign the asymptotic (or limiting) distribution of U_t to U_0 for mathematical convenience, which can provide reasonable approximations and also substantially simplify the expressions of the Laplace transforms of moments. To do so, we first state a proposition in Cox and Miller (1965, p.347) or Cox (1962, p.61), which is a well known result in renewal theory.

Proposition 4.1. *The asymptotic (or limiting) distribution of U_t , denoted by Π , has the density function*

$$f_{\Pi}(u) := \frac{\bar{P}(u)}{\gamma_1} = \frac{1}{\gamma_1} \exp\left(-\int_0^u \rho(v)dv\right), \quad u \geq 0.$$

Π is in fact the limiting distribution of U_t when $t \rightarrow \infty$, and it can serve a reasonable approximation for the distribution of U_t when the underlying process has been running for a relatively long period and is close to the stationary (asymptotic) state (Cox, 1962, Chapter 5, p.61-70).

Now, let us start with finding the asymptotic m^{th} moment of S_t when $U_0 \sim \Pi$, denoted by

$$e_m(t; S_0; \delta) := \mathbb{E}[S_t^m | S_0].$$

Denote the Laplace transform (with respect to time t) of the m^{th} moment of S_t conditional on S_0 by

$$\hat{e}_m(\theta; S_0; \delta) := \mathbb{E}[\hat{e}_m(\theta; S_0, U_0)], \quad U_0 \sim \Pi.$$

Proposition 4.2. *If $U_0 \sim \Pi$, then, we have*

$$\mathbb{E}\left[\frac{e^{\xi U_0}}{\bar{P}(U_0)} \int_{U_0}^{\infty} e^{-\xi v} p(v)dv\right] = \frac{1}{\gamma_1} \frac{1 - \hat{p}(\xi)}{\xi}.$$

Proof.

$$\begin{aligned} \mathbb{E}\left[\frac{e^{\xi U_0}}{\bar{P}(U_0)} \int_{U_0}^{\infty} e^{-\xi v} p(v)dv\right] &= \int_{u=0}^{\infty} \frac{e^{\xi u}}{\bar{P}(u)} \int_{v=u}^{\infty} e^{-\xi v} p(v)dv f_{\Pi}(u)du \\ &= \int_{u=0}^{\infty} \int_{v=u}^{\infty} e^{-\xi v} p(v) \frac{e^{\xi u}}{\gamma_1} dv du \\ &= \int_{v=0}^{\infty} \int_{u=0}^v e^{-\xi v} p(v) \frac{e^{\xi u}}{\gamma_1} du dv \\ &= \frac{1}{\gamma_1} \int_{v=0}^{\infty} e^{-\xi v} p(v) \left(\int_{u=0}^v e^{\xi u} du\right) dv \\ &= \frac{1}{\gamma_1} \int_{v=0}^{\infty} e^{-\xi v} p(v) \frac{e^{\xi v} - 1}{\xi} dv \\ &= \frac{1}{\gamma_1} \int_{v=0}^{\infty} p(v) \frac{1 - e^{-\xi v}}{\xi} dv \\ &= \frac{1}{\gamma_1} \frac{1 - \hat{p}(\xi)}{\xi}. \end{aligned}$$

□

Theorem 4.2. For $U_0 \sim \Pi$, the Laplace transform of the m^{th} moment of S_t conditional on S_0 is given by

$$\hat{e}_m(\theta; S_0; \delta) = \sum_{k=0}^m B_k(\theta) S_0^k, \quad (4.8)$$

where

$$\begin{aligned} B_m(\theta) &:= \frac{1}{\theta + m\delta}, \\ B_k(\theta) &:= \frac{1}{\gamma_1} \frac{1}{\theta + \delta k} \sum_{n=k+1}^m A_n^*(0) \binom{n}{k} \mu_{n-k}, \quad k = m-1, m-2, \dots, 1, 0. \end{aligned} \quad (4.9)$$

Proof. Using (4.1) and Proposition 4.1, we have

$$\hat{e}_m(\theta; S_0; \delta) = \mathbb{E}[\hat{e}_m(\theta; S_0, U_0; \delta)] = \mathbb{E} \left[\sum_{k=0}^m S_0^k A_k^*(U_0) \right] = \sum_{k=0}^m S_0^k \mathbb{E}[A_k^*(U_0)],$$

where

$$\begin{aligned} B_k(\theta) := \mathbb{E}[A_k^*(U_0)] &= \frac{\sum_{n=k+1}^m A_n^*(0) \binom{n}{k} \mu_{n-k}}{1 - \hat{p}(\theta + \delta k)} \mathbb{E} \left[\frac{e^{(\theta + \delta k)U_0}}{\bar{P}(U_0)} \int_{U_0}^{\infty} e^{-(\theta + \delta k)v} p(v) dv \right] \\ &= \frac{\sum_{n=k+1}^m A_n^*(0) \binom{n}{k} \mu_{n-k}}{1 - \hat{p}(\theta + \delta k)} \frac{1}{\gamma_1} \frac{1 - \hat{p}(\theta + \delta k)}{\theta + \delta k} \\ &= \frac{1}{\gamma_1} \frac{1}{\theta + \delta k} \sum_{n=k+1}^m A_n^*(0) \binom{n}{k} \mu_{n-k}, \quad k = m-1, m-2, \dots, 1, 0. \end{aligned}$$

□

Theorem 4.3. For $U_0 \sim \Pi$, the m^{th} asymptotic moment of S_t is given by

$$\lim_{t \rightarrow \infty} \mathbb{E}[S_t^m | S_0] = \frac{1}{\gamma_1} \sum_{k=1}^m \mu_k A_k^{**}, \quad (4.10)$$

where the constants $\{A_k^{**}\}_{k=1, \dots, m}$ can be calculated iteratively from

$$\begin{aligned} A_m^{**} &:= \frac{1}{m\delta}, \\ A_k^{**} &:= \frac{\hat{p}(\delta k)}{1 - \hat{p}(\delta k)} \sum_{n=k+1}^m A_n^{**} \binom{n}{k} \mu_{n-k}, \quad k = m-1, m-2, \dots, 1. \end{aligned}$$

Proof. By the *final value theorem*, we have

$$\lim_{t \rightarrow \infty} \mathbb{E}[S_t^m | S_0] = \lim_{\theta \rightarrow 0} \theta \hat{e}_m(\theta; S_0) = \sum_{k=0}^m S_0^k \lim_{\theta \rightarrow 0} \theta B_k(\theta) = \lim_{\theta \rightarrow 0} \theta B_0(\theta),$$

since

$$\begin{aligned} \lim_{\theta \rightarrow 0} \theta B_m(\theta) &= 0, \\ \lim_{\theta \rightarrow 0} \theta B_k(\theta) &= \frac{1}{\gamma_1} \sum_{n=k+1}^m \binom{n}{k} \mu_{n-k} \lim_{\theta \rightarrow 0} \frac{\theta}{\theta + \delta k} A_n^*(0) = 0, \quad k = m-1, m-2, \dots, 1. \end{aligned}$$

Note that, according to (4.9), we have

$$B_0(\theta) = \frac{1}{\gamma_1} \frac{1}{\theta} \sum_{n=1}^m A_n^*(0) \mu_n,$$

then,

$$\lim_{\theta \rightarrow 0} \theta B_0(\theta) = \frac{1}{\gamma_1} \sum_{n=1}^m \mu_n \lim_{\theta \rightarrow 0} A_n^*(0) = \frac{1}{\gamma_1} \sum_{k=1}^m \mu_k \lim_{\theta \rightarrow 0} A_k^*(0) = \frac{1}{\gamma_1} \sum_{k=1}^m \mu_k A_k^{**},$$

where $A_k^{**} = \lim_{\theta \rightarrow 0} A_k^*(0)$. □

Now, let us start with finding the first moment of S_t conditional on S_0 at time $t = 0$ by inverting its Laplace transform.

Corollary 4.4. *For $U_0 \sim \Pi$, the first moment of S_t conditional on S_0 is given by*

$$e_1(t; S_0; \delta) = S_0 e^{-\delta t} + \frac{\mu_1}{\gamma_1} \left(\frac{1 - e^{-\delta t}}{\delta} \right), \quad (4.11)$$

and the first asymptotic moment is given by

$$\lim_{t \rightarrow \infty} \mathbb{E}[S_t | S_0] = \frac{\mu_1}{\gamma_1 \delta}. \quad (4.12)$$

Proof. Using (4.6) and Proposition 4.1, and setting $\xi = \theta$ in Proposition 4.2, the Laplace transform (with respect to time t) of the first moment of S_t conditional on S_0 for $U_0 \sim \Pi$ is given by

$$\begin{aligned} \hat{e}_1(\theta; S_0; \delta) &= \frac{S_0}{\theta + \delta} + \frac{\mu_1}{1 - \hat{p}(\theta)} \frac{1}{\theta + \delta} \mathbb{E} \left[\frac{e^{\theta U_0}}{\bar{P}(U_0)} \int_{U_0}^{\infty} e^{-\theta v} p(v) dv \right] \\ &= \frac{S_0}{\theta + \delta} + \frac{\mu_1}{\gamma_1} \frac{1}{\theta(\theta + \delta)} \\ &= S_0 \frac{1}{\theta + \delta} + \frac{\mu_1}{\gamma_1} \frac{1}{\delta} \left(\frac{1}{\theta} - \frac{1}{\theta + \delta} \right). \end{aligned}$$

Inverting it immediately gives us (4.11). Note that, we have Laplace transforms

$$\mathcal{L}_\theta \{e^{-\delta t}\} = \frac{1}{\theta + \delta}, \quad \mathcal{L}_\theta \{1\} = \frac{1}{\theta}.$$

Its asymptotic result in (4.12) follows by setting $t \rightarrow \infty$ in (4.11). \square

Unfortunately, in general, it is not possible for us to obtain other conditional moments explicitly beyond the first moments. Therefore, we have to develop numerical methods for estimation.

Corollary 4.5. *For $U_0 \sim \Pi$, the second moment of S_t conditional on S_0 is given by*

$$e_2(t; S_0; \delta) = S_0^2 e^{-2\delta t} + \frac{2\mu_1}{\delta\gamma_1} S_0 (e^{-\delta t} - e^{-2\delta t}) + \frac{\mu_2}{\gamma_1} \left(\frac{1 - e^{-2\delta t}}{2\delta} \right) + \frac{\mu_1^2}{\delta\gamma_1} \frac{\hat{p}(\delta)}{1 - \hat{p}(\delta)} F_4(t), \quad (4.13)$$

where $F_4(t)$ is a function of time t with the Laplace transform

$$\hat{F}_4(\theta) = \frac{1 - \hat{p}(\delta)}{\hat{p}(\delta)} \frac{2\delta}{\theta(\theta + 2\delta)} \frac{\hat{p}(\theta + \delta)}{1 - \hat{p}(\theta + \delta)}; \quad (4.14)$$

The asymptotic second moment is given by

$$\frac{1}{\gamma_1 \delta} \left[\frac{\mu_2}{2} + \mu_1^2 \frac{\hat{p}(\delta)}{1 - \hat{p}(\delta)} \right]. \quad (4.15)$$

Proof. Using (4.5) and Proposition 4.1 and 4.2, the Laplace transform (with respect to time t) of the second moment of S_t conditional on S_0 for $U_0 \sim \Pi$ is given by

$$\begin{aligned} & \hat{e}_2(\theta; S_0; \delta) \\ &= \frac{S_0^2}{\theta + 2\delta} + \frac{2\mu_1 S_0}{(\theta + 2\delta)(\theta + \delta)} \frac{1}{\gamma_1} + \left[\frac{\mu_2}{\theta + 2\delta} + \frac{\hat{p}(\theta + \delta)}{1 - \hat{p}(\theta + \delta)} \frac{2\mu_1^2}{\theta + 2\delta} \right] \frac{1}{\theta} \frac{1}{\gamma_1} \\ &= S_0^2 \frac{1}{\theta + 2\delta} + \frac{2\mu_1 S_0}{\gamma_1} \frac{1}{\delta} \left(\frac{1}{\theta + \delta} - \frac{1}{\theta + 2\delta} \right) + \frac{\mu_2}{\gamma_1} \frac{1}{2\delta} \left(\frac{1}{\theta} - \frac{1}{\theta + 2\delta} \right) \\ & \quad + \frac{2\mu_1^2}{\gamma_1} \frac{1}{\theta} \frac{\hat{p}(\theta + \delta)}{1 - \hat{p}(\theta + \delta)} \frac{1}{\theta + 2\delta} \end{aligned} \quad (4.16)$$

$$\begin{aligned} &= S_0^2 \frac{1}{\theta + 2\delta} + \frac{2\mu_1 S_0}{\gamma_1} \frac{1}{\delta} \left(\frac{1}{\theta + \delta} - \frac{1}{\theta + 2\delta} \right) + \frac{\mu_2}{\gamma_1} \frac{1}{2\delta} \left(\frac{1}{\theta} - \frac{1}{\theta + 2\delta} \right) \\ & \quad + \frac{\mu_1^2}{\delta\gamma_1} \frac{\hat{p}(\delta)}{1 - \hat{p}(\delta)} \hat{F}_4(\theta). \end{aligned} \quad (4.17)$$

The first three terms of (4.17) can be inverted analytically, then, we obtain (4.13). Based on Theorem 4.10, setting $m = 2$, we can calculate

$$A_2^{**} = \frac{1}{2\delta}, \quad A_1^{**} = \frac{\hat{p}(\delta)}{1 - \hat{p}(\delta)} \frac{\mu_1}{\delta}.$$

Substituting A_1^{**} and A_2^{**} into (4.10), we derive (4.15). \square

Corollary 6.3 in Léveillé and Garrido (2001a, p.230) confirms both our results (4.12) and (4.15). Interestingly, from a probabilistic point of view, function $F_4(t)$ in (4.14) can be nicely interpreted as the CDF of a random time τ^* , which can be estimated by the following algorithm:

Algorithm 4.1 (Decomposition Approach). *The random time τ^* can be exactly sampled by the following distributional decomposition:*

$$\tau^* \stackrel{\mathcal{D}}{=} E^* + \sum_{i=1}^I E_i, \quad (4.18)$$

where

- E^* is an exponential random variable of constant rate 2δ , i.e., $E^* \sim \text{Exp}(2\delta)$;
- I is a geometric random variable with success probability parameter $1 - \hat{p}(\delta)$, i.e., $I \sim \text{Geometric}(1 - \hat{p}(\delta))$ with the probability mass distribution

$$\Pr \{I = i\} = \hat{p}^{i-1}(\delta) [1 - \hat{p}(\delta)], \quad i = 1, 2, 3, \dots;$$

- $\{E_i\}_{i=1,2,\dots}$ are i.i.d. random variables with the identical Laplace transform

$$\hat{f}_{E_i}(\theta) := \mathbb{E} [e^{-\theta E_i}] = \frac{\hat{p}(\theta + \delta)}{\hat{p}(\delta)}. \quad (4.19)$$

Proof. The Laplace transform of $F_4(t)$ specified by (4.14) can be rewritten as

$$\begin{aligned} \hat{F}_4(\theta) &= \frac{1}{\theta} \times 2\delta \frac{1 - \hat{p}(\delta)}{\hat{p}(\delta)} \frac{\hat{p}(\theta + \delta)}{1 - \hat{p}(\theta + \delta)} \frac{1}{\theta + 2\delta} \\ &= \frac{1}{\theta} \times 2\delta \frac{1 - \hat{p}(\delta)}{\hat{p}(\delta)} \frac{1}{\theta + 2\delta} \left[\frac{1}{1 - \hat{p}(\theta + \delta)} - 1 \right] \\ &= \frac{1}{\theta} \times 2\delta \frac{1 - \hat{p}(\delta)}{\hat{p}(\delta)} \frac{1}{\theta + 2\delta} \sum_{i=1}^{\infty} \hat{p}^i(\theta + \delta) \\ &= \frac{1}{\theta} \times \sum_{i=1}^{\infty} \hat{p}^{i-1}(\delta) [1 - \hat{p}(\delta)] \left[\frac{\hat{p}(\theta + \delta)}{\hat{p}(\delta)} \right]^i \frac{2\delta}{\theta + 2\delta} \\ &= \frac{1}{\theta} \times \sum_{i=1}^{\infty} \Pr \{I = i\} \left[\frac{\hat{p}(\theta + \delta)}{\hat{p}(\delta)} \right]^i \frac{2\delta}{\theta + 2\delta} \\ &= \frac{1}{\theta} \times \mathbb{E} \left[\left[\frac{\hat{p}(\theta + \delta)}{\hat{p}(\delta)} \right]^I \right] \frac{2\delta}{\theta + 2\delta} \\ &= \frac{1}{\theta} \times \int_0^{\infty} e^{-\theta t} f_4(t) dt, \end{aligned}$$

where $f_4(t)$ can be considered as the density function of the random time τ^* defined by (4.18), and

the associated CDF is given by

$$F_4(t) := \Pr \{ \tau^* \leq t \} = \mathbb{E} [\mathbf{1} \{ \tau^* \leq t \}].$$

The Laplace transform of the CDF is given by

$$\hat{F}_4(\theta) := \int_0^{\infty} e^{-\theta t} F_4(t) dt = \frac{1}{\theta} \hat{f}_4(\theta),$$

where

$$\begin{aligned} \hat{f}_4(\theta) &:= \int_0^{\infty} e^{-\theta t} f_4(t) dt \\ &= \frac{1 - \hat{p}(\delta)}{\hat{p}(\delta)} \frac{\hat{p}(\theta + \delta)}{1 - \hat{p}(\theta + \delta)} \frac{2\delta}{\theta + 2\delta} \\ &= \mathbb{E} \left[\left[\frac{\hat{p}(\theta + \delta)}{\hat{p}(\delta)} \right]^I \right] \times \frac{2\delta}{\theta + 2\delta} \\ &= \mathbb{E} \left[\mathbb{E} \left[e^{-\theta \sum_{i=1}^I E_i} \right] \right] \times \mathbb{E} \left[e^{-\theta E^*} \right] \\ &= \mathbb{E} \left[e^{-\theta (E^* + \sum_{i=1}^I E_i)} \right] \\ &= \mathbb{E} \left[e^{-\theta \tau^*} \right]. \end{aligned} \tag{4.20}$$

So, we have the decomposition (4.18). Note that, E_1, E_2, \dots, E_I have the identical Laplace transform (4.19), and they are well defined random variables, since

$$\hat{f}_{E_i}(\theta) = \frac{\hat{p}(\theta + \delta)}{\hat{p}(\delta)} = \int_0^{\infty} e^{-\theta t} \frac{e^{-\delta t}}{\hat{p}(\delta)} p(t) dt = \int_0^{\infty} e^{-\theta t} f_{E_i}(t) dt,$$

and we have the density function of E_i via the *Esscher transform* (Gerber and Shiu, 1994) (or, *exponential tilting*) as

$$f_{E_i}(t) = \frac{e^{-\delta t}}{\hat{p}(\delta)} p(t), \tag{4.21}$$

and

$$\int_0^{\infty} f_{E_i}(t) dt = \int_0^{\infty} \frac{e^{-\delta t}}{\hat{p}(\delta)} p(t) dt = \frac{\hat{p}(\delta)}{\hat{p}(\delta)} = 1.$$

□

Note that, since $F_4(t)$ can be interpreted as a CDF, we have

$$\lim_{t \rightarrow \infty} F_4(t) = 1.$$

Setting $t \rightarrow \infty$ in (4.13), again, we obtain the asymptotic result (4.15). Alternatively, $F_4(t)$ as a CDF could be estimated by the numerical inversion of Laplace transform (Abate and Whitt, 1992, 1995, 2006), which will be discussed in detail in Section 5.

Corollary 4.6. *For $U_0 \sim \Pi$, the variance of S_t conditional on S_0 is given by*

$$\text{Var} [S_t | S_0] = \frac{\mu_2}{\gamma_1} \left(\frac{1 - e^{-2\delta t}}{2\delta} \right) - \frac{\mu_1^2}{\gamma_1^2} \left(\frac{1 - e^{-\delta t}}{\delta} \right)^2 + \frac{\mu_1^2}{\delta \gamma_1} \frac{\hat{p}(\delta)}{1 - \hat{p}(\delta)} F_4(t). \quad (4.22)$$

Proof. Based on the first moment (4.11) and the second moment (4.13), we have the variance

$$\text{Var} [S_t | S_0] = \mathbb{E} [S_t^2 | S_0] - (\mathbb{E} [S_t | S_0])^2.$$

□

The moments, of course, can be estimated by the *direct simulation* for sample paths of S_t : Say, to estimate the moments of S_t at time $T > 0$, we have to simulate all interarrival times, jump sizes within the time period $[0, T]$; and moreover, as intermediate steps required, we also need to solve all ODEs recursively between two successive jumps, in order exactly simulate the distribution of S_t at an arbitrary time point T . In fact, it is a *path-dependent approach*. However, our *decomposition approach* provides a shortcut, which avoids simulating full paths of the underlying stochastic processes but only needs a few simple random variables as illustrated in Algorithm 4.1. Essentially, we use a Monte Carlo alternative Laplace transform inversion.

4.2 Moments of Discounted Aggregate Claims

Denote the m^{th} moment of L_t^0 by

$$\ell_m(t) := \mathbb{E} [(L_t^0)^m],$$

and the associated Laplace transform by

$$\hat{\ell}_m(\theta) := \int_0^{\infty} e^{-\theta t} \ell_m(t) dt,$$

which can be obtained explicitly as below.

Theorem 4.4. *For $U_0 \sim \Pi$, the Laplace transform of the m^{th} moment of L_t^0 is given by*

$$\hat{\ell}_m(\theta) = \frac{1}{\gamma_1} \frac{1}{\theta + mr} \sum_{n=1}^m \mu_n \tilde{A}_n, \quad (4.23)$$

where

$$\begin{aligned}\tilde{A}_m &= \frac{1}{\theta}, \\ \tilde{A}_k &= \frac{\hat{p}(\theta + (m-k)r)}{1 - \hat{p}(\theta + (m-k)r)} \sum_{n=k+1}^m \tilde{A}_n \binom{n}{k} \mu_{n-k}, \quad k = m-1, m-2, \dots, 1, 0.\end{aligned}$$

Proof. Note that, by setting

$$S_0 = 0, \quad L_t^0 = e^{-rt} S_t, \quad \delta = -r,$$

in the process S_t , we recover the associated L_t^0 . Using this duality property, in general, we have the m^{th} moment of L_t^0 by

$$\mathbb{E} \left[(L_t^0)^m \right] = e^{-mrt} e_m(t; 0; -r).$$

Then, we have its Laplace transform

$$\hat{\ell}_m(\theta) = \mathcal{L}_\theta \left\{ e^{-mrt} e_m(t; 0; -r) \right\} = \hat{e}_m(\theta + mr; 0; -r),$$

where $\hat{e}_m(\theta; S_0; \delta)$ is specified by (4.8). □

Based on the Laplace transform (4.23), as examples, we can compute the first two moments and the variance as below.

Corollary 4.7. *The first moment and the variance of L_t^0 are given by*

$$\ell_1(t) = \frac{\mu_1}{\gamma_1} \left(\frac{1 - e^{-rt}}{r} \right), \quad (4.24)$$

$$\ell_2(t) = \frac{\mu_2}{\gamma_1} \left(\frac{1 - e^{-2rt}}{2r} \right) + \frac{\mu_1^2}{r\gamma_1} \frac{\hat{p}(r)}{1 - \hat{p}(r)} \tilde{F}_4(t), \quad (4.25)$$

$$\text{Var} [L_t^0] = \frac{\mu_2}{\gamma_1} \left(\frac{1 - e^{-2rt}}{2r} \right) - \frac{\mu_1^2}{\gamma_1^2} \left(\frac{1 - e^{-rt}}{r} \right)^2 + \frac{\mu_1^2}{r\gamma_1} \frac{\hat{p}(r)}{1 - \hat{p}(r)} \tilde{F}_4(t), \quad (4.26)$$

where $\tilde{F}_4(t)$ is the CDF of random time $\tilde{\tau}^*$ which can be exactly simulated the same as τ^* via Algorithm 4.1 by replacing δ by r .

Proof. Setting

$$S_0 = 0, \quad L_t^0 = e^{-rt} S_t, \quad \delta = -r,$$

in (4.11), we have the mean (4.24), i.e.,

$$\ell_1(t) = \mathbb{E} [L_t^0] = e^{-rt} e_1(t; 0; -r).$$

Similarly, for the second moment, based on (4.13), we have

$$\ell_2(t) = \mathbb{E} \left[(L_t^0)^2 \right] = e^{-2rt} e_2(t; 0; -r),$$

with its Laplace transform

$$\mathcal{L}_\theta \{ \ell_2(t) \} = \mathcal{L}_\theta \left\{ e^{-2rt} e_2(t; 0; -r) \right\} = \hat{e}_2(\theta + 2r; 0; -r),$$

where $\hat{e}_2(\theta; S_0; \delta)$ is specified by (4.16). Then, we have the Laplace transform

$$\mathcal{L}_\theta \{ \ell_2(t) \} = \frac{\mu_2}{\gamma_1} \frac{1}{2r} \left(\frac{1}{\theta} - \frac{1}{\theta + 2r} \right) + \frac{2\mu_1^2}{\gamma_1} \frac{1}{\theta + 2r} \frac{\hat{p}(\theta + r)}{1 - \hat{p}(\theta + r)} \frac{1}{\theta},$$

which is exactly the same as the last two terms of (4.16) by replacing δ by r . Therefore, we have the second moment (4.26). Finally, it is straightforward to obtain the variance (4.26). \square

5 Numerical Illustration with Applications

To illustrate the applicability of renewal shot-noise processes and our newly-derived results, in this section, we offer four choices for modelling renewal interarrival times: 1) exponential (Exp), 2) gamma, 3) inverse Gaussian (IG), and 4) folded normal (FN) distributions. The first two examples are for actuarial application of discounted aggregate claims. Since the discounted aggregate claims L_t^0 defined by (2.1) can be alternatively interpreted as the present value of aggregate losses from a portfolio in general, we use the third and fourth examples for credit risk and reliability applications, respectively. We commonly assume $S_0 = 1$ and $\delta = r = 0.05$, and the jump sizes follow an exponential distribution of unit rate, i.e., $\mu_1 = 1, \mu_2 = 2$ for all four cases.

For each case, we compute the first conditional moments and variances of renewal shot-noise process and discounted aggregate claims, respectively. Except for the first case of exponential distribution, it is often not easy to obtain explicit expression for $F_4(t)$ in (4.22) and $\tilde{F}_4(t)$ in (4.26). We have to rely on estimating $F_4(t)$ by Monte Carlo *exact simulation* (ES)¹ via Algorithm 4.1, or, numerical inversion (NI) of Laplace transform such as *Euler algorithm* and *Talbot algorithm* (Abate and Whitt, 2006, p.415-416). The detailed implementation of NI we adopted in this paper is explained in Appendix A. However, it is well known that the algorithms for numerical inversion are not perfect, and they are often not reliable when the underlying function has some discontinuity or oscillation, or the function of Laplace transform involves complex special functions. Therefore,

¹ES is a simulation method of drawing an unbiased associated estimator throughout the entire simulation process.

efficient simulation becomes a crucial and more reliable alternative tool for estimation. Based on the fact that the shape of true function $F_4(t)$ or $\tilde{F}_4(t)$ beyond the exponential case is unknown, it is prudent for us to implement the two estimation approaches of ES and NI simultaneously in order to validate each other.

5.1 Example: Poisson Shot-noise Process

If N_t is a Poisson process, i.e., the interarrival times follow a simple exponential distribution, then, S_t is the classical Poisson shot-noise process and explicit results for variances exist. This special case was investigated by Jang (2004), and same results can be recovered here. In fact, this provides a benchmark case that can be used for validating the estimation methods ES and NI for computing $F_4(t)$ and $\tilde{F}_4(t)$ in the conditional second moments and variances. If $\tau_i \sim \text{Exp}(\varrho)$, $\varrho > 0$, with the density function

$$p(u) = \varrho e^{-\varrho u},$$

we have

$$\hat{p}(\theta) = \frac{\varrho}{\varrho + \theta}, \quad \gamma_1 = \frac{1}{\varrho}.$$

From (4.11), we have the first moment

$$\mathbb{E}[S_t | S_0] = S_0 e^{-\delta t} + \mu_1 \varrho \left(\frac{1 - e^{-\delta t}}{\delta} \right).$$

The first moment of discounted aggregate claims (i.e., the actuarial net premium) at present time 0 is given by

$$\mathbb{E}[L_t^0] = \mu_1 \varrho \left(\frac{1 - e^{-rt}}{r} \right),$$

which can be also found in L evell e and Garrido (2001a,b), Jang (2004) and Jang and Kravavych (2004). For calculating the associated variances, from (4.20), we have

$$\hat{f}_4(\theta) = 2\delta \left(\frac{1}{\theta + \delta} - \frac{1}{\theta + 2\delta} \right), \quad \hat{F}_4(\theta) = \frac{2\delta}{\theta} \left(\frac{1}{\theta + \delta} - \frac{1}{\theta + 2\delta} \right),$$

which can be inverted analytically, respectively, i.e.,

$$f_4(t) = 2\delta (e^{-\delta t} - e^{-2\delta t}), \quad F_4(t) = (1 - e^{-\delta t})^2.$$

Hence, based on (4.22) and (4.26), we have

$$\text{Var}[S_t | S_0] = \mu_2 \varrho \left(\frac{1 - e^{-2\delta t}}{2\delta} \right), \quad \text{Var}[L_t^0] = \mu_2 \varrho \left(\frac{1 - e^{-2rt}}{2r} \right).$$




Figure 2: Means and variances of S_t and L_t^0 for the exponential (Exp) case with $\varrho = 1$; the associated detailed numerical results are reported in Table 1

So, $\text{Var}[S_t | S_0]$ and $\text{Var}[L_t^0]$ are equal when $r = \delta$. We plot the conditional means, the true and estimated variances of S_t and L_t^0 for $\varrho = 1$ respectively in Figure 2, with numerical results reported in Table 1. Note that, each point in the variance plots is estimated by the exact simulation (ES) via Algorithm 4.1 with 10^7 replications. For implementing Algorithm 4.1, since an exponential distribution after the exponential tilting (4.21) is still an exponential distribution, i.e.,

$$\hat{f}_{E_i}(\theta) = \frac{\hat{p}(\theta + \delta)}{\hat{p}(\delta)} = \frac{\frac{\varrho}{\varrho + \theta + \delta}}{\frac{\varrho}{\varrho + \delta}} = \frac{\varrho + \delta}{(\varrho + \delta) + \theta},$$

we have $E_i \sim \text{Exp}(\varrho + \delta)$ which can be sampled directly.

5.2 Example: Gamma Shot-noise Process

For a Gamma distribution (including chi-squared distribution as a special case), i.e., $\tau_i \sim \Gamma(\alpha, \beta)$ with density function

$$p(u) = \frac{\beta^\alpha}{\Gamma(\alpha)} u^{\alpha-1} e^{-\beta u}, \quad u > 0,$$

where $\alpha, \beta > 0$ are the shape and rate parameters, respectively, we have the mean is $\gamma_1 = \alpha/\beta$ and the Laplace transform

$$\hat{p}(\theta) = \left(\frac{\beta}{\beta + \theta} \right)^\alpha.$$

With the exact simulation via Algorithm 4.1 and using the parameter setting $(\alpha, \beta) = (2, 2)$, the associated conditional means and variances are plotted in Figure 3 and reported in Table 1. Each point in the variance plots is estimated by the exact simulation (ES) via Algorithm 4.1 with 10^7

Table 1: Means and variances of S_t and L_t^0

Time t	$\mathbb{E}[S_t S_0]$	$\mathbb{E}[L_t^0]$	NI-Var	ES-Var	True-Var	$\mathbb{E}[S_t S_0]$	$\mathbb{E}[L_t^0]$	NI-Var	ES-Var
	<Exp>					<Gamma>			
0.2	1.1891	0.1990	0.3960	0.3942	0.3960	1.1891	0.1990	0.3652	0.3655
0.4	1.3762	0.3960	0.7842	0.7843	0.7842	1.3762	0.3960	0.6860	0.6849
0.6	1.5615	0.5911	1.1647	1.1643	1.1647	1.5615	0.5911	0.9839	0.9790
0.8	1.7450	0.7842	1.5377	1.5411	1.5377	1.7450	0.7842	1.2685	1.2591
1.0	1.9266	0.9754	1.9033	1.9107	1.9033	1.9266	0.9754	1.5443	1.5322
1.2	2.1065	1.1647	2.2616	2.2668	2.2616	2.1065	1.1647	1.8131	1.8044
1.4	2.2845	1.3521	2.6128	2.6110	2.6128	2.2845	1.3521	2.0759	2.0649
1.6	2.4608	1.5377	2.9571	2.9485	2.9571	2.4608	1.5377	2.3333	2.3222
1.8	2.6353	1.7214	3.2946	3.2861	3.2946	2.6353	1.7214	2.5855	2.5695
2.0	2.8081	1.9033	3.6254	3.6152	3.6254	2.8081	1.9033	2.8325	2.8209
	<IG>					<FN>			
0.2	1.1891	0.1990	0.3621	0.3623		2.2371	1.2471	2.1387	2.1371
0.4	1.3762	0.3960	0.6992	0.7012		3.4619	2.4817	4.0478	4.0477
0.6	1.5615	0.5911	1.0390	1.0395		4.6745	3.7041	5.9246	5.8159
0.8	1.7450	0.7842	1.3805	1.3840		5.8751	4.9143	7.7669	7.6319
1.0	1.9266	0.9754	1.7215	1.7283		7.0637	6.1125	9.5747	9.6682
1.2	2.1065	1.1647	2.0605	2.0664		8.2405	7.2987	11.3487	11.5322
1.4	2.2845	1.3521	2.3965	2.4025		9.4056	8.4732	13.0894	13.4175
1.6	2.4608	1.5377	2.7288	2.7363		10.5591	9.6359	14.7973	14.8805
1.8	2.6353	1.7214	3.0568	3.0611		11.7011	10.7871	16.4730	16.5891
2.0	2.8081	1.9033	3.3802	3.3764		12.8317	11.9269	18.1169	18.3422

replications. For implementing Algorithm 4.1, since a gamma distribution after the exponential tilting (4.21) is still a gamma distribution, i.e.,

$$\hat{f}_{E_i}(\theta) = \frac{\hat{p}(\theta + \delta)}{\hat{p}(\delta)} = \frac{\left(\frac{\beta}{\beta + \theta + \delta}\right)^\alpha}{\left(\frac{\beta}{\beta + \delta}\right)^\alpha} = \left(\frac{\beta + \delta}{\beta + \delta + \theta}\right)^\alpha,$$

we have $E_i \sim \Gamma(\alpha, \beta + \delta)$ which can be sampled directly.

5.3 Example: Inverse Gaussian Shot-noise Process

We can also make an immediate application to modelling credit defaults: N_t can be used for modelling the arrivals of credit defaults (of e.g. corporate bonds) in a large credit portfolio, X_i is the loss of the i^{th} credit default, then, L_t^0 can be interpreted as the present value of the total loss of this credit portfolio within the time period of $[0, t]$. Our results for the moments of L_t^0 provided in Section 4.2. tell people the moments of the portfolio loss, which could be useful for credit risk management and measurement. U_t is the time elapsed since the last default occurred². For numerical illustration, we assume the interarrival times between two successive credit defaults follow an

²Since the 2007 financial crisis, default rates of corporate bonds have decreased, as the world economy has emerged from the global financial crisis (GFC) with improving market conditions. However, default rates going forward are dependent on the progress of world economic recovery and growth, as well as oil and commodity prices, fiscal and monetary policy and interest rate fluctuations. Hence, in specific situations like 2007-2008 GFC, the time elapsed since the last default occurred could be an important parameter in credit default modelling. The properties of renewal shot-noise processes and the results newly found in this paper could be also appropriate for modelling credit risk.




Figure 3: Means and variances of S_t and L_t^0 for the gamma case with $(\alpha, \beta) = (2, 2)$; the associated detailed numerical results are reported in Table 1

inverse Gaussian, i.e., $\tau_i \sim \text{IG}(a, b)$ with density function

$$p(u) = \frac{a}{\sqrt{2\pi u^3}} e^{-\frac{(a-bu)^2}{2u}}, \quad u, a, b > 0,$$

where the mean is $\gamma_1 = a/b$ and the shape parameter is a^2 , we have the Laplace transform

$$\hat{p}(\theta) = e^{-[\sqrt{2\theta+b^2}-b]a}.$$

With the exact simulation via Algorithm 4.1 and using the parameter setting $(a, b) = (1, 1)$, the conditional means and variances of S_t and the present value of credit portfolio losses L_t^0 are plotted in Figure 4 and reported in Table 1. Each point in the variance plots is estimated by the exact simulation (ES) via Algorithm 4.1 with 10^7 replications. For implementing Algorithm 4.1, since an IG distribution after the exponential tilting (4.21) is still an IG distribution, i.e.,

$$\hat{f}_{E_i}(\theta) = \frac{\hat{p}(\theta + \delta)}{\hat{p}(\delta)} = \frac{e^{-[\sqrt{2(\theta+\delta)+b^2}-b]a}}{e^{-[\sqrt{2\delta+b^2}-b]a}} = e^{-[\sqrt{2\theta+(2\delta+b^2)}-\sqrt{2\delta+b^2}]a},$$

we have $E_i \sim \text{IG}(a, \sqrt{2\delta + b^2})$ which can be sampled directly.

5.4 Example: Folded Normal Shot-noise Process

Queues and related models are important in solving many complex reliability problems. The renewal shot-noise process and its variations can be considered to deal with the expected busy periods




Figure 4: Means and variances of S_t and L_t^0 for the inverse Gaussian (IG) case with $(a, b) = (1, 1)$; the associated detailed numerical results are reported in Table 1

in terms of queuing system and the virtual waiting times of customers, etc. Due to the similar nature of cashflow structure, our results could also be applied to reliability modelling. N_t accounts the total number of failures of machine components up to time t . X_i is the individual cost of the i^{th} failure, then, L_t^0 can be interpreted as the present value of the total cost within the time period of $[0, t]$. For reliability modelling, we take the *folded normal* distribution (including the *half-normal* distribution as a special case) as an example for modelling the interarrival times of two successive failures, i.e., $\tau_i \sim \text{FN}(\mu, \sigma)$ with density function

$$p(u) = \frac{1}{\sqrt{2\pi\sigma^2}} \left[e^{-\frac{(u-\mu)^2}{2\sigma^2}} + e^{-\frac{(u+\mu)^2}{2\sigma^2}} \right] \quad u \geq 0,$$

where $\mu, \sigma > 0$, we have the mean

$$\gamma_1 = \sqrt{\frac{2}{\pi}} \sigma e^{-\frac{\mu^2}{2\sigma^2}} + \mu \left[1 - 2\Phi\left(-\frac{\mu}{\sigma}\right) \right],$$

and the Laplace transform

$$\hat{p}(\theta) = e^{\frac{\sigma^2}{2}\theta^2 - \mu\theta} \left[1 - \Phi\left(-\frac{\mu}{\sigma} + \sigma\theta\right) \right] + e^{\frac{\sigma^2}{2}\theta^2 + \mu\theta} \left[1 - \Phi\left(\frac{\mu}{\sigma} + \sigma\theta\right) \right]. \quad (5.1)$$

Each point in the variance plots is estimated by the exact simulation (ES) via Algorithm 4.1 with 10^7 replications. For implementing Algorithm 4.1, since an folded normal distribution after the exponential tilting (4.21) is unknown, we need the *acceptance/rejection (A/R) scheme* of Algorithm




Figure 5: Means and variances of S_t and L_t^0 for the folded normal (FN) case with $(\mu, \sigma) = (0, 0.2)$; the associated detailed numerical results are reported in Table 1

B.1, where $\tau_i \sim \text{FN}(\mu, \sigma)$ in Step 1 can be simply sampled via

$$\tau_i \stackrel{\mathcal{D}}{=} |\mu + \sigma V|, \quad V \sim \mathcal{N}(0, 1).$$

With the exact simulation via Algorithm 4.1 and using the parameter setting $(\mu, \sigma) = (0, 0.2)$, the conditional means and variances of S_t and the present value of total cost L_t^0 are plotted in Figure 5 and reported in Table 1.

6 Concluding Remarks

We have mainly studied the Laplace transforms of the conditional and asymptotic moments for renewal shot-noise processes and discounted aggregate claims. A very efficient and general simulation algorithm has been developed for estimating the second conditional moments, and it has been compared with the alternative method of numerical inversion. For applications to the net premium calculation in insurance as well as credit risk and reliability modelling, the first conditional moments and variances for four different distributions of interarrival times have been computed, respectively. In fact, renewal shot-noise processes and the properties found in this paper could be also applicable to a wide range of other fields such as queueing, financial transaction data, computer networks, inventories and storage system, and etc., and we leave them as further research.

Acknowledgments

The authors would like to thank all reviewers for very helpful and constructive comments and suggestions. They also thank Yan Qu for helping the implementation of numerical inversion of Laplace transforms. The corresponding author Hongbiao Zhao would like to thank Yao Xiao for a very careful read and acknowledges the financial support from the *National Natural Science Foundation of China* (#71401147,#91546202) and the research fund provided by Shanghai University of Finance and Economics.

Appendices

A Numerical Inversion of Laplace Transform

There are numerous different schemes available in the literature for Laplace numerical inversions. The aim of using numerical inversion in this paper is for comparing and validating our newly-developed exact simulation algorithms. As finding optimal schemes for numerical inversions is not our main focus here at the current stage, we thereby adopt conventional ones. For the first three cases in Section 5, i.e., exponential, gamma and inverse Gaussian cases, which are easier, we simply apply the classical *Talbot algorithm* (Abate and Whitt, 2006, Section 6, p.416). It works very well by using the existing package of MatLab codes `euler_inversion_sym.m` available at MathWorks. For the folded normal case, which is more complicated due to the special function $\Phi(\cdot)$ in the Laplace transform $\hat{p}(\theta)$ specified in (5.1), we develop our own codes based on the *Euler algorithm* (Abate and Whitt, 2006, Section 5, p.415-416) with the aid of existing C++ package `RcppFaddeeva` from CRAN that can deal with the function $\Phi(\cdot)$ for complex values. Both algorithms involve *tuning (or scaling) parameters*, which are simple deterministic functions of positive integer M , the number of terms for approximating the infinite summation, that controls the associated truncation errors. More precisely, as illustrated in Abate and Whitt (2006), for a given Laplace transform \hat{f} of a function f , i.e.,

$$\hat{f}(s) \equiv \int_0^{\infty} e^{-st} f(t) dt,$$

the underlying function f can be approximated by

$$f(t) \approx f_n(t) \equiv \frac{1}{t} \sum_{k=0}^n \omega_k \hat{f}\left(\frac{\alpha_k}{t}\right), \quad t > 0, \quad (\text{A.1})$$

where the *nodes* α_k and *weights* ω_k are the associated *tuning (or scaling) parameters*, which are complex numbers, and depend on n but not on the transform \hat{f} or the time argument t . These tuning parameters are specified differently by Talbot algorithm and Euler algorithm as follows:

- For the *Talbot algorithm* (Abate and Whitt, 2006, Section 6, p.416), the parameters in the framework

(A.1) are $n = M$, $\alpha_k = \delta_k$ and $\omega_k = \frac{2}{5}\gamma_k$, where

$$\begin{aligned}\delta_0 &= \frac{2}{5}M, & \delta_k &= \frac{2}{5}k\pi \left[\cos\left(\frac{k\pi}{M}\right) + i \right], & 0 < k < M, \\ \gamma_0 &= \frac{1}{2}e^{\delta_0}, & \gamma_k &= \left[1 + i\frac{k\pi}{M} \left(1 + \cot^2\left(\frac{k\pi}{M}\right) \right) - i \cot\left(\frac{k\pi}{M}\right) \right] e^{\delta_k}, & 0 < k < M,\end{aligned}$$

with $i \equiv \sqrt{-1}$.

- For the *Euler algorithm* (Abate and Whitt, 2006, Section 5, p.415-416), the parameters in the framework (A.1) are $n = 2M$, $\alpha_k = \beta_k$ and $\omega_k = 10^{\frac{M}{3}}\eta_k$, where

$$\begin{aligned}\beta_k &= \frac{\ln 10}{3}M + \pi ik, & \eta_k &\equiv (-1)^k \xi_k, \\ \xi_0 &= \frac{1}{2}, & \xi_k &= 1, \quad 1 \leq k \leq M, & \xi_{2M} &= \frac{1}{2^M}, \\ \xi_{2M-k} &= \xi_{2M-k+1} + 2^{-M} \binom{M}{k}, & & & 0 < k < M.\end{aligned}$$

In our paper, we adopt $M = 64$ for the first three cases and $M = 6$ for the folded normal case, which determine the associated tuning parameters, respectively. From extensive numerical experiments, we do find that the estimation results from numerical inversion algorithms could be unstable for some sets of parameters. Indeed, it is a very typical problem for the numerical inversion approach, and it is why we shall advocate using our newly-developed exact simulation approach as an alternative.

B Random Variate Generator for E_i

Algorithm B.1 (A/R Scheme for E_i). *For exactly sampling one random variable E_i in general:*

1. Generate a random variable E_e with density $p(u)$;
2. Generate a uniformly distributed random variable $V \sim \mathcal{U}[0, 1]$;
3. If $V \leq e^{-\delta E_e}$, then, accept and set $E_i = E_e$; otherwise, reject and go back to Step 1.

Proof. To exactly sample a random variable E_i with the density function (4.21), we adopt the acceptance/rejection (A/R) scheme with the envelop density function $f_{E_e}(t) = p(t)$. Then, we can easily find the smallest possible constant K such that

$$\frac{f_{E_i}(t)}{f_{E_e}(t)} = \frac{e^{-\delta t}}{\hat{p}(\delta)} \leq K.$$

Hence, we have $K = 1/\hat{p}(\delta)$, and the acceptance level

$$\frac{f_{E_i}(t)}{Kp(t)} = e^{-\delta t}.$$

Note that, the probability of acceptance is $1/\hat{p}(\delta)$. □

References

- Abate, J. and Whitt, W. (1992). The Fourier-series method for inverting transforms of probability distributions. *Queueing Systems*, 10(1):5–87.
- Abate, J. and Whitt, W. (1995). Numerical inversion of Laplace transforms of probability distributions. *ORSA Journal on Computing*, 7(1):36–43.
- Abate, J. and Whitt, W. (2006). A unified framework for numerically inverting Laplace transforms. *INFORMS Journal on Computing*, 18(4):408–421.
- Albrecher, H. and Asmussen, S. (2006). Ruin probabilities and aggregate claims distributions for shot noise Cox processes. *Scandinavian Actuarial Journal*, 2006(2):86–110.
- Andersen, E. S. (1957). On the collective theory of risk in case of contagion between claims. *Bulletin of the Institute of Mathematics and its Applications*, 12:275–279.
- Beard, R. E., Pesonen, E., and Pentikäinen, T. (1984). *Risk Theory*. Springer.
- Brémaud, P. (2000). An insensitivity property of Lundberg’s estimate for delayed claims. *Journal of Applied Probability*, 37(3):914–917.
- Campbell, N. (1909a). Discontinuities in light emission. *Proceedings of the Cambridge Philosophical Society*, 15:310–328.
- Campbell, N. (1909b). The study of discontinuous phenomena. *Proceedings of the Cambridge Philosophical Society*, 15:117–136.
- Cox, D. R. (1955). The analysis of non-Markovian stochastic processes by the inclusion of supplementary variables. *Mathematical Proceedings of the Cambridge Philosophical Society*, 51(3):433–441.
- Cox, D. R. (1962). *Renewal Theory*. Methuen, London.
- Cox, D. R. and Isham, V. (1980). *Point Processes*. Chapman & Hall, London.
- Cox, D. R. and Miller, H. D. (1965). *The Theory of Stochastic Processes*. CRC Press.
- Dassios, A. and Embrechts, P. (1989). Martingales and insurance risk. *Stochastic Models*, 5(2):181–217.
- Dassios, A. and Jang, J. (2003). Pricing of catastrophe reinsurance and derivatives using the Cox process with shot noise intensity. *Finance and Stochastics*, 7(1):73–95.
- Dassios, A. and Jang, J. (2005). Kalman-Bucy filtering for linear systems driven by the Cox process with shot noise intensity and its application to the pricing of reinsurance contracts. *Journal of Applied Probability*, 42(1):93–107.
- Dassios, A., Jang, J., and Zhao, H. (2015). A risk model with renewal shot-noise Cox process. *Insurance: Mathematics and Economics*, 65:55–65.

- Dassios, A. and Zhao, H. (2011). A dynamic contagion process. *Advances in Applied Probability*, 43(3):814–846.
- Dassios, A. and Zhao, H. (2012). Ruin by dynamic contagion claims. *Insurance: Mathematics and Economics*, 51(1):93–106.
- Dassios, A. and Zhao, H. (2014). A Markov chain model for contagion. *Risks*, 2(4):434–455.
- Davis, M. H. (1984). Piecewise-deterministic Markov processes: A general class of non-diffusion stochastic models. *Journal of the Royal Statistical Society. Series B (Methodological)*, 46(3):353–388.
- Davis, M. H. (1993). *Markov Models and Optimization*. Chapman & Hall/CRC.
- Duffie, D., Filipovic, D., and Schachermayer, W. (2003). Affine processes and applications in finance. *Annals of Applied Probability*, 13(3):984–1053.
- Duffie, D., Pan, J., and Singleton, K. (2000). Transform analysis and asset pricing for affine jump-diffusions. *Econometrica*, 68(6):1343–1376.
- Gerber, H. U. and Shiu, E. S. (1994). Option pricing by Esscher transforms. *Transactions of the Society of Actuaries*, 46:99–140.
- Gerber, H. U. and Shiu, E. S. (1998). On the time value of ruin. *North American Actuarial Journal*, 2(1):48–72.
- Gerber, H. U. and Shiu, E. S. (2005). The time value of ruin in a Sparre Andersen model. *North American Actuarial Journal*, 9(2):49–69.
- Goovaerts, M. J., Laeven, R. J. A., and Shang, Z. (2012). Transform analysis and asset pricing for diffusion processes: a recursive approach. *Journal of Computational Finance*, 16(1):47–81.
- Grandell, J. (1991). *Aspects of Risk Theory*. Springer.
- Jang, J. (2004). Martingale approach for moments of discounted aggregate claims. *Journal of Risk and Insurance*, 71(2):201–211.
- Jang, J. and Kravavych, Y. (2004). Arbitrage-free premium calculation for extreme losses using the shot noise process and the Esscher transform. *Insurance: Mathematics and Economics*, 35(1):97–111.
- Karatzas, I. and Shreve, S. (1991). *Brownian Motion and Stochastic Calculus*. Springer.
- Karlin, S. and Taylor, H. (1975). *A First Course in Stochastic Processes*. Academic Press, New York.
- Klüppelberg, C. and Mikosch, T. (1995). Explosive Poisson shot noise processes with applications to risk reserves. *Bernoulli*, 1(1/2):125–147.
- Léveillé, G. and Garrido, J. (2001a). Moments of compound renewal sums with discounted claims. *Insurance: Mathematics and Economics*, 28(2):217–231.

- Léveillé, G. and Garrido, J. (2001b). Recursive moments of compound renewal sums with discounted claims. *Scandinavian Actuarial Journal*, 2001(2):98–110.
- Léveillé, G., Garrido, J., and Wang, Y. (2010). Moment generating functions of compound renewal sums with discounted claims. *Scandinavian Actuarial Journal*, 2010(3):165–184.
- Li, S. and Garrido, J. (2005). On a general class of renewal risk process: analysis of the Gerber-Shiu function. *Advances in Applied Probability*, 37(3):836–856.
- Linetsky, V. (1997). The path integral approach to financial modeling and options pricing. *Computational Economics*, 11(1):129–163.
- Linetsky, V. (2004). The spectral decomposition of the option value. *International Journal of Theoretical and Applied Finance*, 7(3):337–384.
- Linetsky, V. (2007). Spectral methods in derivatives pricing. In Birge, J. R. and Linetsky, V., editors, *Financial Engineering*, Handbooks in Operations Research and Management Science, pages 223–299. Elsevier.
- Macci, C. and Torrisi, G. L. (2011). Risk processes with shot noise Cox claim number process and reserve dependent premium rate. *Insurance: Mathematics and Economics*, 48(1):134–145.
- Møller, J. and Torrisi, G. L. (2005). Generalised shot noise Cox processes. *Advances in Applied Probability*, 37(1):48–74.
- Picinbono, B., Bendjaballah, C., and Pouget, J. (1970). Photoelectron shot noise. *Journal of Mathematical Physics*, 11(7):2166–2176.
- Rice, J. (1977). On generalized shot noise. *Advances in Applied Probability*, 9(3):553–565.
- Rolski, T., Schmidli, H., Schmidt, V., and Teugels, J. (2008). *Stochastic Processes for Insurance and Finance*. Wiley.
- Ross, S. M. (1996). *Stochastic Processes*. Wiley, New York.
- Schmidt, T. (2014). Catastrophe insurance modeled by shot-noise processes. *Risks*, 2(1):3–24.
- Schottky, W. (1918). Über spontane Stromschwankungen in verschiedenen Elektrizitätsleitern. *Annalen der Physik*, 362(23):541–567.
- Seal, H. L. (1983). The Poisson process: its failure in risk theory. *Insurance: Mathematics and Economics*, 2(4):287–288.
- Shang, Z., Goovaerts, M., and Dhaene, J. (2011). A recursive approach to mortality-linked derivative pricing. *Insurance: Mathematics and Economics*, 49(2):240–248.
- Smith, J. A. (1980). *Point Process Models of Rainfall*. PhD thesis, The Johns Hopkins University.

- Torrasi, G. L. (2004). Simulating the ruin probability of risk processes with delay in claim settlement. *Stochastic Processes and their Applications*, 112(2):225–244.
- Verveen, A. and DeFelice, L. (1974). Membrane noise. *Progress in Biophysics and Molecular Biology*, 28:189–234.
- Willmot, G. E. (2007). On the discounted penalty function in the renewal risk model with general interclaim times. *Insurance: Mathematics and Economics*, 41(1):17–31.
- Willmot, G. E. and Dickson, D. (2003). The Gerber–Shiu discounted penalty function in the stationary renewal risk model. *Insurance: Mathematics and Economics*, 32(3):403–411.
- Woo, J.-K. (2010). Some remarks on delayed renewal risk models. *Astin Bulletin*, 40(1):199–219.
- Zhu, L. (2013). Ruin probabilities for risk processes with non-stationary arrivals and subexponential claims. *Insurance: Mathematics and Economics*, 53(3):544–550.